bony core; FIG. 15, also in front of cartilaginous phalanx, shows oval-shaped bony core in connection with bony ridge, which lies in line with top of phalanx; FIG. 16, shows cartilaginous tip of phalanx invested by bony sheath, lying in contact with irregular mass of bone under phalanx; FIG. 17, section some distance from tip showing bony sheath, with irregular wing-like expansions, investing calcified cartilage of terminal phalanx—on the left side the section has passed through one of the many channels that open on the upper surface; FIG. 18, section in front of the large gap seen in FIGS. 12 and 13, showing the great size of the “wing;” FIG. 19, a still more proximal section on a level with the point where the buttress springs from the investing cap; FIG. 20, through the buttress and the tip of the “wing”—note that the phalanx is still cartilaginous and no longer provided with a complete bony investment; FIGS. 12 and 13 are thirty-two times natural size; FIGS. 14-20, are about fifty times enlarged.

FIGS. 21-25 represent fore-limb of embryo D (35 cm.); FIG. 21, natural size skeleton of left fore-limb from outer or ulnar aspect. The shafts of the humerus (*h*), radius (*r*), ulna (*u*), metacarpals II., III., and IV., and of the first phalanx are now well ossified as compared with embryo C. A small centre is present in the second phalanx and the middle portion of the terminal phalanx (FIG. 25) is also ossified, while the investing cap is well formed, and in some respects resembles the outer portion of the os pedis of the adult. In FIG. 21 the distal end of the ulna (*u*) is seen to articulate freely with the cuneiform (*c*). FIG. 22, a front view, shows metacarpals II. and IV., lying to a considerable extent under cover of metacarpal III.; FIG. 23 represents a front view of the terminal phalanx and its bony cap; FIG. 24 represents the same phalanx as seen from below; and FIG. 25 as seen in section. In FIG. 24 the extent of the ossification of the phalanx proper is indicated by the dark part (*c:c*) as seen through the bony cap; in FIG. 25 the difference between the cap and the ossifying phalanx is easily made out. In the section which FIG. 25 represents, the anterior third of the phalanx, with the exception of the very tip, was comparatively soft, while the part between the anterior third and the partly calcified cartilage was fairly well ossified.

VERMINOUS PNEUMONIA IN THE SHEEP.

By J. M'FADYEAN, M.B., B.Sc., Royal Veterinary College, London.

CONSIDERING the amount of attention that has been devoted to the study of the animal parasites of our domesticated quadrupeds, it is remarkable that authors should be so much at variance regarding the precise species of nematodes commonly found in the lung of the sheep. In order to show the diversity of opinion on this subject expressed by different authors, I shall here quote from a few of the standard text-books in English, French, and German.

Professor Williams, in his work on Veterinary Medicine (sixth edition), says: “In the calf and lamb, bronchial irritation arising from the presence of nematode parasites, termed strongyles, is of frequent occurrence, and is variously termed *The Husk*, *Hoose*, *Phthisis pulmonalis verminalis*, and parasitic bronchitis. In the lamb, the parasite, termed *Strongylus filaria*, is from 1 to 1½ inches long. . . . The anatomical situation of the parasites in the lungs of lambs and sheep is not always the same. In lambs they are found not only in the bronchial tubes, but also in the lung substance, whilst in sheep they are generally encysted in the parenchyma of the lungs, giving

them the appearance of being filled with small tubercular deposits ; indeed the disease was for a long time looked upon as a true tubercular affection . . . The lungs of sheep, which have been the hosts of these parasites, are thickly covered with numerous small nodules, varying in size from a pin-head to a hemp seed, or even larger, and resembling small vesicles or blisters. Some are filled with a clear fluid ; others contain a soft material, consisting of granular matter ; whilst others are hard and gritty—and all contain minute worms coiled upon themselves . . . The presence of these parasites in the lung does not always cause irritation or inconvenience to the host. The lungs of sheep killed in the prime condition are found loaded with them. Occasionally, however, they induce debility, anæmia, and cause death, more particularly in lambing ewes, at or about the period of parturition. Many ewes died from this cause during the lambing season¹ of 1874. In lambs, however, the parasites find their way through the softer lung structures into the bronchial tubes, and there give rise to irritation, and to the symptoms of the “lamb disease.”

It will be gathered from the preceding that Professor Williams has met with only one nematode in the lung of the sheep, viz., the *Strongylus filaria*, and that he holds this parasite accountable for both the bronchitic “lamb disease,” and for the formation of the tubercle-like nodules in the lungs of adult sheep.

The late Dr Cobbold, in his treatise on “The Entozoa of Man and Animals,” published in 1879, says : “The nematodes of the ruminants are both numerous in, and destructive to, their bearers, those infesting the lungs being productive of a parasitic bronchitis, termed husk or hoose. In cattle the lung worm (*Strongylus micrurus*) is particularly fatal to calves, whilst the *S. filaria* attacks sheep, and especially lambs, a larger but less common lung strongyle (*S. rufescens*) is sometimes found associated with the latter.”

In his lectures delivered at the Royal Veterinary College, Dr Cobbold, after describing the *Strongylus filaria*, thus refers to the *S. rufescens* :—“*S. rufescens*, the long worm, or gordian worm. Head unarmed, mouth has three lobes. Vulva immediately in front of the anus. Caudal bursa short, and has a few short rays. Spicules strongly curved. Eggs contain a segmented yolk. Length 6-7 inches. Sometimes found in the lung structure.”¹

Dr Cobbold’s views regarding the *Strongylus rufescens* are stated at greater length in the following passages, which are quoted by Professor Simmonds in a “Report on the Parasitic Lung Disease of Lambs”² :—

“The well-known lamb disease, though generally supposed to be due to the presence of a single species of worm, called the common lung-strongyle, or *Strongylus filaria*, is in reality due to the occurrence of at least two different parasites belonging to the same group.

“For years past it has been known to continental helminthologists that a second species of nematode worm is usually associated with the above-named parasite, the form in question having been originally described by Professor Leuckart, under the title of *Strongylus rufescens*.

¹ For this quotation I am indebted to my colleague Professor Penberthy, who has kindly placed at my disposal his notes of Dr Cobbold’s Course of Lectures delivered during the Session 1880-81.

² Journal of the Royal Agricultural Society of England, vol. x. S.S. Part 1.

"I am the more desirous of calling attention to this fact, since Dr Crisp, an earnest worker in the cause of helminthology, believes that he has discovered a new worm in the lungs of lambs and sheep, which he calls a *gordius*. I entertain no doubt that Dr Crisp's gordian worm is the adult representative of the immature worm found by Professor Brown some years ago; fresh specimens have also recently been examined by Mr Axe and myself. This so-called gordian worm is a large species of strongyle, which, in contra-distinction to the common lung strongyle, I am in the habit of speaking of as the long strongyle, and I think this description is particularly convenient, because the parasite, in its full-grown state, acquires a length of from 6 to 7 inches, or, as Leuckart affirms, it may be a span long."

It thus appears that Dr Cobbold recognised two species of strongyles as being the cause of lung disease in sheep, viz., the *Strongylus filaria*, and the *S. rufescens*, or long strongyle, and it is important to note that the parasite to which he applied these latter designations was 6 or 7 inches long. Unfortunately, I have not been able to gather from Dr Cobbold's writings whether he considered that these two species caused the same or different lesions in the lungs of affected sheep, but, from his reference in one of the passages quoted to Professor Brown's discovery of an immature form of the long strongyle, it is probable that he ascribed the pseudo-tubercles found in the sheep's lung to this parasite. It may be noted, however, that Professor Brown himself did not regard the worm whose discovery is here referred to as belonging to a new species, as will be seen from the following quotations from Harley and Brown's "Demonstrations of Microscopic Anatomy" (second edition, 1876):—

"*Filaria bronchialis*, or *Strongylus bronchialis*, is found in the bronchial tubes of man, and in the lungs of sheep. When accumulated in vast numbers, they are capable of causing suffocation.

"In the lungs of the sheep a variety of this worm is frequently found occupying whitish tuberculous-looking spots, commonly mistaken for tubercle (from which the lungs of the sheep appear to be perfectly exempt). The young of the parasite migrate to the small bronchial tubes, and occasion considerable irritation."

This reference to the parasite enclosed in the pseudo-tubercles is supplemented by a drawing showing the worm coiled up "amidst a quantity of exudation matter, undergoing calcareous degeneration." According to this view, then, the parasite which inhabits the tubercle-like nodules in the sheep's lung belongs to the species *Strongylus filaria*, and not, as Cobbold believed, to the *S. rufescens*, or long strongyle.

Dr Cooper Curtice, in his work on "The Animal Parasites of Sheep," published in 1890, thus summarises his opinions regarding the causes of verminous pneumonia and verminous bronchitis among American sheep.

"The lung worms which cause disease in sheep in the United States belong to two, perhaps three, different species. They are *Strongylus ovis-pulmonalis*, Diesing, the hair-lung worm; and *Strongylus filaria*, Rud., the thread-lung worm. The third species, which has been reported as infesting sheep in Europe, is the hog-lung worm, *Strongylus paradoxus*, an abundant species occurring in the lungs of swine in this country, and while it has never been credited as having been found in our sheep it is to be looked for. As its size and the

disease it causes is similar to that of *Strongylus filaria*, it will not be treated separately.

"The diseases produced by these species of worms are caused by the mechanical injuries the worms inflict on the delicate membranes of the lungs and the clogging up of the air passages by them and the *débris* which they produce. The two forms of disease produced depend on the different size and habits of the two species. *Strongylus ovis-pulmonalis*, being very small, penetrates the air passages to their endings in the bronchioles and infundibuli, and causes disease in them primarily, while *Strongylus filaria*, which penetrates only into the bronchi, creates a disturbance there which produces a solidification of the lung secondarily. The general diagnosis of each disease is no easy matter. The disease produced by *Strongylus ovis-pulmonalis* is characterised by the spongy feeling of the lung and the presence of nodules from the size of a mustard seed to that of a pea scattered under the surface of the dorsum of the lung and at its posterior free edge. These may be connected by a greyish, fleshy, intermediate portion of the lung into patches of considerable size. The parasites can scarcely be seen by the unaided eye; but if small pieces of the affected lung or a tubercle be placed in a shallow dish of water and teased out with needles under a tripod lens, they can be readily seen.

"The disease produced by *Strongylus filaria* and *S. paradoxus* is characterised by the posterior portion usually, or some entire section of the affected lung appearing as a solid, usually red, mass which has lost all of its contained air and is in a state of hepatisation.

"A piece cut out generally sinks in water, while pieces from the former disease float. If in the latter disease the trachea is carefully slit open and the branches traced down to the affected part the parasites will be found in thread-like bunches, completely filling the tubes."

At an earlier part of the same work the author says: "There are two lung parasites of sheep in the Old World that have not been found here, viz.: *Strongylus rufescens*, Leuckart, and *S. minutissimus*, Megnin. The former is so meagrely described that it is probably not a distinct species, but a synonym of *Strongylus* (*Pseudalius*) *ovis-pulmonalis*, Diesing. The latter occurs in Algeria, is well described and figured, and seems to be a well-established species."

Dr Cooper Curtice thus recognises two lung strongyles of the sheep, viz., the *Strongylus filaria*, the cause of verminous bronchitis in lambs, and the *S. ovis-pulmonalis*, which determines the formation of nodular or tubercular lesions in the lungs of adult sheep. He describes the encysted worms, as being sexually mature and full-grown, the length of the male being given as 16 mm., and that of the female as 25 mm. "Escaping from the cysts they make their way into the small air-tubes (bronchioles and bronchi), where the sexes mate and reproduce. The eggs are then laid in surrounding cavities and hatched into young worms, which make their way into the neighbouring air-chambers (infundibula). Afterwards some of these worms may be coughed out of the lungs on to the pastures and infect other sheep." The embryo worm is said to be provided with a very sharp-pointed tail.

The quotations already made will suffice to show the diversity of opinion regarding the etiology of parasitic lung disease of the sheep

on the part of English authors. Let us now see what are the views of German and French authors on the same subject.

In Dr F. A. Zürn's work on the parasites of the domesticated animals (*Die Schmarotzer auf und in dem Körper unserer Haus-äugetiere*, 1882) the lung-worm disease of sheep is ascribed to the *Strongylus filaria*, the main habitat of which is said to be the trachea and bronchi, but adult worms and still more frequently eggs and embryos of the same species are said to be met with in the lung parenchyma, under the pleura. It is not clear whether the latter remark refers to the worm-tubercles of the sheep's lung, but in a note to the article describing the *Strongylus filaria* reference is made to the observations of Utz and Lydtin regarding the lung-worm nodular disease (Lungenwurmknotenkrankheit) of the sheep. Utz found in the lungs of sheep, under the *pleura pulmonalis*, nodules varying in size from a pin's head to a pea, and each containing intricately coiled nematodes. His observations were published in the *Thierärztliche Mitteilungen* for 1880, and in the same year Lydtin wrote that similar worm nodules had been examined by him and Fenzling. The nodules were described by Lydtin as varying in size from a pin's head to a lentil, dark-violet, reddish-brown, or yellow in colour, rather firm in consistence, merging gradually into the surrounding lung tissue, and generally covered by thickened and opaque pleura. At the periphery of the nodules there was a small cavity containing a thread-like round worm about 5·6 mm. long and 0·17 mm. thick. The body of the worm was somewhat thinner in front than behind, the mouth was naked, and succeeded by a small muscular œsophagus. Only females and embryos had been met with, and the latter had a characteristic lancet-shaped caudal process. The worm was thought by Lydtin to be probably identical with the *nematoideum ovis-pulmonalis*, first described by Sandie and Padley in 1849, but Zürn, "as the result of his own investigations," holds that it represents the embryonic stage of the *Strongylus filaria*.

In 1883, Koch, an Austrian veterinary surgeon, published a monograph entitled "Die Nematoden der Schaflunge," in which he held that three (or with the *Strongylus rufescens* of Leuckart, four) different parasites are responsible for the lesions of verminous bronchitis and pneumonia in sheep, viz., the *Strongylus filaria*, the *Strongylus paradoxus*, and a third parasite which he proposed to name the *Pseudalius ovis-pulmonalis*. Koch's memoir adds nothing to what was previously known regarding the first two of these parasites, and the greater part of it is taken up with a description of the so-called *Pseudalius ovis-pulmonalis* and the lung lesions excited by it. These lesions, Koch says, vary according to the number of parasites in the affected lung, but are always characteristic. One observes at various places on the surface of the lungs yellowish or yellowish-grey, round or angular, rather well-defined nodular projections, up to the size of a walnut; either soft and elastic or firm in consistence, and extending in the form of a wedge into the lung tissue. These large nodules are generally found towards the upper edge of the lung, but also at the apex and along the under edge; they are never found isolated in the depth of the lung parenchyma. Besides these larger lesions there are also met with very small, deep yellow nodules, about the size of a hemp seed, sometimes vesicular, sometimes firm in consistence.

The larger nodules when incised allow the escape of a milk-white frothy liquid, in which, with the aid of a lens one can see small worms, some brown (males), some white (females). The length of these worms is estimated by Koch at 20-30 mm. The author's view of the life history of these parasites is given in the following sentences:—"The females lay their eggs in the finest bronchial twigs and in the alveoli, whence the embryos creep out to gain the outer world by the same way as that by which the parent worms have gained access to the lungs, namely, through the large bronchi and the trachea." . . . "The parent worms, however, have fulfilled their purpose after completion of their sexual activity, and they wander in the interstitial lung tissue, most of them until they arrive under the pleura pulmonalis, boring their way through the interspaces of the incomplete cartilaginous rings; they then become encapsuled after the manner of a muscle trichina, calcify, and perish. These encapsuled parent worms are the previously mentioned small nodules of the size of a millet seed, in which we, as a rule, find only one worm repeatedly coiled like a clue; with the steadily progressing calcification the worm becomes destroyed, and hence in old nodules only the rudiments of the parasite are to be found."

Apparently, when Koch began his monograph he was under the impression that the parasite which he describes under the name *Pseudalius ovis-pulmonalis* was a species distinct from the *Strongylus rufescens*, but in the concluding pages he expresses doubt on this point, and suggests that the *Pseudalius* is a metamorphosed and diminished *Strongylus rufescens*, the change in the shape and size of the latter being caused by its penetration into the finest bronchi. He was led to this conclusion by finding in some lungs, along with the *Strongylus filaria*, a few small brownish-red worms, and embryos which were identical in appearance with those which he had previously described as belonging to the *Pseudalius*. One of these worms was identified by Prof. Bugnion as an example of the *Strongylus rufescens*, while Prof. Leuckart declared an incomplete specimen submitted to him as possibly belonging to that species. Unfortunately, Koch does not describe the lesions that were present in the lungs in which he found these worms.

Turning next to the French authors, one finds that Prof. Railliet, in his *Elements de Zoologie Médicale et Agricole* (1886), describes two species of strongyle as being parasitic in the air passages of European sheep, viz.: the *S. filaria*, the cause of verminous bronchitis, and the *S. rufescens*, the eggs of which are laid in the pulmonary alveoli, and by their development there give rise to the formation of "localised areas of inflammation." These lesions "present themselves to the naked eye under the aspect of small greyish tumours, which remain for a long time permeable to air. The tumours occupy especially the surface and borders of the lung; in scrapings obtained from these cut surfaces one finds a large number of eggs and embryos, the latter being remarkable by having their caudal extremity recurved in an undulated point. All these areas may, however, at a certain moment affect the caseous character, and even undergo the cretaceous infiltration; finally, they sometimes finish by being more or less completely resorbed."

As far as can be gathered from these passages, Railliet ascribes

the pseudo-tuberculosis of the sheep's lung to the *Strongylus rufescens*.

Neumann, in his *Traite des Maladies Parasitaires* (1888), refers the parasitic lung lesions of sheep to the *Strongylus filaria*, *S. rufescens*, and *S. paradoxus*. The last of these species is said to be rarely encountered, while the first two are stated to be often present in the same individual, the *S. filaria* being accountable for bronchitis, and the *S. rufescens* for a pneumonic lesion, which ordinarily accompanies the parasitic bronchitis. Generally, it is said, the lesion determined by the *S. rufescens* has the form of a tubercle, varying in size from a millet seed to a pea, greyish-yellow in colour and semi-transparent, or of a more or less intense dark red colour, according to the age and acuteness of the inflammatory phenomena. These lesions owe their production to the eggs deposited by the females of the *S. rufescens* in the pulmonary alveoli.

From this analysis of the views expressed by different authors regarding the strongyles that are parasitic in the air passages of the sheep, it will be seen that the divergence of opinion is mainly with regard to the cause of what may be called the worm-tubercle. Regarding the specific characters of the *Strongylus filaria*, and its relationship to epizootics of bronchitis in lambs, all authors are in agreement. On the other hand, the following different views are held regarding the parasites found in the tubercular lesions :—

1. That they belong to the species *S. filaria* (Williams and Zürn).
2. That they belong to the species *S. ovis-pulmonalis* (distinct from the *S. rufescens*), and represent young worms that have become encysted while acquiring their sexual maturity (Curtice).
3. That they belong to the species *Pseudalius ovis-pulmonalis* (possibly not distinct from the *S. rufescens*), and represent individuals that have become encysted after having discharged their sexual functions (Koch).
4. That they are immature individuals of the species *S. rufescens* (Railliet).

I purpose now to describe here my own observations regarding some of these disputed points.

The Strongylus Rufescens.

I have on a considerable number of occasions during the last five or six years met with this species in the lungs of sheep from different parts of England, Scotland, and Wales. In every case the parasites were without difficulty identified by the description and figures given in Prof. Railliet's work. The characters of the species are there given as follows :—

"Body very slender, and reddish-brown in colour. Head not winged; mouth surrounded by three papilliform lips. The male is 18-28 mm. long; its caudal bursa is excised behind and offers two slight lateral notches in front; the posterior ribs are short and not very distinct, the middle are separated, and the anterior split and reunited (*fendues*). The spicules are curved and striated transversely. The female is from 25-35 millimetres long; its tail is terminated in a blunt point; its vulva is situated immediately in

front of the anus, at the base of a small eminence. Oviparous, eggs ellipsoidal, from 75-120 μ long by 48-52 μ broad."

In most of the cases that have come under my observation the *Strongylus rufescens* was the only species present, but in a few instances it was accompanied by the *S. filaria*. The worms were never numerous present in the bronchi as far as these could be followed with fine scissors, and in some cases only two or three could be found although numerous lesions of the kind described below were present in the parenchyma of the lung. Females were always more numerous present than males.

Lesions Excited by the S. rufescens.—These are very characteristic, and, once seen, are easily recognised even without having recourse to microscopic examination. Plate III. reproduces the appearance of a typical case. The uncut lung shows under the pleura a variable number of greyish patches, which in the collapsed organ are rather more prominent than the surrounding lung tissue. To the touch these patches are less resilient than the normal lung tissue, but they lack the solidity of hepatised or collapsed lung, and when excised, and thrown into water they float. They are irregularly conical in shape, the surface underlying the pleura representing the base of the cone. When incised and scraped they yield a turbid, frothy liquid, and when this is examined under the microscope it invariably shows the eggs and embryos of the *Strongylus rufescens*, and in some cases fragments of the adult worms also. The recently laid eggs are brown, granular, and ellipsoidal, but in a single scraping or section eggs may be found showing all stages of development up to the fully-formed embryo worm. The liberated embryos have a characteristic, curved, sharp-pointed caudal extremity, and in the fresh state they exhibit active movements under the microscope. The bronchi generally contain an excess of frothy mucus, but in the larger tubes this is seldom very abundant unless the *S. filaria* is present along with the *S. rufescens*. The bronchial mucus, even in parts of the lung where the parenchyma is normal in appearance, sometimes shows large numbers of young worms, and in one or two cases I found young worms of both species (*S. filaria* and *S. rufescens*) abundantly present together. In such a case the two species are easily distinguished by the above-mentioned character of the tail in the *S. rufescens*.

The structure of one of the semi-consolidated patches represented in Plate III. is shown in Fig. 5, Plate IV. A small piece of one of these patches was stained in bulk with alcoholic borax carmine, and cut after embedding in paraffin. The figure reproduces a photograph of a small area near the inner limit of the patch—that is, where the diseased and healthy lung tissue merge into each other. The earliest change observable is a broadening of the alveolar septa, and a swelling of the epithelium. The septa become crowded with leucocytes, while the air-cells become reduced in size and partially or completely filled with leucocytes and shed epithelial cells. The terminal bronchioles are affected in precisely the same way, and where the lesion is most advanced the normal appearance of the lung tissue is completely obliterated by a dense infiltration of small round cells (uni-nuclear leucocytes). As might have been expected, the unsegmented eggs are chiefly found in the least altered parts of the patch,

RIGHT LUNG OF SHEEP, showing lesions caused by the *Strongylus myfascia*. The figures 1, 2, 3 indicate three of the invaded patches.

lying singly or in small groups in the air-cells. Eggs in various stages of development and free embryos are found in the more solid part of the patch, but in the latter situation one may find nearly a whole field under a low power ($\times 60$) densely crowded with leucocytes among which one can detect neither eggs nor embryos. It is not improbable that the liberated embryos in general do not remain long in the vesicular tissue of the lung, but soon wander into the bronchi, from which they may be expelled in large numbers during fits of coughing. It appears certain that the young parasites do not remain long enough in the lung to attain to anything like maturity, as even in lungs that give evidence of disease of considerable standing one fails to find any intermediate stages between the adult worm and embryos such as those shown in Fig. 4, Plate IV.

The lesion illustrated in Plate III. and Fig. 5, Plate IV. is undoubtedly the one most commonly excited by the *S. rufescens*, but it is not the only one. In vol. i., p. 139 of this Journal, Mr J. Rowland Dykes described a disease which prevailed among the sheep on a farm in his district (Wellingborough), and which was characterised by very severe lung lesions. From specimens sent to me by Mr Dykes I ascertained that the lesions in question were caused by the *S. rufescens*, but in some of the cases they were much more serious than those commonly caused by this parasite. In nearly every case there was extensive consolidation of the lung; large patches, up to the size of one's hand, being converted into a dirty-white fleshy tissue. There had also been more or less extensive pleurisy, with the consequent formation of fibrous adhesions between lung and chest wall, and in some cases the solidified part of the lung contained phthisis-like cavities filled with thick, yellow, purulent matter. Neither eggs, embryos, nor adult worms could be found in the more solid parts of these lungs, but the surrounding tissue contained both eggs and embryos, and showed alterations similar to those already described (Fig. 5, Plate IV.). The whitish solid parts of the lungs in these cases were found to possess a very remarkable structure, which is represented in Plate IV., Fig. 2. This figure reproduces a photograph of a section made from a piece of lung stained in bulk with alcoholic borax carmine and embedded in paraffin. There is here such a profound departure from the normal structure of the lung that even an experienced histologist might be pardoned for failing to recognise the section as one of lung tissue. The alteration consists in: (1) a great thickening of the alveolar walls, and (2) a peculiar transformation of the alveolar and bronchial epithelium. The former of these changes is due to the development of a delicate spindle-celled connective tissue, while the alveolar transformation consists in the assumption of a cubical or short-columnar shape by the flat epithelial cells that form the lining of the normal air-cells. The cells that line the very small bronchi are also altered to high columnar epithelium, and in some of these the mucus membrane has papilla-like outgrowths projecting into the lumen of the tube. This overgrowth of the bronchial mucous membrane in some cases has a striking resemblance to the lesion excited in the bile-ducts of the rabbit's liver by the presence of the *coccidium oviforme*.

A very similar lesion to the one here described, is found in congenital syphilis of the human subject, and Ziegler, in his *Pathologische*

Anatomie (7th Edition, p. 707), gives an illustration of this syphilitic pneumonia which is almost identical with Fig. 2, Plate IV.

The larger bronchi in pieces of lung showing the before-described lesion of the parenchyma are always profoundly altered, their walls being densely infiltrated with round cells, and their epithelium proliferating and sometimes almost entirely lost. The parenchyma lesion is probably a sort of atrophy of lung tissue that has become functionless from occlusion of its bronchi, or it may be the result of an irritation excited by products absorbed from bronchiectatic cavities and vomicæ in adjacent parts of the lung.

Pseudo-Tuberculous Verminous Pneumonia of the Sheep.

The majority of writers appear to regard the lesion which I intend to describe under the above heading as not distinct from that which is commonly excited by the *S. rufescens* (Plate III.), but such an opinion is not in harmony with my own observations. This pseudo-tuberculous lesion of the sheep's lung is of comparatively common occurrence, and, so far as Great Britain is concerned, it is far more frequently encountered than the form of pneumonia previously described as characteristic of the lung when invaded by the *S. rufescens*. I have never met with it in sheep under a year old, and in most cases the animals in which it was present were ewes of several years old. In this respect it does not differ greatly from the *rufescens* pneumonia, for the latter is generally encountered in adults, or in sheep that are at least a year old, but there is this profound difference between the two lesions,—that the pseudo-tuberculous affection is seldom or never the cause of death, or even of disturbance manifested by any symptom during life, while lung invasion by the *S. rufescens* is generally, if not always, attended by serious consequences, and is frequently fatal. Furthermore, I have never met with a combination of the two types of lesion in the same lung, nor have I seen anything to indicate that the *rufescens* pneumonia would, if the animal had survived, have developed into the tuberculous type.

Verminous pseudo-tuberculosis is generally encountered at the slaughter-house in otherwise apparently healthy sheep, and although its true nature was pointed out in this country by Sandie and Padley as early as 1849, and independently by Prof. Brown in 1866, it is still often mistaken for genuine bacillary tuberculosis. Occasionally an ignorant but enthusiastic meat inspector, when he discovers a lung that is the subject of this lesion, promptly seizes the carcass, and returns the case as one of tuberculosis. This, of course, is generally the mistake of a policeman or other lay inspector, but it is to be feared that a similar error is sometimes made by veterinary surgeons and medical officers of health.

In pseudo-tuberculosis both lungs are affected, and both to the same degree. The tubercles are always distributed throughout the lung, never confined to patches of it. They are always most numerous in the superficial strata of lung parenchyma, but they are by no means confined to that, as some authors would lead one to suppose; on the contrary, they are found in the very depth of the lung tissue. The tubercles are about the size of a common pin's head; they are firm when pinched between the finger and thumb, and difficult to dissect out from the lung texture. When cut, or punctured with a needle,

and squeezed, they generally give exit to a speck of thick yellowish-white matter, and when this is examined under a low power of the microscope it shows a slender but relatively long and intricately coiled worm. The tubercles probably never contain a "worm brood," though sections of the coiled worm embedded in a tubercle may suggest such an idea. Sometimes the tubercles are gritty on section from the presence of calcareous granules. The lung tissue between the tubercles generally appears normal.

The minute structure of these pseudo-tubercles is represented in Plate IV., Figs. 1 and 3. These figures reproduce photographs of two tubercles, both from the same lung, and they illustrate the two commonest varieties. In the one (Fig. 3) the coiled parasite is immediately surrounded by a mass of small closely packed leucocytes with deeply staining nuclei (alum carmine). This mass of leucocytes is sharply limited outwardly by a narrow pale belt made up of "epithelioid" cells and giant-cells. The latter are usually indistinct and imperfect as compared with the giant-cell of the tubercle proper. The exact outline of the giant-cell is often difficult to make out, but when distinct the cell is usually seen to be elongated in the line of a radius of the tubercle, and its nuclei are grouped at its peripheral pole. The giant-cell zone is succeeded outwardly by another which is three or four times as broad as that, and which is composed of vesicular tissue rendered solid by thickening of the alveolar walls, and crowding of the alveoli with round cells, considerably larger than those that immediately surround the parasite. This outer zone passes gradually into the still permeable lung tissue.

In the second form of tubercle (Fig. 1), which appears to represent a later stage in their formation, the worm is embedded in a homogeneous or granular, diffusely staining, structureless material, obviously resulting from the death and degeneration of the pus-like cells that compose the central zone in the younger tubercle. This may be limited outwardly by a giant-cell zone, or that may be absent and its place taken by a few strata of spindle cells, which pass by transitional forms into the round cells of the outer zone. Occasionally the entire tubercle is composed of small round cells, closely packed together, except at the periphery, where they crowd the still recognisable air vesicles.

The lung tissue between the tubercles is generally almost normal, save that it shows a very distinct hypertrophy of the non-striped muscular tissue. This tissue is greatly but irregularly increased in thickness in the small bronchi, and from there it appears to extend backwards, even into the walls of the air-cells. The medium-sized and large bronchi do not share in this hypertrophy of the bronchial muscle.

Such are the macroscopic and microscopic appearances of these pseudo-tubercles, and in conclusion a word or two may be said about the parasite itself. I do not know upon what grounds those who maintain that it represents an encysted stage of the *S. filaria* (Williams, Zürn) have arrived at that opinion, for the worms found in these tubercles, in my experience, have never attained to their full sexual development, and do not possess characters by which they can be certainly recognised as belonging to the *S. filaria*. Nor can they be certainly referred to the *S. rufescens*. The assumption that the parasites belong to the latter species, that they acquire sexual maturity while within the tubercle, and that they leave the tubercles

to copulate in the bronchi, is entirely opposed to the observed facts. If that view of them were correct one ought to find remains of pseudo-tubercles in every lung showing the lesions depicted in Plate III., but such is not the case. Besides, a glance at Fig. 3, Plate IV., will show how improbable it is that these imprisoned parasites could, by their own exertions, escape from the tubercles. Koch's opinion that the encysted parasites are fully developed worms belonging to the *S. rufescens*, or to his *Pseudalius ovis-pulmonalis*, is difficult to understand on any other assumption than that he has described for the encysted form the fragments of adult examples of the *S. rufescens* which he obtained by scraping patches such as those shown in Plate III. Certain it is that if he found in worm-tubercles like those shown in Figs. 1 and 3, Plate IV., sexually mature parasites, such as he describes under the name of *Pseudalius ovis-pulmonalis*, the observation stands by itself. That view of the encysted worms is also negatived by the fact that there is no correspondence, such as one would expect in that case, between the distribution of the pneumonic patches (Plate III.) and the pseudo-tubercles throughout the lung. The former lesions are always irregularly disseminated in each lung, and often unequally distributed in the two lungs; but the pseudo-tubercles are always present throughout both lungs, and never have a patchy distribution.

Reviewing all the evidence, there does not appear to me to be sufficient grounds for assigning the immature worm that is found in the pseudo-tubercle of the sheep's lung to either the *S. rufescens* or the *S. filaria*. The ordinary distribution of the tubercles rather suggests that they may have formed round parasites developed from eggs arrested in the pulmonary arterioles, like the pseudo-tubercles caused by the *S. vasorum* of the dog. In this connection it is interesting to note that Prof. Williams (*Veterinary Medicine*, 6th Edition, p. 772) states that he has on two occasions seen the *S. filaria* in the cavities of the heart and in the blood-vessels. Further observations ought to be made whenever the opportunity occurs, in order to discover whether adult worms of that or any other species are present in these situations in cases of pseudo-tuberculosis.

DESCRIPTION OF PLATE IV.

FIG. 1.—Pseudo-tubercle from Lung of Sheep. *a*. Segment of the worm; *b*. Zone in which the cells have undergone degeneration.

FIG. 2.—Chronic Lung Lesion caused by the *Strongylus rufescens*. *a*. Delicate spindle-celled tissue developed in the alveolar septa; *b*. A metamorphosed alveolus with columnar epithelium.

FIG. 3.—Pseudo-Tubercle from Lung of Sheep. *a*. Coils of central worm; *b*. Zone of closely packed leucocytes; *c*. Narrow zone of giant-cells; *d*. Outer zone in which alveolar structure is still faintly discernible; *e*. A normal bronchus.

FIG. 4.—Two Embryos of *Strongylus rufescens* in a scraping from one of the pneumonic patches seen in Plate III.

FIG. 5.—Eggs and Embryos of *Strongylus rufescens* in a scraping from one of the pneumonic patches seen in Plate III.

FIG. 6.—Section from the deeper part of one of the Pneumonic Patches shown in Plate III. *a*. Part of an adult worm in an infundibulum; *b, b, b*. Eggs lying in air cells; *c*. More solid part of the patch, where air-cells and alveolar septa are infiltrated with small round cells.

For description of figures, see opposite page.