cited in the text, as obtained in the case which forms the basis of these few observations.

I subjoin two delineations of the appearance of the aortic arch and its branches in this instance,—one as seen *in situ* from the front, and the other as observed on looking at it from the left side, which may help to render my description the more readily intelligible.

Fig. 1.

Fig. 2.

Fig. 1. The Aortic Arch, &c., as seen from the Front.

- a. Arch of the aorta.
- b. Bodies of the dorsal vertebrae.
- c. Intercostal arteries.
- 1. The right carotid artery.
- 2. Left carotid artery.
- 4. Right subclavian artery.

(The left subclavian is in this figure concealed by the left carotid artery, behind which it arises.)

Fig. 2. The Aortic Arch, &c., as seen from the Left Side.

- a. Ascending aorta.
- 1. Right carotid artery.
- 2. Left carotid artery.
- 3. Left subclavian artery.

PRACTICAL OBSERVATIONS.

By JOHN CHIPPENDALE, Esq.

I FIND that in a short notice which I addressed to you on the subject of idiopathic erysipelas, I neglected to say, that of the mixture of infusion of senna and decoction of cinchona which I recommended, and which should be made in such proportions as to produce eight ounces, one ounce is a dose.

As I believe you are always the best pleased when the communications of your correspondents are of a practical nature, I will avail myself of this opportunity of conveying a few observations of that kind to you, trusting they may be of use to some of your readers.

In your publication of this day I find a note from Dr. Allnatt respecting the "Best Remedies for Neuralgia." Conversing, some short time ago, with a professional friend on the subject of this disease, he told me that he had found considerable advantage to accrue, after everything else had failed, from frictions with a strong infusion of tobacco, to which he adds the potassio-tartrate of antimony. These appear to me to be so opposite in their nature that I should scarcely think of employing them in

conjunction; but I have tried the infusion of tobacco in two cases, and have, in both of them, met with complete success. It appears to me that an extract might be prepared from the tobacco, which, being mixed with simple cerate, would be a more convenient form for frictions.

One of your correspondents, Mr. M. K. O'Shea, in speaking of fractures of the bones of the forearm, advocates the prone position of the limb when the fracture is put up. He will find that doctrine inculcated in Mr. Lonsdale's excellent "Treatise on Fractures," which (I speak from memory, not having the book by me) must have been published some five or six years. There is, however, another important point which, Mr. O'Shea has not mentioned, which is, that a wedge-shaped pad should be placed between the bones, to counteract the efforts of the pronator quadratus muscle.

I will close with a practical remark on catheterism. Sir Astley Cooper says that when it is desired to introduce a catheter into the bladder during the existence of enlargement of the prostate gland, an instrument of elastic gum may be selected, which being introduced as far as to the obstruction caused by the enlarged gland, the stilette may be partially withdrawn, by which process the point of the catheter will be raised, and may then be pushed on into the bladder. This is a very useful hint, but I would beg to add to this a caution to junior practitioners, to the effect that the stilette having been thus partially withdrawn, there should not be any attempt made to return it whilst the instrument remains in the passage; for if the gland is only laterally enlarged the point of the catheter may be so diverted from the straight line that the eye may allow the stilette to pass outside, and thus lacerate the parts. This happened to me some years ago, although the result was fortunate, for, by this manœuvre, an abscess, of which I did not even suspect the existence, was punctured, a large quantity of pus was discharged, and my patient recovered; so that I acquired a share of glory which did not really belong to me. But it is not every case which might terminate thus fortunately.

Great Queen-street, April 6, 1844.

MIDWIFERY REPORT.

DURATION OF LABOUR AND LENGTH OF THE FUNIS.

By G. Y. HUNTER, Esq., Surgeon, Margate.

IN my letter which you inserted in THE LANCET of the 5th of Nov., 1842, I promised to send you the result of my continued investigation connected with the above subject, and, therefore, I now forward a table of 123 midwifery cases, in which I have been able to measure the funis. You will see that in some instances I have added the age of the parents, in hopes of aiding the inquiry which has been commenced by the German writers, for the purpose of ascertaining how far the sex of the infant depends upon the father's age; a question which might soon be set at rest if the registrar-general could be induced to direct the district-registrars in England and Wales to insert, in the proper column of each book of births, the age of the father and mother, whenever it could be obtained. As I have also, in the accompanying report, noticed the deviations from nature, it may be considered as an appendix to the table of 2250 cases published at page 537 of THE LANCET, of January 7th, 1843. In conclusion, I shall briefly observe that in my opinion neither the great length nor extreme shortness of the funis is likely either to expedite or retard delivery, at least, when the contraction of the uterus is strong enough to expel the fœtus, it being clear to me that a very short funis would be soon torn from the placenta, as in case numbered nine (an event which I have known to happen before), or the placenta ejected along with the child, which I have often seen occur in labours where the pregnancy had advanced as far as the seventh month. If my life should be prolonged to the close of the year, I hope again to transmit you the result of my observations in this most useful and highly-important branch of our profession.

Number of each case.	Whether 1st or 2nd pregnancy, or otherwise.	Duration of labour in hours.	Length of funis in inches.	Sex.		Age of parents.		Funis twisted round the neck.	Born with cauls.	Abortions; period in weeks.	Premature labours; period in months.	Remarks.
				Male.	Female.	Father.	Mother.					
1	2nd	3	28	—	—			—				
2	6th	9½	30½	—	—			—				
3	11th	3	35	—	—			—				
4	2nd	1½	25	—	—			—				
5	7th	1½	24	—	—			—	—			
6	3rd	1½	18	—	—							
7	8th	1½	21	—	—							
8	7th	5½	25	—	—							
9	4th	14	21	—	—							{ Funis torn from the placenta, which was quickly expelled.
10	4th	12	33½	—	—							
11	10th	1	48	—	—			—				
12	6th	2	20	—	—			—				
13	10th	1		—	—					10		
14	1st	16	29	—	—	24	20					
15	6th	1	34	—	—	42	33	—				
16	2nd	7	23½	—	—	35	32					
17	3rd	20	21½	—	—							
18	6th	1½	22½	—	—	35	24					
19	3rd	1½	27½	—	—	48	35					
20	7th	1½	27½	—	—	42	38					
21	6th	1	27	—	—	37	35					
22	3rd	1	18	—	—	26	25					
23	1st	3	33½	—	—	35	28					
24	8th	1	23½	—	—	39	42					
25	4th	1½	19½	—	—	35	32					
26	4th	3	19½	—	—	40	31					
27	1st	5		—	—							
28	1st	12		—	—		21					
29	4th	1½	42	—	—	28	28					Annular palate.
30	6th	1	20	—	—	40	40					
31	2nd	1½	27½	—	—	25	23					
32	6th	1	34	—	—	53	39					
33	2nd	1	17	—	—							
34	4th	1½	28	—	—	35	33					
35	5th	1	15	—	—	29	23					
36	9th	1½	29	—	—	35	35					
37	4th	1½	23	—	—	25	24					
38	1st	7	13	—	—		18				6	Dead some time.
39	4th	3	16	—	—							
40	3rd	1	31	—	—	27	24					
41	2nd	4	21	—	—	30	33					
42	6th	2	32	—	—	35	30					
43	5th	2½	24	—	—	31	26					
44	4th	1	27½	—	—	31	29					
45	1st	3	20½	—	—	29	28					
46	2nd	1½	20	—	—	25	23					
47	10th	1½	34	—	—	44	42					
48	5th			—	—							
49	1st	2	12	—	—	37	32					Club-feet; funis torn off.
50	4th	1½	22½	—	—							
51	3rd	2	24	—	—	34	33					
52	7th	18	40	—	—	41	40					Breech presentation.
53	4th	1½	30	—	—							
54	4th	1½	30	—	—	25	29					
55	3rd	3	29	—	—	50	52					
56	4th	1	36	—	—	25	22					
57	7th	1½	38	—	—	29	30					
58		1½	38	—	—		24				7	
59	4th	1½	31½	—	—	33	27					
60	5th	3	37½	—	—	33	30					
61	2nd	1½	31	—	—		32					
62	1st	2	23	—	—	25	26					
63	4th	4	24½	—	—	33	33					
64	1st	2		—	—	23	23				5	
65	6th	1½	22½	—	—	32	32					
66	3rd	1½	39½	—	—	26	28					
67	9th	2		—	—							
68	2nd	1½	32	—	—							
69	2nd	1	31½	—	—	28	26					
70	7th	1	30	—	—							
71	4th			—	—	27	27					
72				—	—					6		
73	3rd	1½	33½	—	—	30	28					

Number of each case.	Whether 1st or 2nd pregnancy, or otherwise.	Duration of labour in hours.	Length of funis in inches.	Sex.		Age of parents.		Funis twisted round the neck.	Born with cauls.	Abortions; period in weeks.	Premature labours; period in months.	Remarks.
				Male.	Female.	Father.	Mother.					
74	8th	1 1/2	35 1/2			35	34					
75	3rd	1 1/2	36									
76	1st	3 1/2	37 1/2									
77	3rd	5	32			37	32					
78		3	36 1/2									
79	4th	1 1/2	22 1/2									
80		2	28 1/2									
81	1st	3	33 1/2									
82	1st	1 1/2	27									
83	5th	1 1/2	24			39	42					
84		2	32 1/2									
85	3rd	1 1/2	32 1/2									
86	1st	3	31			26	22					
87	7th	2	32 1/2									
88	7th											
89	8th	1	{ 24 } { 25 }								7	
90	2nd	1 1/2	21 1/2									Twins.
91	1st	1 1/2	42									
92	6th	1 1/2	28									
93	2nd	1 1/2	24									
94	3rd	1										
95	1st	5 1/2	38			23	25			10		
96	2nd	1 1/2	30			32	35					
97	6th	1	20			26	27					
98	3rd	13	32			22	24					
99	1st	6	26			39	34					
100	8th	1 1/2	21			22	21					
101	2nd	1 1/2	22									
102	2nd	1 1/2	27									
103	8th	1 1/2	39									
104	4th	1 1/2	39									
105	5th									10		
106	3rd					30	32				7	Dead some time.
107	8th	1	18			41	41				8	Dead some time.
108	4th	16	33			25	24					
109	5th	3	22								7	
110		2 1/2	32									
111	4th	1	30									
112	4th	1	38									
113	3rd	2	{ 20 1/2 } { 26 1/2 }									Twins ; one breech presentation.
114	7th	1 1/2	20 1/2									
115		1 1/2	22									
116	7th	1 1/2	32									
117	1st	3	30									Head embraced by the child's hands.
118	3rd	1 1/2	31									
119	1ts	44	24									
120	3rd	6	30									{ Hour-glass contraction ; removed { placenta.
121	5th	5	27									
122	8th	1 1/2	28 1/2									Face presentation.
123	2nd	2	36									
		385	3175	58	63			11	2	4	8	

Average length of funis, 28 1/2 inches. Average hours of labour, 3h. 7m.

PRACTICAL OBSERVATIONS
ON THE
CURATIVE EFFECTS OF BELLADONNA
IN CERTAIN
AFFECTIONS OF THE NERVOUS SYSTEM.

By R. S. HUTCHINSON, M.D., M.R.C.S., Senior Physician
to the General Hospital near Nottingham, &c.

(Continued from last volume.)

TETANUS.

THE morbid affections of the true spinal system of nerves divide themselves, first, into those which affect the nerves of sensation only, as in the great class of neuralgiæ ; secondly, into those attacking the motor nerves

alone, as in chorea, and perhaps epilepsy ; and, thirdly, into those which morbidly affect both the sensor and motor nerves, as in tetanus, hydrophobia, spasmodic tic, and laryngismus stridulus. The simple inflammatory affections of the medulla spinalis and its membranes are not here included ; but, it may be remarked in passing, that a rheumatic inflammation, acute, subacute, and chronic, of the theca vertebralis, has a much more common pathological existence than is ordinarily admitted, some observations and cases of which I published in THE LANCET several years ago ; and it is probable that this may frequently constitute the cause productive of some or other of the disorders of the true spinal system previously mentioned.
The beneficial influence of the belladonna over the neuralgiæ has constituted the subject of preceding com-