Reviews.

Die Geschwülste des Nervensystems; Hirngeschwülste, Rückenmarksgeschwülste, Geschwülste der peripheren Nerven—Eine Klinische Studie. Von DR. LUDWIG BRUNS. Mit 31 Abbildungen im text. Karger, Berlin, 1897 (Williams and Norgate, London), 12s. Pp. 388.

ALTHOUGH from the striking character of the symptoms cases of tumour of the brain or spinal cord must always have excited great interest, yet in former times this interest lacked practical bearing, partly from the hopeless nature of the illness, and partly from the want of sufficient data on which to make an accurate diagnosis of the tumour during life. Three causes have within late years greatly stimulated investigation in this field; first, the use of the ophthalmoscope; secondly, the discovery of the localisation of function in certain parts of the brain; the one giving in the presence of optic neuritis a certainty to the general symptoms of brain tumours which could not be obtained before, and the other establishing the reliability of the local symptoms; and thirdly, the demonstration that tumours of the brain and cord could be successfully removed by operation. The rapid advance in the knowledge of these tumours that has followed has resulted from the increased attention thus directed on the subject. Of this knowledge and its most recent developments, this book gives a very complete account, and the reader may turn to it with confidence that nothing of importance has been omitted. The author has taken pains to bring his information fully up to date, and his own experience and wide reading qualify him to deal authoritatively with his subject.

With regard to the pathology of brain tumours the gliomata are attributed to overgrowth of neuroglial tissue, and the author thinks that they must be sharply differentiated from the sarcomata, both on histological grounds and on those of their origin from an ectodermal structure. An important practical distinction from most sarcomata consists in the infiltrating character of the

gliomata, so that they may take the form of that part of the brain in which they lie, and from lack of demarcation be incapable of removal. Nerve-fibres and cells within the limits of the tumour may retain their functional capacity, so that the symptoms may be slighter than would seem likely from the extent of the tumour.

Dr. Bruns thinks that traumatism is doubtful as the only exciting cause of a brain tumour, but considers that in syphilis and tubercle it may, by forming a *locus minoris resistentiæ*, determine the seat of a syphilitic or tubercular growth. A traumatism may, however, first call attention to the existence of a tumour, either by causing hæmorrhage into a glioma, or by so altering the surroundings or conditions of any tumour that it takes on active growth, and in each case thus first gives occasion for distinct symptoms to arise. Whilst admitting the cogency of these arguments, however, it seems doubtful whether the frequent history of injury to the head shortly before the appearance of symptoms of a brain-tumour is thus fully explained by them.

A chapter follows on the general effects of the presence of a tumour on the brain itself, on the cranial nerves, and on the ventricles, which will be found of much interest and value.

Dr. Bruns adopts the mechanical theory of the origin of optic neuritis, which he thinks amply sufficient to account for this symptom, supported as it is by recent observations on the subsidence of optic neuritis after operations to relieve increase of intra-cranial pressure. He agrees with Oppenheim's statement that about 90 per cent. of all cases of optic neuritis are due to the presence of an intra-cranial tumour. He accepts the division of the symptoms of a brain tumour into general and focal as one of great practical importance and founded on a sound pathological basis, but points out that the line between them must not be too sharply drawn, as "general" symptoms may under special circumstances have a focal or local significance, *e.g.*, those due to the early appearance of internal hydrocephalus in tumours of the posterior fossa, and on the other hand symptoms usually focal in import may be in some cases "general," as when pressure on the olfactory nerves or optic chiasma arises from increase of the general brain pressure. The characteristic mental change in intra-cranial tumours is considered to be mental dulness or stupidity; marked psychical symptoms are specially frequent in multiple tumours.

Other things being equal, the smaller the growth and the slighter the general symptoms the more certain is the local

diagnosis. In the localisation of tumours considerable importance is to be attached to increased tenderness of the skull to percussion. When the tenderness is general over the skull there is likely to be a general increase of internal pressure; this is apt to occur early in the illness in large tumours in any situation, and also in tumours of the cerebellum. Local tenderness, if it exists over the position that other signs indicate as the seat of the tumour, is a guide to its localisation, and tends to show that it lies in or near the cerebral cortex, or in the membranes, though this sign may be present in deep-seated tumours. A tympanitic percussion note, and *bruit de pôt fêlé* are also sometimes met with, and, if these signs are localised, and correspond to the position assigned to the tumour on other evidence, are valuable in diagnosis. The author states that he holds tenderness on percussion, tympanitic note, or *bruit de pôt fêlé*, to be such valuable signs that, if, for instance, other symptoms indicated the motor convolutions as the seat of a tumour, the presence of these local signs over a neighbouring area, say, in the frontal or parietal region, would be sufficient to indicate one of the latter regions as the position in which to trephine. Both signs are most evident and pronounced in children, and are sometimes met with in other conditions than tumour. In the section devoted to the consideration of the symptoms and signs evoked by tumours in special regions of the brain, full details are given of those proper to a growth in each region. It is stated that a tumour of the optic thalamus can be diagnosed with some certainty if symptoms of hemiplegia with distinct disturbances of sensibility, perhaps with hemiopia, and especially with one-sided athetosis and paralysis on the same side of the facial nerve muscles for mimetic or emotional expressions, but not for voluntary movements, are present. Dr. Bruns thinks that it will only very seldom be possible to make a diagnosis of a tumour of the corpus callosum; he denies that there is any especial tendency to mental changes in tumours of the præfrontal area, and when such symptoms are present he attributes them to the fact that a tumour in this region can grow to a large size before it endangers life, an argument which does not apply to those observed cases in which a frontal tumour of moderate size has produced early pronounced mental symptoms. He lays stress, however, as an aid to diagnosis, on the hilarious or humorous mental state, and also on the ataxia which may be observed in cases of tumour in this situation. A useful table which is given of the accompanying symptoms in cerebellar and frontal ataxia

shows how closely the group of symptoms may resemble one another in each case.

Whilst he admits the occasional presence of sensory disturbances in tumours of the central convolutions, especially the occurrence of ataxy and disturbance of the power of localisation of the sense of position, he thinks that there is great individual variation in this respect in different persons, the difference being unaccountable unless it is due to the greater extent of sensory (than motor) representation in the cortex, and hence the greater possibility of compensation.

With regard to the diagnosis of the side on which a cerebral tumour is situated, it is pointed out that large tumours may excite symptoms by compression of the opposite hemisphere.

Two forms of ataxy in cerebellar tumours are recognised: (1) the staggering or drunkard's gait, and (2) a gait and movements like those of locomotor ataxy; and ataxy is considered to be more constant in tumours of the vermis than in those of the hemispheres, but cannot be taken as a guide to the position of the tumour in the cerebellum. Tumours of the middle peduncle or of the lateral lobe compressing the middle peduncle are stated to cause movements of rotation of the body round its longitudinal axis.

Cerebellar tumours are noted to be extremely seldom latent.

There is a very good section on the Diagnosis of Tumours at the Base. The author considers that an intra-medullary tumour of the pons can only be certainly distinguished from one lying outside it if the symptom of loss of conjugate movement of the eyes to one or other side has been present from the beginning. With regard to the attainment of an accurate local diagnosis, this can be best accomplished in tumours of the motor centres, of the base (posterior and middle fossa), and of the pons, though in many cases of tumour of the posterior fossa one has to be content with that diagnosis without more accurate localisation. In tumours of the speech regions, cerebellum, medulla oblongata, 4th ventricle, white matter of the occipital lobe, the local diagnosis is less certain. Those of the frontal lobe come in this group, if the local evidence and signs of affection of neighbouring parts be clear. Such a diagnosis is uncertain or impossible in growths of the parietal and right temporal-sphenoidal lobes, of the corpus callosum, centrum ovale, and great ganglia.

In cases in which the general and local symptoms are pronounced, the diagnosis is a certain one in 80 per cent.

With regard to surgical treatment, practically the only treat-

ment in intra-cranial tumour, the author states that taking everything into consideration, 60 per cent. of tumours in which a local diagnosis is possible are unsuited for operation, because they are not accessible to the surgeon. In about 32 per cent. of cases of intra-cranial tumour a certain general and local diagnosis is possible, and the tumour is seated in an accessible position. The *exact* conditions present can, however, never be ascertained until the tumour is exposed, and unavoidable hindrances due, for instance, to the size and extent of the growth, its situation and character, must be considered. The 32 per cent. is thus reduced to about 8 per cent. of all cases; in this 8 per cent. the position of the tumour is correctly diagnosed, it is accessible, and there are no hindrances at the operation to complete extirpation. Deducting again from this cases in which unavoidable surgical accidents occur, and the 4 per cent. of fully successful cases given by Oppenheim is nearly reached. These statistics were of course compiled before the recent results obtained at the National Hospital were published by Professor Ferrier; these give a much more favourable aspect to operation.

Taking all things into consideration, Dr. Bruns dissuades from operation in cerebellar tumours. He speaks strongly in favour of trephining as a palliative measure in cases of irremovable intra-cranial tumour, especially if blindness threatens.

The whole question of operative treatment, and the advice to be given to patients in this respect, is treated in a masterly manner. Dr. Bruns would put as clearly as possible the exact conditions before the patient and his friends, and in a suitable case would advise an operation, but would not urge it. With regard to intra-cranial syphilis he says that in his experience the lasting cure of gummatous processes by anti-syphilitic remedies is much more seldom than is generally stated in the text-books; in most cases the improvement is temporary.

The section of the book devoted to Tumours of the Cord is also a very complete one, and gives a full account of their symptomatology and pathology. The regional or segmental diagnosis is entered into very thoroughly, and is brought well up to date; special stress is laid on the importance of Sherrington's discovery of the overlapping in the distribution of the nerve roots; the author thinks that even five roots may in man supply fibres to a single area of sensory distribution. The researches of Head or Thorburn on segmental distribution are also fully utilised. The differential diagnosis between tumours of the spinal column, those within the spinal canal but extradural, and those growing in the

cord itself, receives a full discussion. The directions for localising the position of the growth are very clear and good.

Dr. Bruns especially mentions the frequency of cancer of the vertebræ secondary to mammary cancer; these secondary deposits may occur after so long an interval as eight years after a complete and apparently successful operation on the breast. He thinks that the outlook is so hopeless in cases of tumour of the spinal cord that operation is not only justifiable but obligatory in every case in which a tumour of the spinal cord can be diagnosed with certainty. He gives a short account of the published cases of this operation; the first successful one was the memorable case of Gowers and Horsley. In all there are twenty reported cases of operation for an "occult" spinal tumour; it is interesting to note that the diagnosis was correct in every case, and in eighteen the tumour was found at the operation. In six, or 30 per cent., cure or great improvement resulted, in two there was slight or temporary improvement. In twelve cases the patient died, in nine soon after the operation from shock, hæmorrhage or sepsis. Considering the grave nature of the disease and the difficulties involved in the operation, this may be considered as a good result.

The concluding portion of the book deals with tumours of the peripheral nerves in the same thorough manner. The classification of neuromas into true and false neuromas is shown to be misleading, in that these growths spring from the epi-, endo-, or peri-neurium, most often from the latter, and are all, therefore, false neuromas in Virchow's sense. It is not denied that a growth of nerve substance sometimes takes place in these tumours, but with the exception of many cases of amputation neuroma, it almost always takes a small part in the construction of them. The nerve fibres are most apt to be damaged in sarcomas; in ordinary neuromas, although the nerve bundles are separated from each other and broken up, the nerve fibres often retain their medullary sheaths, and presumably their functional capacity for a long time. The important fact is mentioned that malignant neuromas (sarcoma) often start on small peripheral twigs, and the recurrent growths take place nearer and nearer to the centre until they reach parts no longer accessible to the knife.

I have only alluded to a few points of present interest, but the book itself will be found to give a very thorough account of the whole subject in all its bearings, and will well repay a careful perusal. It is written in a clear style, in a highly interesting

manner, and has evidently been well thought out. The author gives full credit to the work of others, and his cordial appreciation of the labours of English-speaking observers, which he repeatedly quotes and refers to, must be particularly gratifying to us.

The absence of an index is a serious defect to the book as a work of reference, for which it is otherwise very well qualified to serve. This will no doubt be rectified in a future edition.

J. MICHELL CLARKE.

Jahresbericht über die Leistungen und Fortschritte auf dem Gebiete der Neurologie und Psychiatrie. 1. Jahrgang. Bericht über das Jahr 1897. 1 vol. large 8vo, pp. 1508. Karger, Berlin, 1898.

THIS work has just reached us, whilst the last sheet of this number of BRAIN was going through the press. It will prove an indispensable companion to every working neurologist, whose labour in looking over the literature of the subject in hand will be greatly facilitated. It is intended to give a full *résumé* of all the neurological contributions of any value published during the past year; and the names of Drs. Mendel, Flatau and Jacobson, which figure on the title page, are a guarantee that this aim will be reached. The list of working contributors is also promising. The great advantage of a yearly *Bericht* over a *Centralblatt* is that the papers abstracted are arranged under distinct headings, and that the work being done more systematically, the reading of every chapter gives a much clearer notion of the progress made during the year. On the other hand the information is necessarily somewhat belated, so that it is by combining the two sources of information that the student will obtain all the profit derivable from these most praiseworthy efforts of our laborious German collaborators in the field of neurology. It would obviously be a hopeless task to try, in a short notice, to give an idea of the enormous mass of material accumulated in these 1,500 pages. The reader will, however, have no difficulty in finding what he wants, for the divisions and sub-divisions of the work are clear, and the type and paper fairly good. The subjects embrace the whole field of neurology and psychiatry, from the methods of colouring of the nervous elements to questions of criminality and the like. It only remains for us to thank most heartily the editors, authors and publisher who have given us a work of such supreme practical value in our labours, and to hope that their efforts will be further encouraged by the reception this *Jahresbericht* will receive at the hands of Neurologists.

A. DE W.