alveolar spaces is essential but difficult in the absence of the help of the thoracic excursion. A limited dead space is, as has been stated by Yandell Henderson, essential for the retention of the required amount of carbon dioxide.

My experience has been, however, that in the use of a positive pressure apparatus the tendency is to the over-retention of CO₂ from an excessive dead space resulting in excess of CO₂, slowing of the pulse and dyspneic respiration, rather than a tendency to lack of CO₂ (acapnia), cardiac tetanus, and apnea, described by Henderson as the symptoms of hyperventilation.

I have done six experiments by the method of Volhard and Sollman introducing oxygen by catheter through a tracheotomy wound to the bifurcation of the bronchi, with both chests wide open and lungs collapsed.

The alveoli of a two-thirds collapsed lung can thus be ventilated and the animal kept alive. Not only is a direct access of oxygen to the partially collapsed alveoli thus established, but a ready escape and flushing out of carbon dioxide is also provided through the tracheotomy wound around the catheter. In one of two attempts I found it possible to maintain oxygenation by similarly introducing air and ether mixture by tracheal catheter, with both chests opened and both lungs collapsed.

The secret of the success of these tracheal experiments with semi-collapsed lungs is the absolute prevention of a dead space for carbon dioxide retention. In order to avoid tracheotomy and introduce air by the mouth, the lungs must be kept inflated by resisting the outflow of the air compression. The aerating surface is thus increased and the ventilating air is kept in motion as in the tracheal experiments.

I have also observed that when the lungs are held in expansion by a strong resistance to the escape, together with a low supply of air compression, that dyspneic signs of incomplete ventilation come on, and from these observations I have been led to attempt every possible means of improving the gas exchange in the alveoli and thus preventing an absolute dead space. I have, therefore, increased the pressure and rapidity of flow of the afferent "fresh air" by using higher compression, up to about three pounds, yielding a cubic foot of air each minute.

The resistance to the efferent air I have correspondingly diminished so that the lung inflation has remained at its original level. This profuse flushing out of the alveoli is, therefore, a principle which I recommend to prevent the over-accumulation of carbon dioxide, and since its employment I have seen no tendency to dyspnea or cyanosis.

SUMMARY.

The results of thirty operations under positive pressure with 70 per cent. recoveries were encouraging.

Thirty operations and twenty-five experiments proved that emphysema and circulatory changes do not occur with a properly regulated apparatus.

In the use of positive pressure the escape of deoxygenated air must be amply provided for.

I would express my thanks to Prof. Walter B. Cannon for valuable suggestions in the blood-pressure experiments; and to George Adams Leland, Jr. (Harvard Medical School, Class of 1911), for his able assistance.

THE POSITIVE PRESSURE METHOD OF ARTIFICIAL RESPIRATION

WITH ITS EXPERIMENTAL APPLICATION TO THE SURGERY OF THE THORACIC ESOPHAGUS.*

NATHAN W. GREEN, M.D.

AND

J. W. DRAPER MAURY, M.D.

NEW YORK.

It is not the intention in this paper to go over the history or to cover the literature of artificial respiration. There are two distinct procedures included in the term. The first applies to postural methods, the second applies to that produced by apparatus. The latter is used in the surgery of the chest. This may again be divided into the positive pressure and the negative pressure methods. It makes little difference which of these is employed. It is in accordance with obvious surgical principles to apply the most expedient.

Matas,¹ Brauer,² Tuffier,³ F. T. Murphy,⁴ H. H.

Fig. 1. Result of sectioning operation using silk sutures to make the anastomosis: 1, point of anastomosis which was sequestered in the abdominal cavity below diaphragm 3; 2, the normal esophago-gastric opening closed with silk sutures.

Janeway and others have employed positive pressure. Sauerbruch⁵ has been the chief advocate of negative pressure. Woillez,⁶ in 1876, also employed a negative-

* Read in the Section on Surgery and Anatomy of the American Medical Association, at the Fifty-ninth Annual Session, held at Chicago, June, 1908.

1. Matas, R.: Tr. South. Surg. and Gynec. Assn., xii, 52-82; Artificial Respiration by Direct Intra-Laryngeal Intubation, Am. Med., Jan. 18, 1902.

2. Brauer, L.: Mitt. a. d. Grenzgeb. d. Med. u. Chir., 1904, xiii, 496.

3. Tuffier, T.: L'Ouverture de la plèvre sans pneumothorax, Presse méd., Jan. 27, 1906.

4. Boston Med. and Surg. Jour., April 13, 1905, clii, No. 15, pp. 428-431.

5. Sauerbruch, F.: Mitt. a. d. Grenzgeb. d. Med. u. Chir., 1904, xiii, 400-480; München. med. Wchnschr., Jan. 2, 1906; Zentrbl. f. Chir., 1905, xxxii, 82; Beitr. z. klin. Chir., 1905, xvi, 405-494. Stetten, DeWitt: The Experiments of Sauerbruch in the Field of Esophageal Surgery, New York Med. Jour., June 10, 1905.

6. Bull. de l'Acad. de méd., series 2, 1876, v., 613.

pressure box. Its application was, however, different from that of the Sauerbruch chamber.

As the technic of intrathoracic work shall improve, and as the underlying difficulties shall be more thoroughly understood, artificial respiration as an aid will become more general. The possibility of intervention in thoracic visceral disease has been made clear by Tuffier, Maffei⁷ and others. Dr. Sauerbruch's classic work has given a new impetus to the interest in surgical intervention in the chest.

There are difficulties in the surgery of the chest that are not present elsewhere. The chest is a more or less rigid cavity in which the viscera lie deeply anchored. The condition of negative pressure exists. The walls remain rigidly outstanding after either partial collapse of the lung or the removal of part of the enclosed viscera. This latter causes a failure of the viscera to fill the chest at once completely after it has been opened.

There is an apparent tardiness in the serous membranes of the chest to form rapid and plastic adhesions as compared with the peritoneum. Whether this is real or only due to the other conditions within the thorax has not yet been worked out. As to the fibrous adhe-

side of the chest usually means collapse of the lung on the other side as well, unless the lungs are kept inflated by some artificial means, and frequently we have found a communication of one pleural cavity with that of the opposite side. Prof. George S. Huntington regards these openings as artifacts. Furthermore, even if not patent, the mediastinum in dogs is very thin and easily movable, and does not offer the resistance to atmospheric pressure, when one side is opened, that is found in the human being. On this account it would seem that operations for certain conditions in the human subject will prove more successful than the same class of operations for experimental purposes in dogs. This may seem an extraordinary statement, as it is the popular belief that dogs are more resistant to septic infection than are human beings. Our researches, however, have shown that dogs are not as resistant to infection as is commonly believed. In operations in the chest we may

Fig. 2. Side-tracking operation by means of a Murphy button: 1, is on the edge of the normal esophageal opening; 2, is on the edge of the new stoma; 3, is the diaphragm; 4, is the Murphy button impacted in the pyloric end of the stomach, causing death two weeks after operation.

sions, it is well known that they may be very extensive. In other words, speaking surgically rather than histologically, there are two degrees of adhesions: the first is the plastic form, the second is the fibrous or organized. The rapid development of the first-mentioned is the element which makes a large part of the intra-abdominal work possible. The rate of formation of plastic adhesions within the chest is apparently slower. There is a marked susceptibility of the pleural cavity to infection.

After operations in the chest, as elsewhere, septic manifestations may occur. When these have once occurred, it is very hard to establish drainage. This has been our experience in the Research Laboratory. Infection, when present on one side, may rapidly involve the other. We have found in dogs that opening one

Fig. 3. Side-tracking operation by means of a Murphy button containing an obturator in the male portion: 1, normal esophageal opening into stomach; 2, new stoma; 3, diaphragm. (It must be understood that in these operations the cardiac portion of the stomach was drawn up through the diaphragm into the chest and became thereafter an adventitious thoracic viscus.)

get infection from the skin, the severed bronchial tubes, or the opened hollow viscera. Operations on the lungs appear to be less frequently followed by empyema than operations on the alimentary canal. In work on the lungs it is important to have all the cut ends of the bronchial tubes closed in to prevent it. Any method that accomplishes this well should succeed. In a recent article by Samuel Robinson,⁸ of Boston, very good results have been reported. In a previous paper,⁹ one of us has outlined a technic for the prevention of sepsis from the bronchi. In work on the esophagus, sepsis more frequently follows, unless special points in technic applicable to the chest are carefully observed. Whether this is due to the aspirating of air down the esophagus

7. Jour. de Chir. et Annal. de la Soc. belge de Chir., 1906, vi, 380-390.

8. Ann. Surg., February, 1908, xlvii, No. 2.

9. Green, N. W.: Surg., Gynec. and Obst., May, 1906, li, No. 5, 512-538.

while it is opened, thus contaminating the field of operation, or whether it is due to subsequent leakage from some mechanically imperfect closure is hard to determine. It may be from both. H. H. Janeway has suggested that it may, in some degree, be due to the suction of the negative pressure aiding intra-thoracic seepage along the stitches, or retrograde leakage between them.

Dr. Sauerbruch has had very favorable results with the aid of some mechanical device such as the Murphy button. We think that this is due to a more perfect closure of the anastomosis wound and of the lymphatics than can be made by stitches, for when properly applied it makes a tight apposition of the visceral surfaces.

Fig. 4. Special button for esophago-gastric anastomosis (side-tracking operation); A, gastric half of button; B, esophageal half; C, floating ring (elastic) for maintaining continuous pressure on tissues (this is made now as a spring ring); D, forceps for releasing spring which engages needles, E, E; F, complete button partially closed; G, method of uniting halves of button by perforating walls of esophagus and stomach by the needles (the portion of the walls thus included between the halves of the button undergoes necrosis from pressure and sloughs out in a few days). The new stoma shown in Fig. 6 is the result of such a process.

Fig. 5. Apparatus for applying positive pressure in artificial respiration; A, ether chamber filled with gauze; B, intubation tube; C, C, C, valves used to direct current of air either through ether-chamber or directly to intubation canula; D, safety valve; E, inlet for fresh ether; F, air-filter; G, clamp for holding tube in place.

This, unlike the union made by stitches, is an uninterrupted pressure circle. That made by stitches, on the other hand, even if they be continuous, must have points of varying pressure throughout its circumference. These theoretical points have been suggested to us by the difference in mortality, in practically comparing the two methods. Our own experience leads us to believe that some mechanical device which will establish an anastomosis with absolutely no soiling of the serous membrane, forming at the same time a tight joint, will

prove of the greatest assistance in rendering operations of this nature free from disastrous infection. Dead spaces within the chest should be overcome as much as possible, at the termination of the operation by increased air pressure. The absence of omentum with its coapting power is an unfavorable feature which may hinder success. We have carried omentum into the chest and applied it to the wound in one instance. The constant motion of the diaphragm may be a factor which militates against proper union after operation. We have stretched the diaphragm on the operated side, and also interrupted the function of the phrenic nerve in an effort to overcome this.

The difficulty of establishing drainage may, in part, be overcome in the human by some sort of aspirating device.¹⁰ In our experimental work on dogs this is hardly applicable.

Fig. 6. Stomach removed from dog ten days after operation: 1, glass rod passing through esophagus and normal cardiac orifice of stomach; 2, new stoma made by button shown in Fig. 4; 3, diaphragm.

The indications for operation on the esophagus are: stricture of the esophagus, carcinoma of the lower end or of the esophago-gastric juncture, diverticulum of the esophagus, and foreign bodies within the esophagus.

The operations that we have undertaken in the Surgical Research Laboratory on the esophagus may be classified under two types. In one, the esophagus was sectioned, reuniting the proximal end with the cardia of the stomach (Fig. 1). In the other, there was a side-tracking of the cardiac orifice similar to the method of Biondi.¹¹ The sectioning operation was done with

10. Seidel, H.: *Deutsch. med. Wchnschr.* Feb. 22, 1906.
Bryant, J. D.: *Surg., Gynec. and Obst.*, 1906, iii, 296-302.
11. Gosset, A.: *Rev. de Chir.*, 1903, xxviii, 694.

the aid of sutures and of the Murphy button. The results have not been encouraging. The greater amount of traumatism and the more protracted duration, together with the difficulty of making a perfect closure, have limited our success. The side-tracking method has been accomplished with the aid of the twine triangular stitch elaborated in this laboratory,¹² and with the aid of the Murphy button (Figs. 2 and 3). We have also made use of a special button (Fig. 4) designed (by N. W. G.) with the idea of causing an anastomosis with no further opening of the viscera at the time of operation than two needle-like punctures made after the serous surfaces have been coapted.

We have used during these procedures an apparatus for artificial respiration described in previous communications.¹³

It may be well here to present a diagram of it (Fig. 5). Ether has always been the anesthetic used. The loss of body heat was always considerable, but this, in part, has been overcome by warming the ether vapor before sending it into the air-passages.

It may be of interest here to describe in detail one type of procedure which has seemed to us particularly satisfactory. The animal was first placed under ether and the intralaryngeal canula inserted and held in place by a clamp, pinching the tube just oral to the hyoid bone. Through the mouth were introduced, first into the stomach and then into the esophagus, the halves of the button shown in the diagram. The esophageal portion, being the one with the needles attached, was introduced and guarded by a long alligator-jaw forceps. This was left in place while the chest was opened at the level of the eighth rib on the left side. The eighth rib was removed, stripping from it its periosteum, as suggested by Janeway. This gives a much better opening than the intercostal incision and makes it far easier to close the chest at the completion of the operation. The amount of air introduced was regulated by direct inspection of the lung. The seventh and ninth ribs were pressed apart by a self-retaining retractor and care was taken to protect the wound from contamination by a suitable covering. The diaphragm was opened in its tendinous aponeurosis, and the cardiac portion of the stomach drawn up through the opening into the chest and sutured in place to the edges of the diaphragmatic wound. This suture was a continuous one and several reefing stitches were taken in the stomach to one in the diaphragm. This close stitching of the stomach was found to be necessary on account of the tendency of the abdominal viscera to cause fatal prolapse into the chest. We were careful not to include the blood vessels of the stomach in our sutures. Feeling through the stomach wall, we grasped the first portion of the button, previously introduced, and drew it through the new diaphragmatic opening into this cul-de-sac. The forceps were then removed from the esophageal half, the surfaces of the stomach and esophagus brought in apposition and the two halves of the button pressed tightly together. They were held by the spring which engages the serrations at the side of the needles. This completed the intrathoracic part. The chest wall was then closed in layers. The operation here described consumed fifty-five minutes. At the end of ten days the

dog was given a lethal dose of ether. The stoma shown in the photograph was found present (Fig. 6).

We wish to express our appreciation to Mr. Arthur G. Sullivan for his assistance, to Drs. H. H. Janeway and N. B. Leggett for many valuable suggestions and to Dr. Robert M. Brown for his skillful mounting of the specimens.

152 West Fifty-seventh Street—264 West Fifty-seventh Street.

PRESENT STATUS OF SURGERY OF THE THORAX

AND THE VALUE OF THE SAUERBRUCH NEGATIVE PRESSURE PROCEDURE IN THE PREVENTION OF PNEUMOTHORAX.*

E. F. SAUERBRUCH, M.D.

Oberarzt and Chief of the Surgical Poliklinik, University of Marburg.
GERMANY.

When I was accorded the honor of an invitation to deliver an address before the Section on Surgery and Anatomy, of the American Medical Association, it occurred to me that it would be a suitable opportunity to present the modern methods employed for opening the thoracic cavity and to give you the results of many personal experiences obtained during the past four years in the use of my pneumatic cabinet. The inefficiency of the methods in vogue up to that time prompted me at the solicitation of my then chief, von Mikulicz, to endeavor to devise a method of operation which would obviate the dangers incident to thoracic surgery. I began by analyzing the methods used to prevent the occurrence of pneumothorax, especially the method of artificial respiration which had been used extensively and successfully in animal experiments for more than a hundred years, and, more recently, in an improved manner, by Matas and several others in the human being. I soon became convinced that this method possessed too many disadvantages ever to become practical. That led me to experiment with negative pressure, basing my work on the known principles of the difference between the intrabronchial pressure and the pressure at the surface of the pleura, about 7 to 8 mm. Hg, which serves to keep the lungs distended. If the normal relationship between these pressures is disturbed, as by opening the thorax, the lungs collapse and the result is a pneumothorax, which may be followed by serious disturbances, such as dyspnea, displacement of the diaphragm and heart, vagus reflexes, etc. Believing that these complications would not occur if the normal relationship of these pressures could be maintained, I was led to carry out the following experiments: The ends of a glass cylinder were sealed with gutta percha paper. One end was perforated by three openings, two small ones, and one large one; the opposite end was perforated by a single large opening. The animal was placed in the cylinder, so that the head protruded from one of the large openings, and the legs and lower part of the body through the other opening. I passed my hands in through the small openings, endeavoring to keep the cylinder airtight. Next, the cylinder was exhausted sufficiently to reduce the air pressure 10 mm. Hg; then I opened

12. Maury, J. W. Draper.: Studies from the Laboratory for Surgical Research, Columbia University, 1907, 1.

13. Green, N. W., and Maury, J. W. Draper.: Ann. Surg., Oct., 1907, xlv1, 544-548.

* Read in the Section on Surgery and Anatomy of the American Medical Association, at the Fifty-ninth Annual Session, held at Chicago, June, 1908.