


This article was downloaded by: [York University Libraries]
On: 02 February 2015, At: 01:42
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer
Street, London W1T 3JH, UK



Annals and Magazine of Natural History: Series 7

Publication details, including
instructions for authors and subscription
information:

<http://www.tandfonline.com/loi/tnah13>

VI.—Notes on the Cubomedusæ

F.S. Conant

Published online: 22 Sep 2009.

To cite this article: F.S. Conant (1898) VI.—Notes on the Cubomedusæ, Annals and Magazine of Natural History: Series 7, 1:1, 31-40, DOI: [10.1080/00222939808677920](https://doi.org/10.1080/00222939808677920)

To link to this article: <http://dx.doi.org/10.1080/00222939808677920>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands,

costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

secured on the 7th November, which ate 18 frogs, cast its skin on the 7th December, 1896, 13th January and 27th February, 1897.

"Tree-snake (*Dendrophis pictus*) ate 94 frogs up to 21st January, and died on the 22nd January. Cast its skin on the 2nd April, 6th May, 28th June, 27th July, and 29th October. A second specimen was procured on the 13th February, which ate 18 frogs.

"Russell's viper (*Vipera Russellii*), young, ate 67 mice, 2 small rats, 5 squirrels, and 4 frogs. Cast its skin on the 16th June, 31st July, 6th October, and 27th December, 1896.

"Russell's viper (*Vipera Russellii*) ate 27 rats and 5 squirrels. Cast its skin only once, *i. e.* 17th November, 1896.

"*Echis carinata*.—Two specimens were exhibited on the 29th October, and have up to date eaten only 8 frogs.

"The two aquatic snakes added were *Tropidonotus quincunciatus* and *Atretium schistosum*; the latter is called in Tamil 'Neer nagan,' *i. e.* water cobra, from its power of dilating its head. Both the snakes eat frogs and fishes. *Tropidonotus* laid 36 eggs on the 26th February, 1897."

VI.—Notes on the Cubomedusæ.

By F. S. CONANT*.

IN June and July of 1896, while the Marine Biological Laboratory was stationed at Port Henderson, in Kingston Harbour, Jamaica, two species of Cubomedusæ were found in unexpected abundance. Of these one was very much like the *Charybdea marsupialis* of the Mediterranean, the other presented characteristics midway between those of the Charybdeidæ and the Chirodropidæ—the two families into which Hæckel divides the Cubomedusæ—so that for it I have erected a new family under the name of the Tripedalidæ. The names that have been chosen for the two Jamaica forms are *Charybdea xaymacana* and *Tripedalia cystophora*.

A.—SYSTEMATIC.

Hæckel's classification, as given in his 'System der Medusen,' is as follows:—

* From the 'Johns Hopkins University Circulars,' November 1897, pp. 8-10.

CUBOMEDUSÆ (Hæckel, 1877).

Acraspeda with four perradial marginal bodies, which contain an auditory club with endodermal otolith-sac and one or several eyes. Four interradial tentacles or groups of tentacles. Stomach with four wide perradial rectangular pockets, which are separated by four long and narrow interradial septa. They belong to the subumbrella, and are developed from the endoderm of the stomach-pockets, so that they project freely into the space of the pockets.

Family I. *Charybdeidæ* (Gegenbaur, 1856).

Cubomedusæ with 4 simple interradial tentacles; without marginal lobes in the velarium, but with 8 marginal pockets; without pocket-arms in the 4 stomach-pockets.

Family II. *Chirodropidæ* (Hæckel, 1877).

Cubomedusæ with 4 interradial groups of tentacles; with 16 marginal pockets in the marginal lobes of the velarium, and with 8 pocket-arms, belonging to the exumbrella, in the 4 stomach-pockets.

(This family is represented in American waters by a species of *Chiropsalmus* found at Beaufort, N.C., and identified by H. V. Wilson as *Ch. quadrumana*. Before Claus's paper on *Charybdea marsupialis* was received at the laboratory, Professor Wilson had gone over precisely the same ground on *Chiropsalmus*, but with results so similar to those of Claus that, when the latter's article came, he did not finish for publication the notes and drawings he had made.)

To accommodate the second species found in Kingston Harbour another family must be added to the preceding.

Family III. *Tripedalidæ* (1897).


Cubomedusæ with 4 interradial groups of tentacles, each group having 3 tentacles carried by 3 distinct pedalia; without marginal lobes in the velarium; with 16 marginal pockets; without pocket-arms in the stomach-pockets.

Description of *Charybdea xaymacana*. (Fig. 1.)

Genus *Charybdea* (acc. to Hæckel): *Charybdeidæ* with 4 simple interradial tentacles, with pedalia; with velarium suspended, with velar canals and 4 perradial frenula. Stomach flat and low, without broad suspensoria. Four

horizontal groups of gastric filaments, simple or double, tuft- or brush-shaped, limited to the interradial corners of the stomach.

Fig. 1.



Charybdea xaymacana.

Reference letters in Figures.

af, furrow on inner surface of exumbrella underlying the adradial ridge on the outer surface; *afr'*, furrow on outer surface, setting off the area in which the sensory niche lies; *ct*, canal of the tentacle; *go*, gastric ostium; *ifr*, interradian furrow on outer surface; *n*, nerve-ring on the subumbrella; *pe*, pedalium; *ph*, phacellus; *pr*, proboscis; *r*, reproductive organ; *sn*, sensory niche; *su*, suspensorium; *sub*, subumbrella; *vc*, velar canals.

Species: Charybdea xaymacana.

Bell a four-sided pyramid, with the corners more rounded than angular, yet not so rounded as to make the umbrella bell-shaped. The sides of the pyramid parallel in the lower two thirds of the bell, in the upper third curving inward to form the truncation; near the top a slight horizontal constriction. Stomach flat and shallow. Proboscis with four

oral lobes, hanging down in bell-cavity a distance of between one third and one half the height of bell; very sensitive and contractile, so that it can even be inverted into the stomach. The four phacelli, epaulette-shaped, springing from a single stalk. Distance of the sensory clubs from the bell-margin one seventh or one eighth the height of bell. Velarium in breadth about one seventh the diameter of the bell at its margin. Four velar canals in each quadrant; each canal forked at the ends, at times with more than two branches. Pedalia flat, scalpel-shaped, between one third and one half as long as the height of bell. The four tentacles, when extended, at least eight times longer than the bell. Sexes separate. Height of bell 18–23 millim.; breadth about 15 millim. (individuals with mature reproductive elements). Without pigment. Found at Port Henderson, Kingston Harbour, Jamaica.

As may be seen from the above, *C. xaymacana* differs only a little from the *C. marsupialis* of the Mediterranean. Claus mentions in the latter a more or less well-defined asymmetry of the bell, which he connects with a supposed occasional attachment by the proboscis to algæ. In *C. xaymacana* I never noticed but that the bell was perfectly symmetrical. *C. xaymacana* is about two thirds the size given by Claus for his examples of *C. marsupialis*, which were not then sexually mature. It has 16 velar canals instead of 24 (32) as given by Hæckel, or 24 as figured by Claus. Difference in size and in number of velar canals are essentially the characteristics upon which Hæckel founded his 'Challenger' species *C. Murrayana*.

Description of *Tripedalia cystophora*. (Fig. 2.)


Genus *Tripedalia*: for the present the characteristics of family and genus must necessarily be for the most part the same. The genus is distinguished by having 12 tentacles, in 4 interradial groups of 3 each; velarium suspended by 4 per-radial frenula; canals in the velarium; stomach with relatively well-developed suspensoria; 4 horizontal groups of gastric filaments, each group brush-shaped, limited to the interradial corners of the stomach.

Species: *Tripedalia cystophora*.

Shape of bell almost exactly that of a cube with rounded edges; the roof but little arched. Proboscis with 4 oral lobes; hanging down in the bell-cavity generally more than

half the depth of the cavity, and at times even to the bell-margin. In the gelatine of the proboscis an irregular number (15-22) of sensory organs resembling otocysts, from the presence of which comes the specific name. Phacelli brush-shaped, composed of from 7 to 13 filaments springing from a

Fig. 2.

*Tripedalia cystophora.*

single stalk in each quadrant, or, rarely, from two separate stalks in one of the quadrants. Distance of the sensory clubs from the bell-margin about $\frac{1}{5}$ or $\frac{1}{4}$ of the height of bell. Breadth of velarium about $\frac{1}{6}$ the diameter of bell at margin; with 6 velar canals on each quadrant; the canals simple,

unforked. Pedalia flattened, shaped like a slender knife-blade, about half as long as the height of the bell. Tentacles at greatest extension observed $2\frac{1}{2}$ times the length of pedalia. Sexes separate. Height of bell in largest specimens (reproductive elements mature) 8 or 9 millim. Breadth same as height, or even greater. Colour a light yellowish brown, due in large part to eggs or embryos in the stomach-pockets. The reproductive organs especially prominent by reason of their similar colour. Found in Kingston Harbour, Jamaica.

It will be seen from the above that *Tripedalia* possesses two of the characteristics of the Charybdeidæ and two of the Chirodropidæ. The small size of *T. cystophora* is worthy of note in connexion with the fact that of the twenty species of Cubomedusæ given by Hæckel in his system only two are smaller than 20 millim. in height, and those are the two representatives of Hæckel's genus *Procharagma*, the prototype form of the Cubomedusæ, without pedalia and without velarium. While *Tripedalia* has both pedalia and velarium, it may be, perhaps, that its small size, taken in connexion with characteristics just about midway between the Charybdeidæ and the Chirodropidæ, indicate that it is not a recently acquired form of the Cubomedusæ.

B.—HABITAT.

The Cubomedusæ are generally believed to be inhabitants of deep water, which come to the surface only occasionally. Both of the Jamaica species, however, were found at the surface of shallow water near the shore, and only under these circumstances. Whether these were their natural conditions, or whether the two forms were driven by some chance from the deep ocean into the harbour, and there found their surroundings secondarily congenial, so to speak, can be a matter of conjecture only. *C. xaymacana* was taken regularly a few yards off-shore from a strip of sandy beach not ten minutes' row from the laboratory at Port Henderson. It was seen only in the morning before the sea-breeze came in to roughen the water and to turn the region of its placid feeding-ground into a dangerous lee-shore. Some of the specimens taken contained in the stomach small fish so disproportionately large in comparison with the stomach that they lay coiled up, head overlapping tail. The name *Charybdea*, then, meaning a gulf, rapacious, seems to be no misnomer. It is worth mentioning that the digestive juices left the nervous system of the fish intact, so that from the stomach of a *Charybdea* could be obtained beautiful dissections, or, rather, macerations, of the brain, cord, and lateral nerves of a small fish.

Charybdea is a strong and active swimmer, and presents a very beautiful appearance in its movements through the water. The quick vigorous pulsations contrast sharply with the sluggish contractions seen in most Scyphomedusæ. When an attempt is made to capture one it will often escape by going down into deeper water—as, indeed, do other jellyfish. Escape from observation is all the more easy by reason of the entire absence of pigment, except for the small amount in the sensory clubs. The yellowish or brownish colour usually stated as common in the Cubomedusæ is nowhere present in *C. xaymacana*.

The environment in which *Tripedalia* was found is still more unlike that in which Cubomedusæ have been taken heretofore. It was obtained in two localities, both of which were cut off from the main body of the harbour, and so from the ocean, by peninsulas and islands covered with a growth of mangrove. The water was shallow and discoloured with organic matter, and the bottom for the most part a black mud. The regions were virtually mangrove-swamps. It was under the shelter of the overhanging mangrove boughs, among their half-submerged roots, that the *Tripedalia* was found thriving in large numbers. Conditions more strikingly unlike those of the pure deep sea could hardly be imagined.

T. cystophora is as quick and vigorous a swimmer as the species of *Charybdea*. In one case as many as 110 pulsations were counted in a minute. Its brownish-yellow colour enables it to escape observation not infrequently in the discoloured water.

C.—ANATOMY.

An abundance of material of both species was preserved in several ways and has furnished subject-matter during the past year for investigation of the adult anatomy of the Cubomedusæ. The results have been embodied in a paper whose publication I hope will not be greatly deferred. Meanwhile the mention of one or two points may not be out of place.

The parts to which especial attention was given were the vascular lamellæ, or cathammal plates, and the nervous system. The vascular lamellæ, being definite areas where the two walls of the gastro-vascular system have come together and fused, are interesting because they show how the present structure of the gastro-vascular system of the Cubomedusæ has come about from a simpler undivided condition. It was hoped that their study might throw definite light on the affinities of the group. The results, however, were unsatisfactory, for while one part of the system of

lamellæ pointed strongly toward a relationship with such Hydromedusæ as, for example, *Liriope* (Trachomedusæ), another part, around the margin of the bell, gave evidence equally as good to the effect that the velarium was formed by the fusion of marginal lobes and that the view which considers it not homologous with the Hydromedusan velum is therefore correct. The question as to the affinities of the Cubomedusæ must wait until the development is known.

The nervous system has been described by Claus*, and more recently in greater detail by Schewiakoff†. With the latter's conclusions as to the structure of retina and vitreous body of the complex eyes on the sensory clubs I am unable to agree. Schewiakoff's conception of the structure is in brief as follows:—

a. The retina is made up of two types of cells, pigment and visual, which are figured as alternating regularly. The pigment-cells are cone-shaped; the visual are spindle-shaped, with their nuclei lying in the swollen central portion of the spindle at a lower level than the nuclei of the supporting pigment-cells.

b. From the visual cells extend rod-like processes into the vitreous body (which lies between the retina and lens), lying in canals in the vitreous body.

c. In the vitreous body separate cone-shaped streaks of pigment overlie the pigment-cells, which do not, however, form part of those cells.

d. Apart from these pigment-streaks and the rod-like processes of the visual cells, the vitreous body is structureless, probably a secretion of the pigment-cells.

The conclusions reached upon the same points by the study of the two Jamaica species are:—

a. There is not good evidence of two distinct types of cells in the retina—cone-shaped pigment-cells and spindle-shaped visual cells, with the nuclei of the latter at a lower level than those of the former.

b. The rod-like processes in the vitreous body exist, though not referable to a special type of cell in the retina.

c. The cone-shaped streaks of pigment in the vitreous body belong to the underlying pigment-cells, in fact are direct continuations of them, and at their distal ends they are prolonged into fibrous processes lying in canals of the vitreous body exactly like the visual rod-like processes of Schewiakoff.

* "Ueber *Charybdea marsupialis*," Arb. aus d. zool. Inst. d. Univ. Wien, Bd. ii. Heft 2 (1878).

† "Beiträge zur Kenntniss des Acalephenauges," Morph. Jahrb. Bd. xv. Heft 1 (1889).

d. The vitreous body, over and above these "visual fibres" and "pigment-streaks," is not a homogeneous secretion, but is composed of prisms of gelatinous refracting substance, each with a denser central fibre. The size of the cross-sections of these prisms corresponds with that of the cross-sections of the majority of retinal cells. It is probable, therefore, that they represent the outer ends of retinal cells.

From the above it will be seen that I find in the vitreous body two structures in addition to Schewiakoff's visual processes—namely, similar processes from the "pigment-streaks," and the prisms that make up the mass of the body. I am inclined to believe that the retina contains at any rate two, and perhaps three, kinds of cells, but that these are distinguishable only by that part of them which lies in the vitreous body, and are not distinguishable in the retina as two alternating types, as figured by Schewiakoff.

In another point also I am unable to agree with Schewiakoff. In one region of the sensory club is found a mass of peculiar large, closely packed, polygonal cells, which he affirms to be ganglion-cells, as against Claus, who considers them supporting cells. In my sections they show nothing whatever to suggest a resemblance to ganglion-cells. No processes could be made out, and the body of the cell consisted of a beautiful cytoplasmic reticulum with thickenings at the nodes of the rather coarse meshes.

The sensory organs in the proboscis of *Tripedalia* are simple cysts of ciliated epithelium lying in the gelatine. The long cilia support and keep in constant motion an otolith of ragged outline, in which no trace of cellular structure was found. This is the only case, so far as I know, of the occurrence among the Medusæ of a free unsuspended otolith in a ciliated sac.

D.—DEVELOPMENT.

Females of *Tripedalia* were found with stomach-pouches or pockets filled with developing ova, so that I am able to offer a few facts preliminarily on the development of the Cubomedusæ. The mature eggs pass from the ovaries into the stomach-pouches, and there develop up to the stage of free-swimming planulæ. In with them are found floating masses of cells, showing here and there a nematocyst, whose structure points clearly to an origin from the ovary also. These masses circulate about among the developing ova, and serve for protection and apparently also for nutrition, since the cells are found vacuolated in masses from individuals containing well-advanced embryos. All the eggs or embryos in one

individual are at the same stage of development. Gastrulation was not observed in the living material, but sections show a stage in which central cells lie clearly marked off from a surrounding ring of smaller surface-cells, and the probability is that these endodermal cells were formed by delamination. When the spherical mass is beginning to elongate into the ellipsoidal form of the planula a large migration of cells from the surface into the centre takes place, not limited to one pole. The relation of the central cells earlier seen and of these in-wandering cells to the definitive endoderm was not made out. The young are set free from the parent as ciliated planulæ, having pigment spots on the posterior end. The planulæ swim about actively on the surface for a day or two, their motion being both translation and rotation. Then they gradually lose the forward motion and rotate on their own axis in one spot. At this point they settle down and the pigment spots which were on the posterior surface are found to migrate into the interior. In a day or two they bud out two tentacles, and shortly afterwards two more, though young hydrazes with three and five tentacles were by no means uncommon. In this condition they lived for three weeks in the aquaria without undergoing further development.

Search for scyphistomas in the region where the jellyfish were found was fruitless. The youngest specimens taken in the tow were of practically the adult structure, so that the important intermediate stages remain unknown. The youngest jellyfish were interesting, however, in that only the middle tentacle in each group was developed, and that the sensory clubs were situated almost on the margin, and were not yet enclosed in niches.

VII.—*On new Mammals from Western Mexico and Lower California.* By OLDFIELD THOMAS.

THE British Museum has received from Mr. W. W. Price two collections of mammals—one from Sinaloa, W. Mexico, obtained by Mr. P. O. Simons, and the other from the Cape region of Lower California, collected by Mr. D. Coolidge.

In the former collection occur the remarkable pigmy skunk with uninterrupted dorsal lines (*Spilogale pygmaea**) exhibited recently at the Zoological Society, and the jaguarondi described below. The remainder of the collection consists mainly of

* Diagnosed in Abstract of Meeting of Nov. 30. Published Dec. 1, 1897.