spots were found, the spleen was not enlarged, and the serum reaction for typhoid was twice negative. The temperature at first fell and the abdominal tenderness and distension abated. The temperature then gradually mounted again; some tenderness was found below the right costal margin and in the right lumbar region; the upper level of liver dulness in the axilla was two ribs higher than it should be. Aspiration of the chest in the axilla was negative, but incision of the right loin far back evacuated pus from the right kidney well. The liver dulness gradually descended and an uneventful recovery ensued.

To enter into a detailed description of the characteristics and limits of abscesses in the lesser sac of the peritoneum and those in the pelvis would require more time than is available for this address.

The treatment of intraperitoneal abscess is essentially surgical. When there is reason to suspect one of these collections it should be drained. The process of localisation takes from seven to ten days or more. The dangers of secondary diffuse peritoneal infection, of infection of the pleural sac and lung, and of portal pyæmia are always present.

The fallacies which attend the use of the aspirator needle for the diagnosis of subphrenic collections I have already mentioned. The needle, if used, should only be introduced after a close consideration of the history and careful weighing of the clinical signs. It may give accurate information as to the nature of the pleural exudate and quite unreliable indications as to the abdominal condition. It should, needless to say, never be used for puncture below the ribs, nor yet below the diaphragm on the left side. The presence of the liver on the right is in some measure a protection. In a doubtful case where the needle has failed, as it often does, or the patient's progress and symptoms point to suppuration, the boldest course, of exploring by surgical incision, is the safest and most successful in the long run. The most suitable incision varies with the position, size, and apparent cause of the collection.

An extensive right subphrenic abscess, caused by recent rupture of a gastric or duodenal ulcer, is usually explored first from the front on account of the causal lesion. Should the collection be localised posterior transpleural drainage is usually adopted. The ascending subphrenic abscess due to appendicitis, often associated with a collection in the right renal well, may be treated by an incision high up and far back in the flank, but should the subphrenic collection be of any magnitude it is unsafe to rely on the flank incision alone, and better to open the subphrenic space higher up by the transpleural route, steps being taken to shut off the pleural cavity. In one intractable case, complicated by empyema, the condition, which appeared hopeless, was successfully treated by Mr. C. S. Wallace, who excised the ninth rib from the posterior axillary line forwards to the costal margin, a procedure which afforded excellent drainage of the pleura and the whole subdiaphragmatic space.

The left subphrenic abscess is treated on the same lines as the right. The anterior incision may be necessary for the exploration of a perforated stomach, but for a localised collection of pus far back around the spleen the transpleural route should be taken. The locality of the abscess is best exposed by resection of rib, not higher than the seventh.

A point worth mentioning is the position in which patients suffering from appendicitis should be nursed. There is much to be said for the Fowler position, the back being elevated by pillows and the nates supported by a bolster placed transversely beneath the thighs and tied to the bed-frame. This position has the merit of preventing ascending infection, but the state of the pelvis should not be lost sight of, since it favours drainage into this cavity.

Cases of intraperitoneal inflammation well repay careful watching and continued efforts to limit the infection. The first operative procedures are not always successful, but this must not discourage further attempts. Success may only be attained by persistent efforts to reach and drain the various infected localities.

Devonshire-place, W.

THE APOTHECARIES' HALL OF IRELAND.—The quarterly examinations will take place on the following dates:—First professional, Monday, April 18th; second professional, Thursday, April 21st; third professional, Monday, April 25th; and final professional, Wednesday, April 27th.

THE TREATMENT OF PSORIASIS.¹

BY E. G. GRAHAM LITTLE, B.A. CAPE UNIV.,
M.D. LOND., F.R.C.P. LOND.,

PHYSICIAN, WITH CHARGE OF THE SKIN DEPARTMENT, TO ST. MARY'S HOSPITAL AND TO THE EAST LONDON HOSPITAL FOR CHILDREN AND DISPENSARY FOR WOMEN; LECTURER ON DERMATOLOGY, ST. MARY'S HOSPITAL MEDICAL SCHOOL; MEMBER OF THE SENATE OF THE UNIVERSITY OF LONDON.

PSORIASIS is one of the most ancient of skin diseases. Naaman the Syrian, commonly called Naaman the leper, probably in reality suffered from this, and the ages have revealed no remedy comparable in swiftness and certainty with the immersion in the River Jordan, which we are told cured him. "Lepra" was the name for certain clinical types of psoriasis until quite recently; Bateman's Atlas, published in 1840, has a plate (No. VII.) of "Lepra vulgaris, the common leprosy," which is certainly psoriasis as we know it. The "leprosy" of the Middle Ages must have been largely composed of psoriasis, and it is sad to reflect how much avoidable misery the victims of this disease must have endured from that unhappy confusion. The sum of suffering for which psoriasis is responsible still remains very great, for it is in our climate, next to eczema, the commonest of skin diseases.

The frequency of its occurrence in a general dermatological clinique may be estimated from the following figures. Of 11,500 cases of general skin diseases recorded in my department at St. Mary's Hospital, 465 were instances of psoriasis—i.e., almost exactly 4 per cent. Dr. Radcliffe Crocker's statistics quoted in his text-book (718 in 10,000 cases) give a much higher percentage; the difference in the two sets of figures may perhaps be accounted for by the fact that in my records care is taken to exclude the entry of the same individual case more than once, recurrent attacks not being recorded.

Psoriasis was included in Dr. Crocker's six diseases of the skin with which every practitioner should be quite familiar. A detailed exposition of some forms of treatment may, therefore, be acceptable to you, for its management requires the utmost patience and resourcefulness on the part of both the patient and the medical practitioner. Inasmuch as its causation remains still completely obscure, it is impossible to devise a rational treatment which would aim at removing the cause, and we are obliged to be satisfied with the teachings of experience as to what remedies are most successful in removing the eruption at any particular time. The pendulum of practice has swung to and fro between the claims of internal and external medication and a combination of both. It may be said that the present methods of treatment are more directed to external than to internal therapeutic measures, the long abuse of arsenic in this disease having produced a reaction, which has perhaps gone beyond its just limit, against all internal treatment.

Diagnosis.

A few observations on diagnosis may fitly precede the consideration of therapeutics, which must rely upon diagnosis. Psoriasis is usually one of the least difficult of diseases of the skin to recognise, since it follows such a characteristic and typical distribution and course. Cases, however, in which these are anomalous are not infrequent, and attention to certain points will be of essential aid to the practitioner in forming his conclusions.

Character of the initial lesion.—This is always a papule in its earliest stages, of conical shape, and of the same colour as the surrounding skin, from which it is, however, quite appreciably raised; later the papule becomes pink and enlarges, but retains an outline which is circular, not polygonal as in lichen planus. It is also pinker than the papule of lichen planus, in which a blue tint is added to the red, and than the papule of syphilis, in which a brown shade mingles with the red. What has been described as the "bleeding test" may be applied to the papule at this stage; it consists in gently removing the uppermost scale either with the finger-nail or, better, with a small scalpel; a pink, shiny surface comes into view, and if further removal of epithelium is effected by scratching or otherwise, a number of minute bleeding points are seen. The explanation of this phenomenon probably lies in the histology of psoriasis; the

¹ Paper founded on a lecture delivered at the Medical Graduates' College and Polyclinic.

superficial blood-vessels are dilated, and the increased vascularity accounts for the pinkness of the surface. In the accompanying diagram the position of the first and second hypothetical scratches is indicated by the interrupted horizontal line; the second, by taking off the tops of the papillæ exposes the bleeding blood-vessels from which blood exudes more freely by reason of their dilatation. The papule rapidly enlarges in some cases to form a scaly patch. The scalliness may become enormously exaggerated, as in the so-called psoriasis rupioides, and is always in excess of the scalliness of

Papule of psoriasis (semi-diagrammatic). A, Zone of adherent scales (Parakeratosis). B, Zone of hypertrophied epidermic downgrowth (Acanthosis). C, Superficial plexus of vessels with papillary branches. 1, Position of initial scratch, scales removed. 2, Position of later scratch, papillæ with dilated vessels exposed.

any other skin disease. The scales are white, laminated or stratified, and easily detached from the skin as silvery flakes.

The *distribution* is, in adults, so regular as to afford material aid to diagnosis. The extensor surfaces of the limbs, and especially the skin covering the joints, together with the scalp, may be said to be the typical sites for the eruption to first appear. In children, however, the limbs are quite often free, while the trunk is chiefly affected. The hands and face are generally spared, and the mucous membranes are never in any way attacked. The nails are sometimes implicated by the disease; they become thickened, the free edges brittle, and the enamel of the surface dull; often minute depressions, like the markings on a thimble, appear on the nail, and are very characteristic. The hair is seldom in any way affected; extensive psoriasis of the scalp may coexist with a good head of hair; occasionally, if the scalp is left untreated for long periods, the hair becomes thinned and falls off. The history of family inheritance of the disease and the chronicity and seasonal recurrence of the eruption are features so frequent in psoriasis as to afford when present strong suggestion of this diagnosis.

Treatment.

In considering in detail the therapeutic measures to be adopted in any individual case much will depend on the question whether the patient can remain in bed or is obliged to go about the ordinary avocations of life during his treatment. I propose to deal with treatment therefore under two heads: (1) what may be called ambulatory psoriasis, and (2) "lectual," which is good Latin for "confined to bed."

Ambulatory psoriasis.—Here the patient has to confine his treatment to his sleeping hours and must remain fairly presentable in his person in order to allow of his continuing his daily routine of work. It is important to keep the latter requirement in view and to avoid the risk of disfiguring the exposed parts of the body or to know means of concealing the disfigurement when it is inevitable. The face and hands, and in women who wear evening dress the shoulders and upper part of the chest, will thus have to be specially considered. Local treatment is often difficult under these circumstances, and in such cases internal medication has its most legitimate sphere. But with some ingenuity and care local measures, even including ointments which discolour the skin, may be used even in these positions.

Of all the local applications employed in psoriasis none can compare in efficiency with chrysarobin, which, in spite of all its drawbacks, I never willingly withhold. The methods of applying this are very numerous. Chrysarobin is soluble in chloroform, ether, and benzene, and a saturated solution in the first of these with guttapercha is known as traumaticin. The guttapercha addition forms an adhesive paint which may be used for small isolated patches, especially

on the face, but it is the least efficacious, if the most agreeable, method of using this drug. A far better way is to add chrysarobin powder to a fairly adhesive ointment base, such as the following: kaolin levigati $\mathfrak{Z}$ ii., amyli $\mathfrak{Z}$ iv., lanolin. $\mathfrak{Z}$ iv., paraffin. fluid. alb. $\mathfrak{Z}$ ii.; to this is added chrysarobin xv.-xxx. gr. The ointment thus resulting may be applied to the face and the forehead where the hair joins the skin—a very frequent position for obstinate patches of the disease. It should be used at night and removed in the morning either with vaseline or with benzene; the latter is especially useful on hairy parts, and all trace of the ointment should be removed before washing the parts with soap, which in combination with chrysarobin produces purple discolouration of the hair.

The patient must be warned of the danger of conjunctivitis from contact of the conjunctiva with chrysarobin. On the hands the most convenient covering is a pair of cotton gloves, such as Tommy Atkins wears, which may be used again and again. The more the ointment soaks into the glove the more efficacious it becomes. On the covered parts of the body the adhesive ointment mentioned can be left *in situ*, and the proportion of chrysarobin increased to half a drachm to the ounce; it will not come off with mere rubbing, so that the bath need not be prohibited.

For small patches, especially on the face and for resistant patches elsewhere, a very convenient form of application is by the ointment stick—the "*Baculum chrysarobini*" of St. Mary's Hospital Pharmacopœia: Chrysarobin, 3; white wax, 2; lanolin, 5. These sticks may be made of any required calibre according to the dimensions of the patch to be treated; a convenient thickness is that of a lead pencil. The stick is rubbed on the patch for a few seconds, and the smear thus produced is left for as long as may be possible. The cleanliness of the process and the ease with which the stick may be carried about if wrapped in tin-foil, make this a very suitable form for patients while travelling or on the move.

The ointment used for parts which have to be exposed should always be a weak one containing not more than 15 grains of chrysarobin to the ounce. When inconvenient redness results, as it often does even with these weak formulæ, it may be disguised quite effectively by painting over the reddened area, either with the finger or a brush, a very thin film of Pellanthum, either plain or containing 3 per cent. ichthyol, or a combination of the plain and ichthyolated preparation may be required; when mixed, as one mixes colours on a palette, in suitable proportions, the tint of the surrounding skin may often be very closely matched and the redness quite obliterated. The film may be at once removed by wetting with water, and it is quite harmless to the skin. Before pressing out the contents of the Pellanthum tube it is well to ensure mixture within the tube by gently stirring with a needle; the material exuded should be of a uniform creamy consistence. When the film is too thick it may be thinned with a little water.

Where very obstinate small patches (but not on exposed parts) resist the treatment described, a good method of clearing them (suggested by Dr. A. Whitfield) is to paint the surface with saturated solution of chrysarobin in chloroform, allow the film to dry, and apply over it and a little beyond the margin of the deposit left after evaporation of the chloroform some Leucoplast plaster which is left in position until it comes off, in three or four days, when the process may be renewed.

The disadvantages of the treatment by chrysarobin are not to be ignored, but I am convinced by much experience that most patients are willing to put up with its drawbacks for the sake of the greater rapidity and certainty of its action. It is true that the personal and bed linen become disagreeably stained and that the aspect of the ointment is unæsthetic, but these are minor drawbacks. A more serious one is the risk of producing conjunctivitis when the drug is used in the neighbourhood of the face; this may generally be completely avoided with care, the essential precaution to be taken being to keep the ointment from being conveyed to the eye either by the hands or by continuity with the parts of the face so treated. Where the forehead is affected it is well to apply chrysarobin under a bandage which will keep the ointment in the position desired, for it must be remembered that the movements of the head on the pillow may spread the ointment on the pillow-case, which may then

come in contact with the eye. Or more comfortably a face-mask of lint may be used, with apertures made for the nose, mouth, and eyes, and kept in position by tapes going round the head. In spite of all precautions conjunctivitis sometimes will occur and the drug must at once be omitted and the eye treated with astringent lotion—e.g., acid. borici gr. vi.; sod. sulpho-carbolati gr. x.; aq. rosæ ad ℥ i.

Psoriasis of the scalp.—Since chrysarobin stains the hair a purple colour it is seldom possible to use it on the scalp, and one has to be satisfied as a rule with less efficacious applications which do not stain hair or skin. Inasmuch as the hair of the head tends to increase the adhesion of the scales, it is particularly important to remove the scales before using any treatment. This may be done with soft soap, of which a pleasant and cleanly method of use is the following. A lotion consisting of equal parts of sp. vin. rectif. and sapo mollis with 15 minims to the ounce of liq. carb. deterg. is rubbed into the scalp with a small sponge, left to soak for half an hour, and washed away with warm water. This may with advantage be applied every night for a week preceding the application of any ointment to the scalp. When the scales have thus been in some measure removed the following ointment may be used nightly, the head being washed about once a week with the soap lotion mentioned: Sulphur. gr. xv.; camphoræ gr. xv.; liq. picis carb. ℥ xxx.; acid. carbol. gr. xv.; resorcin gr. xxx.; adip. benz., paraff. moll. aa. ℥ ss. This treatment is quite pleasant when the hair can be kept short, but becomes difficult in women, who dislike the greasy effect on their hair. For such cases an ointment which can be used at night but readily washed out in the day is better. The following formula suggested by Brocq I have found very useful: Ol. cadin. pur. 50 grammes; ext. quillaia 5 c.c.; glycerini amyli 45 grammes; essentia caryophylli 2 c.c. Mr. Andrews, the senior dispenser at St. Mary's Hospital, has kindly furnished me with the following notes as to the preparation of this formula: "The extract of quillaia is made by taking 100 parts of quillaia bark; the bark is powdered and is damped with alcohol (49 per cent.) and packed in the percolator. More spirit is poured in till it begins to drip from the bottom. Percolation is then stopped and the mixture is left to stand for 24 hours and then percolated. The first four-fifths of percolate is collected and put aside. Go on percolating with more spirit (49 per cent.) until the drug is exhausted. Evaporate this weak percolate to a soft extract and dissolve this latter in the reserved four-fifths and make up to 100 with more 49 per cent. alcohol. 5 cubic centimetres of this extract are sufficient to emulsify the oleum cadini. The method of mixing it is as follows. The extractum quillaia prepared as above is added to the oleum cadini and shaken; the emulsion thus resulting is added a little at a time to the glycerinum amyli in a mortar and rubbed until thoroughly mixed, and, lastly, the oil of cloves is added." This forms a not very unpleasantly smelling ointment which can be rubbed into the scalp and readily removed by washing. It does not stain the hair or the linen permanently. It should be rubbed into the scalp every night and washed out in the morning. Women, however, may use it less frequently as their hair is difficult to dry.

Another formula with oil of cade, somewhat less difficult and cheaper to make, is the following. It is, however, much more evil-smelling than the formula just cited, but is equally efficacious and easily removed by washing. Lard, 50 grammes; potash, 17.50 grammes; aq. dest., 150 grammes. Dissolve potash in water, add to lard previously put in a weighed water bath; mix thoroughly; add 10 cubic centimetres of absolute alcohol. Heat until a soap is formed; add to this 20 grammes of lanolin; concentrate the whole until the weight is 150 grammes; stir till cold and keep in an air-tight jar. Of this base take 20 gr.; add resorcin, 1 gr.; sulph. precip., 1 gr.; and ol. cadini, 5 gr. Mix well.

The disagreeable odour of these remedies may in some cases render their use impossible, and one is obliged to resort to less offensive measures. Of these, mercurial ointments are probably the best; an ointment of calomel 1 part to 3 parts lard; or the ammoniated mercurial ointment with gr. xv.—xxx. of resorcin, or acid. salicylic. may be tried. The possibility of salivation with the use of these strongly mercurial preparations must be carefully borne in

mind and their administration stopped on the appearance of any symptoms of mercurialisation.

Where chrysarobin cannot be tolerated.—Occasionally in very acute and itchy cases the application of chrysarobin, even in minimal doses, is intolerable; but it is surprising how seldom it is really necessary on this account to withhold it. Certain patients undoubtedly have an idiosyncrasy of intolerance, and in such cases probably the next best substitute is oil of cade, which may be given in varying degrees of strength, from purity to a proportion of 1 of oil of cade to 8 of olive oil. Oil of cade may also be added to unguentum acidici salicylici in the proportion of one or two drachms to the ounce, and this is a useful formula for very scaly cases. In the more erythematous and itchy cases the oil of cade may be usefully combined with glycerine of starch in the proportion of a quarter to a half of oil of cade, with a few drops of essence of peppermint (about 1 per cent.). In the same manner as was mentioned in speaking of the chrysarobin treatment, the identical dressings should be kept in use throughout; the more they are soaked the better they act.

Oil of cade may also be used in the form of *baths*, and these are especially useful in acute cases. Balzer's formula has been much praised and is an excellent method. Add the yolk of an egg to 50 grammes of oil of cade in a mortar, a sufficient quantity of extract of quillaia to make an emulsion, and 250 grammes of distilled water. This mixture is then added to the bath and the patient is immersed in the water for half an hour.

Another method of using tar baths which I have found useful in acute itchy psoriasis is thus applied. The patient is painted over with the following lotion:—Picis carbonis, 3; sp. vin. meth., 2; ether, 1 (balneum picis, St. M. H.). This is allowed to dry on the body, and the patient is then immersed in a warm bath for half an hour, after which he is carefully dried with warm towels. The same formula may be used with the following addition: Lotion for baln. picis, ℥ vi.; bals. Peru, ℥ vi.; coumarin, gr. xxxvi.; musk, gr. xii. This is painted on the psoriasis patches and left on at night. Salicylic acid (gr. xv.—xx. to ℥ i.) increases its efficiency. The strong odour, which, however, is not altogether disagreeable, makes this method not always easily applicable.

X rays.—Very chronic resistant patches of the disease may be favourably influenced by exposure to X rays for short periods. My experience with this treatment is small and not altogether encouraging, the disease having returned in the site treated very soon after the cessation of treatment.

Other methods of treatment:—

Light baths.—Hyde of Chicago promulgated recently an interesting theory that psoriasis was due to the deprivation of the skin of exposure to sunlight, instancing in favour of this view the comparative rarity of the appearance of lesions on the face and hands, the absence of psoriasis in black races who expose their bodies very largely to the sun, and the improvement of actually diseased persons on treatment with prolonged sun baths. In the great Republic it would appear that tracts of country still remain where man can fish and shoot and live in *naturalibus*, and Hyde reports good results with this resumption of a fig-leaf costume—unhappily, only possible in Arcadian climes. In the only way in which we can imitate this treatment, the subjection of the body to electric light baths, it is probable that much of the effect is due to the heat as well as the light. Turkish baths or Russian baths (dry heat and vapour) are also especially useful in chronic cases where the patient is obliged to lead a sedentary life and rarely perspires freely and in cases where itching is a troublesome symptom. It is probable that one reason why psoriasis usually does better in warm climates than in cold is because sweating is more free, and it should be the object of a patient with chronic psoriasis to sweat all over at least once a week, either through exercise, which is naturally the preferable way, or by means of a hot-air or vapour bath. The home cabinets, of which there are many kinds on the market, offer a quite convenient means of taking these baths and form a useful adjuvant of the ambulatory chrysarobin treatment detailed above.

Spa treatment.—Psoriatics are usually sent to spas where arsenical waters are obtainable, and the treatment consists essentially in the internal action of arsenic. But prolonged immersion in hot water alone seems to be beneficial in certain types of the disease. At Leuk, for example, the psoriatic patients are immersed for several hours a day in the natural

warm spring water of this spa. I have noted good results from the treatment by the Aix-les-Bains douche-massage routine, and Dr. Stanley Rendall has assured me that he is convinced of its efficacy in this disease. Probably the sedative effect of the routine of these watering-places has a beneficial influence on the general health, and thus indirectly on the eruption. The effect of sea-bathing is curiously variable. On the whole, in my opinion the balance swings rather in the direction of harm than good, and the salt water unquestionably increases the irritation of the itchy types. But certain individual cases have apparently benefited by sea-bathing, which is not uncommonly actually prescribed.

Local psoriasis.—In all instances where the disease is very acute and extensive, in many where this is not the case, but the patient is desirous of being speedily cleared of his eruption, and whenever it is attempted to produce a final cure, confinement to bed is a most essential part of the treatment. Lately Dr. Norman Walker has suggested the application of chrysarobin all over the body, not merely on the diseased surfaces, in patients who can be kept entirely in bed. I have tried this method in several cases and am satisfied of its value. The point aimed at is the desquamation under chrysarobin of the whole surface and the treatment is kept up until this result is achieved. The patient is wrapped from the neck to the toes in lint thickly smeared with an ointment consisting of half a drachm of chrysarobin to the ounce of vaseline and the lint is kept in position by bandaging. Fresh ointment is applied to the same lint every day until the skin begins to flake off everywhere, the body becoming deeply stained. The psoriasis patches usually resist desquamation longer, and the proportion of chrysarobin may have to be increased on these areas. To cure the actual disease it is necessary that desquamation should occur. When this has happened, and not before, the patient receives a hot bath, fresh clothes are provided, and it is as well to sterilise all the clothes worn previously to the treatment. In using this method the face and scalp cannot generally be included, and these may be treated as mentioned under the heading of ambulatory psoriasis. But the forehead may with careful bandaging be treated by the continuous method, and the scalp in those persons who will sacrifice their hair may with advantage be likewise soaked in the ointment. The average time of detention in bed is four weeks, and the freedom from recurrences may be so prolonged as to amount to actual cure.

I am of opinion that in a first attack, and especially in young patients, it is advisable to attempt this radical method and give the patient the best chance of checking the disease. It is a matter of common observation that the more entirely the last vestige of psoriasis is removed from the body the more protracted are the intervals of freedom between the recurrences: an absolutely complete removal of eruption would probably be attended by an absolutely complete cure. No untoward results have as yet been noted in the course of this treatment. The main precaution to be taken is to avoid contact of the eyes with the ointment, an accident which may easily happen, especially in young people at night. Masks made of lint, with openings for eyes, nose, and mouth, and kept in position with tapes tied round the head, will generally prevent this accident. Similarly, the genital organs, especially of female children, should be protected from contact with the chrysarobin dressings.

Internal treatment in psoriasis.—In no disease has the abuse of arsenic, so commonly and so unhappily regarded as a panacea for all skin diseases, been so general as in psoriasis, and there is undoubtedly a reaction now against arsenic as well as other internal remedies. There is no doubt that in certain cases, in which this result cannot be, however, predicted, it acts with great effect, apparently being alone responsible for the clearance of an eruption. This is especially the case in psoriasis of children and in first attacks. But it cannot be regarded in any sense as a specific for the disease, as mercury, for example, is in syphilis, and the use of arsenic does not prevent the appearance and the spread of an eruption even while the drug is taken in full doses. The danger of giving a patient with an essentially chronic disease like psoriasis a medicine containing arsenic—for the taking of the medicine may become as chronic as the disease—must be borne well in mind. My old teacher, the late Dr. John Cavafy, never felt happy in giving arsenic to out-patients, and there is much to be said for his caution.

More reliance is usually to be placed on external than

internal remedies; of the internal remedies arsenic, in spite of its uncertain effect, remains the least uncertain. The drug should be pushed rapidly and stopped if the effect is not produced; from 5 to 15 minims may be given thrice daily, preferably in liquid form, well diluted, and immediately after meals so as to avoid the irritation which may result from its introduction into an empty stomach. Fowler's solution is probably still the best method of administering it. Certain organic compounds of arsenic, such as the cacodylates, and later the arsenates, have had a certain vogue, which is apparently not likely to be permanent, the effects being singularly uncertain and irregular. In the present state of our knowledge the substitution of atoxyl, the most widely used of these preparations, for the older forms is not encouraged by some very serious results which have been recorded of its action. It is difficult to say how far the natural springs containing arsenic in solution owe their effects to the arsenic or to other bodies, including radio-active elements which are found in association. The amount of arsenic as estimated by careful analyses in most of those waters is extremely small and their effect is probably a very mixed one.

Besides arsenic, salicin, mainly through Dr. Radcliffe Crocker's advocacy, has been much used, and with apparently very irregular results. I have personally not been persuaded of its efficacy, and I have given it an extensive trial in doses of 20 grains three times a day, prolonged for many months. Copaiba resin, given in cachets, is very strongly praised by my friend, Mr. Arthur Shillitoe, as efficacious in clearing an eruption. Potassium iodide in very large doses—from 60 to 600 grains per day—has been recommended, chiefly by Scandinavian writers. Thyroid extract in a very few cases, the selection of which seems capricious, acts well, but it is too dangerous a drug to give in large doses as a routine treatment in a chronic disease.

Diet and general management.—The psoriatic is commonly a fresh-coloured robust person. Sir George Humphry indeed described psoriasis as the "hall-mark of health." Many subjects of this disease are, however, nervous, highly-strung persons, and indeed a large body of opinion favours a "neurotic" theory of causation. It is notable that outbreaks of eruption will often coincide with some mental stress, and psoriasis is a common disease in asylums for the insane. The mischievous effect of stimulants in promoting an eruption has often been noted. Long hours of sleep and "the simple life" in general, with plain diet, are the best regimen. It has been thought that vegetarianism is beneficial and Bulkely in particular urges the importance of this combined with a large intake of milk. Opinion, however, is very divided upon the question whether diet really exerts any influence at all, and I am personally no advocate of any special feeding in psoriasis; restriction of alcohol is probably more important than restriction of meat, which is so commonly insisted upon. It ought to be borne in mind by the practitioner that it is nearly always possible to disperse an eruption even if he cannot cure the disease, and the patient must often be helped and encouraged to persevere with treatment by the confident assurance that relief at any rate is within his reach and that the interval of freedom from recurrences will be lengthy in proportion to the completeness of the dispersal of the actual eruption.

Wimpole-street, W.

THE MANCHESTER CONSUMPTION HOSPITAL.—

It is now 35 years since the Manchester Hospital for Consumption and Diseases of the Throat and Chest was founded. It began in a small way, and when Sir William Crossley became chairman 30 years ago there were six beds at Bowdon. Now there are 140 at the Delamere Sanatorium, for which Manchester is indebted to Sir William Crossley, and at Bowdon. Still, however, the call is for more beds, as patients are kept waiting so long that the time for successful treatment is sometimes passed before they can be admitted. It is proposed to add 12 beds at Delamere, where there is room for more if the funds could be found. And the cost of a bed, which in the early days of the institution was £1000, is now only £100 or even less. Here, then, is another opportunity for the rich and charitable. The Lord Mayor, Mr. C. Behrens, who has been for many years the honorary secretary of the hospital, was in the chair at the annual meeting on March 14th, and made an earnest appeal for more funds.

ON A CASE OF HEART-BLOCK.¹

BY J. G. EMANUEL, B.Sc., M.D., B.S. LOND.,
M.R.C.P. LOND.,

HONORARY PHYSICIAN, QUEEN'S HOSPITAL, BIRMINGHAM.

THE case I wish to bring under notice is that of a tailor, 56 years of age, a small man, 5 feet 1½ inches in height, and weighing 7 stones 1 pound. He has rather a waxy complexion, but there is generally a slight flush on each cheek, that contrasts with what would otherwise be a pale face. He appears to be a feeble little man, lacking physical power and force. On the other hand, he attends regularly to his business, and this necessitates his standing about all day. He has been married 26 years and has had a large family. Seven are living, six died at birth, and there was one miscarriage. His family history is good; the father is living, aged 81 years, and the mother died when 74 years old; one sister died from phthisis, aged 37 years; and four brothers and three sisters are living.

The patient had rheumatic fever when he was 20 years of age, evidently a severe attack, for he was in bed for six weeks. He states that he got up too soon, and at any rate he had to return to bed for a further period of six weeks with an attack of pleurisy. Ever since this time his left side has been weak and he feels pain in the side on slight exertion. For over 20 years he has been subject to giddy feelings and to some sensations which he calls "turns," lasting a second or so and coming on at any time, even while in bed. During these last he says "he loses himself for a second," and he sometimes flushes—he never becomes pale. About Christmas in 1908 he suffered from very bad "giddy turns" and was compelled to keep in bed. During these attacks he felt that he was sinking away, and on one occasion he thought he was dying. On one day he had as many as a dozen in an hour and a half. He describes himself as sometimes having pains over the præcordium which catch his breath, but only last a second. These pains he distinguishes from other pains in the chest which go through to the back in between the shoulders, and which he himself puts down to indigestion. Whether the præcordial pains are anginal or not, it is difficult to say; they never shoot into the neck or down the arm. For many years he has been short of breath on exertion.

Twelve months ago the patient was first brought to me by Mr. L. Kirkby Thomas, as for some weeks he had been losing strength and energy and the "giddy turns" had been exceptionally severe and harassing. From this time until Easter he had to give up his work and he spent most of the time in bed. Before this he had never had to give up his occupation, although he had often been obliged to stay at home for a day or two. He has now been very comfortable for six months and able to go to business every day. His summer holiday of four weeks was spent at St. Anne's, when he walked every day a number of miles in short and easy stages. His digestion, however, is not good and he has to be careful in his diet.

On examination the cardio-vascular system alone presents any abnormality. In point of size the heart is normal, being if anything rather small. There is no absolute cardiac dulness to be detected in consequence of the emphysematous condition of the lungs, and the cardiac impulse is neither visible nor palpable; this may possibly be due to the same cause. On auscultation the heart sounds are most clearly audible at the apex, but at all the areas they are very weak. The first sound is accompanied by a short blowing systolic murmur, the second being free from murmur. On feeling the pulse one is struck by its extreme slowness, while the force and volume are considerable. On five different occasions during the past 12 months I have found the rate to vary within very narrow limits, being 42, 31, 36, 40, and 42 per minute. The rhythm is not quite regular, but the force of each beat is very constant. The heart sounds are synchronous with the pulse, and there are no sounds audible in the long intervals between the beats. The blood pressure is high. On Nov. 4th, as measured by the manometer, it was equal to 180 millimetres of mercury. The radial arteries are thickened and the brachials pulsating and tortuous. There are no signs of cyanosis or of oedema of the extremities. The urine is free from albumin and sugar, and there is no enlargement of the liver or spleen. The pupils are equal and react to light and accommodation. The knee-jerks are exaggerated.

It will be seen from the above account that while in this case the history and symptoms would seem undoubtedly to indicate the existence of a severe heart lesion the physical signs show no sufficient evidence of this. The heart is small, the pulse is full and strong, and though there is a systolic murmur audible at the apex, this in itself would not be sufficient to account for the frequent syncopal attacks and persistent dyspnoea on any but the slightest exertion. The most prominent feature is the bradycardia, and this, in conjunction with the symptoms, suggested to my mind that we might possibly have here a case of what is known as heart-block, or perhaps better called Stokes-Adams or Adams-Stokes disease.

By heart-block is meant a state in which, in place of the usual rhythm the auricles and the ventricles contract more or less independently, the disease varying from incomplete heart-block, in which the normal relation is disturbed, to complete heart-block, in which the auricles and the ventricles contract quite independently of each other. The nature of this condition has been clearly established by experiments on the lower vertebrates, of which those of Stannius are the earliest. These experiments consisted in ligaturing, in the frog, the connexion between the sinus and the auricle, known as the first Stannius ligature, and the connexion between the auricle and the ventricle, known as the second Stannius ligature. The result of the first ligature is that the sinus continues to beat, but the auricle and the ventricle, after a pause, begin to beat with a rhythm independent and slower than that of the sinus. The result of the second ligature is that the sinus and the auricle continue to beat as before, but the ventricle, after a pause, begins to beat with its own inherent rhythm, which is slower and independent of that at which the sinus and the auricle are beating.

Recent experiments are those by Erlanger on the mammalian heart. Erlanger practically carried out the second Stannius ligature. He ligatured, or rather compressed, in a dog the fibres which are known to connect the auricles with the ventricles in the mammalian heart, forming what is now known as the auriculo-ventricular bundle.

The experimental work of Gaskell has demonstrated that the conduction of stimuli through the heart from the auricle to the ventricle depends upon muscular continuity and not upon any nervous structures, and that the auriculo-ventricular bundle is a bundle of specialised muscle fibres that carries on this conduction of stimuli from the auricle to the ventricle. In a paper by Professor Aschoff of Freiburg² will be found a very full and detailed account of the topography and histology of this bundle. Aschoff and his pupil, Tawara, have worked this out in various animals (dog, cat, sheep, calf, rabbit, rat, guinea-pig, pigeon) as well as in man.

I shall content myself with reminding you that the bundle is not a fiction of the imagination but, briefly, has the form of an inverted "Y," which commences in the auriculo-ventricular node situated in the wall of the inter-auricular septum near the opening of the coronary sinus into the right auricle, and passes down in the inter-auricular septum, through the fibrous auriculo-ventricular ring into the inter-ventricular septum where it divides at the lowest part of the pars membranacea septi into two branches, one passing into the wall of the right, and the other into the wall of the left ventricle. Finally, the fibres of each branch spread out under the entire endocardial lining of the ventricular cavities—their final ramifications being what have long been known as Purkinje's fibres. It may be mentioned that the muscular bundle above described is supplied by a special branch of the right coronary artery.

Now Erlanger³ by compressing this bundle of fibres in the dog's heart carried out in the mammal what the second Stannius ligature did in the frog, and this is the result he obtained. With very slight compression the interval between the contraction of the auricle and the ventricle was increased. With further compression a ventricular beat was occasionally missed and this occurred at regular intervals—for example, after every tenth auricular beat. With still further compression a beat was missed after every ninth, eighth, seventh, sixth, and so on until every alternate auricular beat failed to find any response on the part of the ventricle. The condition now produced may be diagrammatically represented thus, and is known as a 2:1 rhythm. (See Fig. 1. The

¹ A paper read before the Midland Medical Society on Dec. 1st, 1909.

² Brit. Med. Jour., Oct. 27th, 1906.

³ Ibid., Oct. 27th, 1906, p. 1111.