effective and less costly than other and more generally accepted methods for the remedy of certain disabilities, when intelligently applied. It is in the hope that serious consideration will be given by the State to the adoption of these means not merely for sailors and soldiers disabled in this war but for those engaged in other public or national services of the future that I write this note. At a very large command dépôt, of which I was appointed to the medical charge, I found, on taking over, a great amount of indoor treatment being done with not very satisfactory results, but on sweeping out much apparatus and substituting outdoor physical training a striking improvement followed. The system of open-air physical training was therefore developed to its fullest extent, supplemented by hydro-therapy and massage for a small percentage of cases. Dr. Frank Radcliffe, who contributed a letter to THE LANCET on the Value of Physical Treatment,¹ ably assisted me in this work and had charge of those men requiring special or individual attention. The vast majority of the men were dealt with in larger squads or classes under special instructors who had been trained at the Army Physical Training Schools at Winchester or Aldershot. Graduated exercise has done an immense deal for tuberculosis (the question of auto-inoculation apart), and I am confident that, properly supervised, outdoor physical exercise will refit a large number of men who, treated under other conditions, would drift into a state of pauperism.

I am, Sir, yours faithfully,

J. H. P. GRAHAM,

Manchester, Dec. 23rd, 1918. Lieutenant-Colonel, R.A.M.C. (S.R.).

THE MEDICAL EXAMINATION OF AVIATION CANDIDATES.

To the Editor of THE LANCET.

SIR,—In your issue of Dec. 14th Dr. G. A. Sutherland rightly says that a proper selection of men fit to fly cannot be made unless the examiner is also fit to make a proper selection. He then goes on to show how the examiner should be trained in this work, laying down principles which should act as a guide. With these excellent principles all will agree; the practical application of them, however, on the part of the examiner is by no means easy. It takes time to make an experienced assessor, able to weigh one set of functions against another. The beginner is apt to be over-cautious in estimating the character of the pulse and in imagining cardio-vascular debility. In certain men with large muscles it is difficult to feel the radial pulse when the arm is raised, and it is necessary to press deeply with the fingers. When raising the arm to estimate the character of the pulse the elbow must be bent to prevent the muscles becoming tense. These are some of the practical points liable to be overlooked by beginners.

Standards of aerial fitness must be adapted to the varying conditions. At the beginning of the war it was thought wonderful for an aeroplane to travel 60 miles an hour and to rise to a height of 10,000 feet in half an hour. Now it is usual for airmen to travel at three times that speed and to ascend in one-sixth the time, and the end is not yet. Now, when we subject the human body to such sudden changes in oxygen pressure and temperature it is reasonable to devise some test to decide whether a rapid ascent to a high altitude can be tolerated. This is a question where physiology can be usefully combined with clinical medicine. If a simple test can be devised it should certainly be used. Amongst the borderland cases we want evidence which will throw light on the functional efficiency of the cardio-vascular and nervous systems. Dr. Sutherland suggests that to stand the candidate on one leg with his eyes shut for 15 seconds, to observe whether his outstretched fingers are tremulous and whether his knee-jerks are normal is sufficient evidence for the examiner to go upon. These are all important points, no doubt, but why not test also the cardio-respiratory centres in the medulla? This can be done thus:—

The candidate is directed to expire so as to empty the chest and then to take a deep inspiration and blow through a tube so as to raise a column of mercury to the height of 40 mm. and keep it there for 40 or 50 seconds. The experiment may produce considerable embarrassment, with marked flushing of the face, and when asked to describe his sensations the candidate will often say that he felt a "bursting in the head" or that "he felt giddy." This may be termed an abnormal answer.

¹ THE LANCET, 1918, ii., 860.

On the other hand, a youth who has good nervous control and a good cardio-vascular system will hold his breath against 40 mm. Hg pressure, with little or no discomfort for the same period, and his answer will be that he "wanted to breathe" or he "wanted more air"—a normal answer. During the holding of the breath the pulse-rate in the two cases varies: in the normal boy the rate will increase from 20-30 beats per minute and stay there until the end of the experiment, whilst in the abnormal case the rate will increase by 60 or more beats per minute, and before the end of the experiment it will drop to a rate lower than at the commencement. I have watched the manometer test in a great variety of cases, and I feel sure that it has its value and should form part of our examination, because it affords evidence of the stability of nerve centres which cannot be elicited in any other way; but, of course, it should never take the place of a thorough physical examination.

The Medical Research Committee have published some statistics on this subject which show that successful pilots who are fit can pass this test, whilst those who are "stale" or who have suffered from crashes cannot. More work is, no doubt, required before the test can be generally accepted, but at this early stage in the problems of flight we must try to keep our minds open with regard to the tests for flying men.

A number of chamber experiments have been carried out on men who have suffered from faintness or other flying disabilities at certain heights. The conditions prevailing at these varying heights, such as pressure and lack of oxygen, have been reproduced, and it is remarkable how the same symptoms appear again at approximately the same heights. These results seem to me to be too suggestive to be passed over in silence.—I am, Sir, yours faithfully,

H. RONALD CARTER,

Temp. Capt. R.A.F.; late Member, Aviation Candidates Medical Board.
Dec. 17th, 1918.

ADVANCES IN THE TREATMENT OF FRACTURES.

To the Editor of THE LANCET.

SIR,—In their letters in THE LANCET of Dec. 14th and 21st Dr. R. Scot Skirving and Sir W. Arbuthnot Lane have brought to notice again the great importance of Major M. Sinclair's work on the treatment of fractures. The military surgeon has found the methods he has advocated of the greatest value, and a large sum of money in pensions has been saved by their adoption. No other method of splinting can compare with Major Sinclair's for immobilising the parts, extending fractures, and at the same time allowing of easy access for dressing the wound. On his net frame cases that would be otherwise a perfect nightmare to sisters become quite easy to dress and nurse, and the patient ceases to live in constant dread of the next dressing, or drawing of his sheet. By fitting the patient's boot on to a Thomas's splint a cheap and easily obtained ambulatory splint is produced which allows the limb to be used and greatly improves its nutrition. One can scarcely enter any ward without seeing some patients being treated by one or other of Major Sinclair's original methods of extension.

We are, Sir, yours faithfully,

JAMES TAYLOR, Major, R.A.M.C.,

District Consulting Surgeon, Aldershot Command;

W. S. EDMOND, Major, R.A.M.C.,

Dec. 23rd, 1918.

Surgeon i/c Division, Cambridge Hospital.

THE CAUSES AND INCIDENCE OF DENTAL CARIES.

To the Editor of THE LANCET.

SIR,—For upwards of 20 years I have on several occasions insisted that the appalling prevalence of dental caries and "adenoids" in this country is due to remedial dietetic causes. It was therefore with great interest that I read Major W. E. Nickolls Dunn's letter in your issue of Dec. 21st, calling attention to this matter. I have again and again referred to the prosaic fact that there are among the inhabitants of this country some 200 million carious teeth, as many alveolar abscesses (pyorrhoea alveolaris), and some 30 million root abscesses—for the most part preventable by the simplest possible means. Well may Major Dunn exclaim that it "is impossible to exaggerate the state of the teeth in this country." It constitutes, indeed, a national disgrace. Dr. Sim Wallace has time after time debated this matter with me, and we have long felt that the remedy can only come through the medical profession.

Three great problems confront the Ministry of Health—the problems of food, housing, and venereal diseases—all, I believe, capable of solution by the application of common