by hypothesis molecules are incapable of exerting *direct* force on one another, not being in contact). The medium is exerting the force, and will continue to exert it over a certain range, hence it possesses the potential energy; when released it will do work and transfer its energy to the steel, in the kinetic form. If the spring overshoots its mean position, the energy is re-transferred and transformed. A perfectly elastic bounding ball has all its energy transformed into potential, at the middle of every period of contact with the obstacle from which it rebounds.

One case possesses perhaps a little more difficulty, viz. the case of a bullet fired into dough—when the body exhibits no recoil: how can the energy of the bullet be said to be transformed into potential now? Only by remembering that heat-motion is a vibration of some kind, and that when a vibration is excited by a blow or by friction, *strain* is the effect first produced, and afterwards the recoil. Think, for instance, of exciting a tuning-fork or a string by bowing or striking it.

Alternations from kinetic to potential may be rapid in some cases, slow in others: no matter; all I have stated is that change of form is necessary and universal whenever energy is transferred, *i. e.* whenever any kind of activity is exhibited by any known kind of material existence.

University College, Liverpool,
May 16, 1885.

LIV. *On the Paths of Electric Energy in Voltaic Circuits*
Appendix to Paper on the Seat of the Electromotive Forces in
the Voltaic Cell. By Prof. OLIVER LODGE*.

[Plates IV. & V.]

THE main conclusions to which I have been led with regard to the potentials of metals and the seat of E.M.F. may be very briefly stated thus:—

(1) A metal is not in general at the same potential as the air in contact with it; the difference of potential (or contact-force) between any given clean metal and air being calculable, at least approximately, from thermo-chemical data, though there is no known way of experimentally observing it.

(2) Putting two metals into contact equalizes their poten-

* Communicated by the Author. For the sake of completeness it may be convenient here to mention that a report of a discussion on Seat of E.M.F., at Montreal (Brit. Assoc., Sect. A), is published in the 'Electrical Review' for Nov. 22, 1884, and that a more elaborate discussion on the same subject, by the Society of Telegraph Engineers and Electricians, will be reported in their volume of 'Proceedings' next issued after the present date.

tial, their surfaces becoming charged with the electrostatic charges proper for effecting this equalization of potential.

(3) The air surrounding a pair of different metals in contact is therefore in a calculable condition of dielectric strain, especially near the junction of the two metals; in other words, there is in the air a slope of potential which is most rapid in the neighbourhood of the junction, and is there easily observed electroscopically.

(4) This slope of potential observed by Volta, measured by Kohlrausch &c. &c., near the junction of any two metals, has caused observers to imagine that the metals themselves differed in potential, and accordingly to postulate an E.M.F. or contact-force resident at the junction in order to maintain such difference of potential in a conductor.

(5) No such force exists at a metallic junction; because, if it did, the passage of a current across such a junction would produce violent reversible thermal effects, and none such are observed.

(6) This non-existent force has been imagined, and has been utilized to account for the greater part of the E.M.F. in a voltaic circuit; in fact, it has itself been considered to be the main E.M.F. in such a circuit; although it was manifestly unable to account for the energy of the current.

(7) The real seat of the E.M.F. in a voltaic or any other circuit must be where conversion of energy, from some other form (chemical, thermal, &c., &c.) into the form known as an electric current, occurs.

The object of the present Appendix or Supplement to my paper is to supply an accidental omission in the history of the subject, and more particularly to exhibit and illustrate the mode of regarding conducting circuits which I advocate, by diagrams of energy-paths, or *erg-odes* for those who like Greek barbarities.

The accidental omission from the history of the subject is a notice of Dr. Gore's communication to the Royal Society of November 1883*, in which he makes an attempt to bring liquids into a thermoelectric series.

The contact-force between a metal and a liquid is shown to vary with temperature; and the results indicate that this contact-force is a complex one, not simply depending on the chemical relations of the substances in circuit, *i. e.* on the cor-

* Gore, Proc. Roy. Soc. No. 233, vol. xxxvii. p. 251: "Some relations of Heat to Voltaic and Thermoelectric Action of Metals in Electrolytes." See also former papers in Phil. Mag. January 1857; Proc. Roy. Soc. 1878, p. 513, No. 188; 1879, No. 199, and 1880, No. 208.

rodibility of the metals, but involving some other more purely physical force also :—much as Bouty surmised from his experiments on the generation of heat at a metal-liquid junction when a current was driven across it.

A number of tables of the order of metals in different solutions of different strengths and at different temperatures are given, together with some numerical values.

It is possible that by an attentive study of Dr. Gore's work, and of other work in the same direction, a more complete account could be given of the exact relation existing between what may be called the thermal and the chemical contact-forces at metal-liquid junctions ; if indeed any valid distinction can actually be drawn between them.

At present, at any rate, I do not attempt this; but proceed to the other matter spoken of, viz. the illustration of my views on the voltaic circuit by a diagram of its energy-paths.

Prof. Poynting has taught us, in a paper of the very greatest interest and power*, that modern views of *vis a tergo*, as opposed to action at a distance, lead necessarily to the idea of continuity in the existence of electric, as well as of all other, energy, and hence to a study of the paths along which it moves from one part of the field to another. He has shown that these paths are the intersections of the magnetic and the electrostatic equipotential surfaces, and that the rate of flow of energy is proportional to the product of the electrostatic and electromagnetic intensities at every point.

This view of the electromagnetic field, though no doubt implicitly involved in Maxwell's equations, yet urgently needed to be dragged forth and exhibited ; and it is difficult to imagine how the developing process could have been done better than it is done in Prof. Poynting's memoir.

The special cases by which his memoir is illustrated are not, however, in exact accordance with the views of the electric circuit which I am advocating ; and in particular the diagram of the curves for a voltaic current is, in my view, wrong. I cannot say I thought it wrong at the first glance ; I was pleased to see that Mr. Poynting held what I consider the correct view, that the main seat of the E.M.F. in a simple pile is at the zinc-acid junction and not at the zinc-copper, and that his diagram embodied this view.

But when I came to draw the curves myself for this case, taking account of the step of potential which I believe to exist between a metal and the air, I found the energy-paths refuse to start spreading out from the zinc-acid junction, as we had

* Poynting, Phil. Trans. 1884, pt. ii., "On the Transfer of Energy in the Electromagnetic Field."

both at first sight expected them to : they insisted on diverging from the zinc-copper junction, as a Voltaist would wish them to. This behaviour is really in perfect accord with my theory; though the fact is the cause of Voltaists' views concerning seat of E.M.F. They locate the force at the zinc-copper junction just *because* the energy-paths diverge thence and appear to start thence.

Prof. Poynting's diagram, therefore, not only differs from my theory, a thing which is natural and unimportant, but it is discrepant with experimental fact.

Rigid experimental truth would have led him to spread out his energy-paths from the metallic junction, not from the acid junction as he has done. True such a proceeding would have landed him in difficulties and apparent absurdity, and he would have been compelled to ask, "How can all the energy start from an inert metallic junction?" And he might well have considered such a conclusion absurd, as it is.

The reconciliation between sense and fact is simple. The lines do, indeed, diverge and spread out from this junction, but they do not *start* thence. They all start from the zinc-acid junction; but they at first creep along close to the surface of the zinc, investing it like the coats of an onion; and some continue to creep along the copper too, but some do not. More lines invest the zinc than the copper; and the lines in excess are given off at the junction, and thence diverge through the air.

An electroscopic experimenter seeking the seat of E.M.F. traces these lines, home as he thinks, to the zinc-copper junction, and hence regards this as their origin and the seat of the force. It is not. The lines do not start thence; they start at the real seat of E.M.F. and source of all the energy, though their close-lying continuations between metallic junction and acid junction are impossible to trace out experimentally.

With this preliminary, the diagram of energy-paths for a voltaic circuit in Plate IV. almost explains itself.

The lines are loci of constant electrostatic potential; and I happen to have chosen a circuit whose external resistance is twice its internal. The junction of the two metals is placed at random wherever its peculiarities may be most conveniently observed. The liquid in which the metals are immersed is only faintly indicated by the outline of its trough, and, for simplicity, it is supposed to be one differing in no electrostatic respect from air. Dilute sulphuric acid of a certain strength has been shown by Prof. Clifton to satisfy this condition almost exactly.

The particular metals for which the diagram is intended to

be drawn are zinc and copper. Now thermo-chemical data lead me to conclude that the step of potential between clean zinc and air is 1·8 volt [the exact decimal not being insisted on except for purposes of illustration] ; hence, if an equipotential line be drawn for every tenth part of a volt, 18 of these lines would closely invest clean zinc always, and form a boundary between it and the air. In the diagram (Pl. IV.) lines are only drawn for every fifth of a volt, and consequently 9 of them envelop the zinc.

As for copper, it is probable that the step of potential between it and air is ·8 of a volt ; and accordingly four lines invest the copper at every point in the diagram. They are not always *the same* four lines—that is of no consequence ; the essential thing to remember in drawing the diagram is to keep the number constant at every point for the same metal. Such a metal as gold or platinum may be nearly free from lines.

It is needless to say that the lines have to be drawn with spaces between them for distinctness, but that in reality the step of potential is of the most sudden description, and all investing lines ought to lie close to the mathematical surface of the metal.

Now, since nine lines coat the zinc, and only four the copper, it is manifest that five lines must leave the metallic surface at the junction, and must spread out into the air. These five lines correspond to the one volt* Volta effect observed near a junction of clean zinc and copper, and which has been called a difference of potential between zinc and copper.

Of the five lines thus starting to spread out from the junction, some make for points on the external circuit, and others go through the internal circuit. Those which go to the external circuit can by no means go through it, or into it, for the potential of the copper differs by ·8 volt from that of the air ; but they can go round it, and leave it on the other side just as if they had gone through it. Moreover, a line does go through, or into, the copper at the very point where the air-line abuts against it, viz. the line of potential ·8 volt lower. So the transfer of energy into the conductor of the external

* Different observers give different values for this number ; from ·5, Kohlrausch, to between ·76 and ·89, Ayrton and Perry and Clifton, up to ·92, Pellat ; but since tarnish or dirt on the zinc surface will certainly lower it, and it is difficult to suppose any zinc surface absolutely clean, I have not thought it necessary to modify the theoretical value of one volt, which at any rate serves well enough for illustration, though it may turn out to be about ten per cent. too high.

circuit is complete, except for just the metallic skin, and everything goes on *in the outer field* as if the energy-paths really started from both sides of the zinc-copper junction, and some of them cut through the metal of the external circuit with degradation into heat.

Note, however, that the heat-energy in a wire is not really thus carried through the air : it creeps along the surface of the wire, some of it perpetually diving into the wire, and becoming converted into heat.

The energy which liberates hydrogen from the copper-acid surface, moreover, does not reach it in the way Mr. Poynting has supposed. Some of it crawls over the current the whole distance ; all of it goes some distance with the current.

In the electrolyte, however, this is not so ; and the heat which there makes its appearance is produced at the expense of energy which has arrived *via* the air from the zinc-copper junction. This is on the supposition that no contact-force exists between the electrolyte surrounding the plates and the dielectric medium (air) enveloping the rest of the circuit. In cases where such a contact-force does exist, some energy will creep along the air-liquid surface, just as it does over the surface of a metal.

In the diagram one of the five lines here spoken of is doubled back upon itself, so as never to leave the surface of the zinc : the spaces between the lines are the real paths of the energy of course, not the lines themselves ; the lines only indicate the directions across which no energy goes. There are five spreading-out energy-paths, then, and it is most instructive to watch the energy arising at the zinc-acid surface flow in its first narrow channels, then broaden out through space, and then cramp itself again into its tube, by which it reaches the copper surface, and is transformed into energy of chemical separation ; or, again, to watch that in the channels touching the copper fritter itself gradually away into heat.

Energy-paths in Secondary Circuits. (Plate V.)

The four figures in Plate V. are intended to indicate the transfer of energy to a secondary circuit, *i. e.* one in which an induced current is being generated.

Fig. 1 shows a steady electromagnetic field disturbed by the presence of a stationary copper channel. The energy simply flows round it, as Prof. Poynting has said. Fig. 2 shows a field no longer steady, but with the potential changing : the energy-paths moving, say, toward the right. Each line, as it comes up,

folds itself round the circuit, and sends part of one of the previously investing paths to the inside, where it displaces part of one of the original inside paths, which at once begins to close up. Fig. 2 shows it half closed, on the hypothesis of absolute suddenness. Fig. 3 the same thing, on the supposition that energy possesses something like inertia. The induced current is produced by energy flowing along the semi-displaced path through the copper. If the change of field stops, fig. 2 is supposed quickly to revert to fig. 1, with an opposite transference of energy. Three stages in the change of field are shown in fig. 4 : or this figure may be regarded as showing three lines in a uniform field disturbed by a moving conductor.

The arrows are drawn as if the energy flowed right away again without loss, or at least without complete loss. It is easy to reverse one of the arrows, so as to make energy flow up from both sides to the secondary circuit; but it is not easy to see how energy can reach the copper and be dissipated when its path does not pierce the copper at all. The lines have to be thought of as first embracing the circuit, then as moving away, having left a portion enclosing the circuit behind, and finally, more lines having come on, as sinking into and through the copper, and shutting themselves up; and so on perpetually, as long as the field is changing. The energy thus reaches the secondary circuit a little behindhand always, which is not wrong. If all the energy is dissipated in the act of passing through the copper, arrows ought not to be drawn on the inside paths which are shutting up; because though they are still paths for energy, it does not follow that they are made use of; in fact, they cannot be, if all the energy is extinct.

Probably, however, what all these diagrams better represent is the steady state of a field disturbed by the presence of a *perfect* conductor in which there is no dissipation whatever—a conductor like the molecules postulated in Ampère's and Weber's theory of magnetism and diamagnetism.

Voltaic Energy-paths. (Plate IV.)

Returning to the Plate showing voltaic-energy path: the diagram seems to me to constitute the most absolute and complete reconciliation between the truth in the opposing views of the voltaic circuit.

It not only justifies modern voltaic experimenters in their recognition of the slope of air-potential near a junction, but it partially justifies even the view of old Volta that this

Phil. Mag. S. 5. Vol. 19. No. 121. June 1885. 2 L

junction is the source of all the energy. So far as the external circuit is concerned, the metallic junction *is* the source of the energy ; all the energy does in a manner start thence, though it does not start *originally* thence. It reaches the junction from another place, its real source, but it travels thither by infinitely unrecognizable paths.

Does the diagram, then, justify modern Voltaists altogether? Certainly not. It cuts the ground from under the feet of those who have asserted that there is an E.M.F. at the junction of zinc and copper—a contact-force which is the cause of the observed difference of air-potential in the neighbourhood. And it annihilates those who have seen in this imaginary and non-existent force any means of propelling the current in a voltaic circuit.

On the other hand, the diagram of course justifies all those experimenters who have regarded the zinc-acid junction as the main seat of energy-transformation, and therefore of electromotive force. At the same time it exhibits the crudity of some of their statements ; and, just in so far as it upholds the truth and consistency of the Voltaists with regard to a part of the phenomenon, it contradicts some of the opposing statements made by chemical theorists with regard to the Volta effect. In particular, it lends no support to the customary “chemical” view, which sets down the whole Volta effect as an accident due to unavoidable chemical action, and so practically ignores it.

Imagine the lines investing the metals to be microscopically close to them, and then look at the diagram as truly exhibiting the facts of nature : it irresistibly suggests a voltaic or contact theory of the voltaic pile. It was right, therefore, for the first observers of voltaic phenomena to found a contact theory, and to adhere to it tenaciously, as in some sense true. It was also right for later philosophers, imbued with notions of energy, to feel the doctrine that the metallic junction was the source of power to be absurd and incomprehensible, and to found a chemical theory.

In the conflict both sides have made mistakes ; but it is the essence of truth in both views which has given them such vitality and endowed their upholders with such energy of persistence.

University College, Liverpool,
May 17, 1885.