SPINAL TRANSPLANT *

HENRY BASCOM THOMAS, B.S., M.D.

Attending Orthopedic Surgeon, St. Lukes Hospital and Cook
County Hospital
CHICAGO

This article was suggested by Dr. E. Wyllys Andrews, and is a plea for fair judgment, favorable or unfavorable, of the autobone transplant of Dr. Albee for tuberculous spondylitis.

The treatment of tuberculous vertebrae by the non-operative method in the hands of most orthopedists terminates satisfactorily to the patient and to the sur-

Fig. 1.—Patient bending and touching the fingers to the floor. Notice the stiffness and absence of gradual even curve in this back. Compare with Figure 2, a normal back.

geon in those cases which he can keep under control for a long period of time — say two or three years. There are cases, however, in which future health and bread-earning ability depend on:

1. Such control as the operative method seems in some instances to give.

2. A shortened period of recumbency which in a number of my cases has apparently been due to the added support of the bone splint. It is only fair to give the facts regarding the results, good and bad, of the work in this field, to those who may help to judge of the merits and disadvantages of such treatment, and it is hoped that those who have operated in one or two cases with unfavorable results will not consider worthless this new, under-the-skin splint before they have the data of many cases, based on observations extending over some years.

Many adults frequent the hospital at ages ranging from 30 to 45 years whose history is much like the following cases and whose condition is due to what I call lack of control sufficient to establish and continue treatment to firm ankylosis, which means a working physical condition.

CASE 1.—A woman aged 39, who appeared much older, gave a history of painful and deformed back since girlhood, and related many periods of illness which confined her to bed and prevented school attendance and work. She always insisted that she was well as soon as the recumbent position, cast or brace had eased the back pain, and it was impossible to continue treatment at any one time more than two months. Extreme emaciation and multiple sinuses are the results. We should ask ourselves if operative measures would have established control in this patient in her youth and prevented, at least in an indirect way, the invalidism she has carried for life.

A second case will further illustrate what I mean by control.

CASE 2.—A girl aged 14, who had been under treatment in recumbence, cast and brace, at various times for two years, each time left the hospital, discarding cast and brace in the face of every effort to the contrary, and returned each time only after she was unable to work longer. During one stay in the hospital she developed a condition diagnosed as tuberculous meningitis, but recovered, and I therefore doubt the diagnosis. At her last appearance at the hospital, she was prepared for operation the day she entered, the operator reasoning that her only hope to escape the fate of Patient 1 was in this procedure. Otherwise, the comfort of a Bradford frame would have influenced her to walk out of the hospital prematurely. Chest examination revealed pulmonary tuberculosis. The Roentgen ray showed caries of the bodies of the sixth and seventh dorsal vertebrae, and there was slight

Fig. 2.—Normal back showing the gradual, even curve, unlike that in the grafted spondylitis, as in Figure 1.

angular kyphosis and very painful, rigid back. She was operated on, kept in bed three months, and allowed to be up and about without support at the end of four months.

Technic of Operation.—An iodine preparation was followed by an 8-inch elliptic incision including the fourth, fifth, sixth, seventh and eighth dorsal spinous processes, which were split and fractured at the base with knife and chisel, about one-third their thickness being turned down. The soft interspinous tissues were hollowed out, so as to form a groove $5\frac{1}{4}$ inches long, $\frac{3}{4}$ inch deep and $\frac{1}{2}$ inch wide for implantation of a section of the tibia. Bleeding was controlled by means of hot sponges.

* Read before the Chicago Medical Society, Jan. 30, 1914.

An elliptic incision was made over the left tibia about 5 inches in length, the anteromedial surface of the tibia bared, and the tibialis anterior muscle pushed outward $\frac{1}{8}$ inch. Incisions were made through the periosteum lengthwise for $5\frac{1}{4}$ inches on each side of the anterior border of the tibia, the periosteum included, and a wedge-shaped section of the bone $5\frac{1}{4}$ inches long by $\frac{1}{2}$ inch deep was excised. This transplant was then cut transversely in three or four places to allow it to be bent easily, and a hole was drilled in both ends.

Strong catgut on an aneurysm needle was introduced through one side of the split fourth spinous process, then

Fig. 3.—Case showing knuckle in high dorsal region, but on account of the prominent knuckle best adapted for Albee's method of operation.

through the drilled hole in one end of the transplant, and thence through the other section of the spinous process, thereby securely anchoring the upper end of the transplant between the split spinous processes. The other end of the transplant was secured in the same manner, and made to fit into the groove between the split spines. This groove was made slightly to the left of the median line. The transplant was then covered over by the paraspinal tissues, the suturing material being heavy chromic catgut. Tension sutures of silkworm gut were used to close the wound and the skin approximated with Michel clips. Dry dressings were applied with a circular pad of felt, preventing pressure on the incision. The leg incision was closed by means of fine catgut for the periosteum and black silk for the skin. As soon as the patient was removed from the operating-table, and before she recovered from the effects of the ether anesthesia, she was placed on a Bradford frame.

VARIATIONS IN TECHNIC

The elliptic incision over the tibia should not be used for the back operation. It has obvious advantages, but, over the tibia, it has no place. Here, the best technic, for comfort, shortness of incision, least disturbance of muscle and perfect healing, is to incise directly over the anterior border of the tibia, dissect the muscle back slightly on the outer border, protecting the periosteum, mark out the size of the bone by cutting down on it through the periosteum, according to the measure obtained from the back; then this bone should be either

sawed or chiseled out, leaving the periosteum intact. It will be found that, in the young patients, one accustomed to chiseling the bone out finds this a quicker and more satisfactory way than sawing. The ends and that part of the graft—usually the middle—which is embraced by the split halves of the most prominent process in the knuckles, should be very deeply and firmly fixed to secure the best results; otherwise, the ends spring up so that in some of the cases the finger could work the skin of the back under the ends of the graft. This aids absorption to take place before bone formation begins. The best fixation I have succeeded in obtaining has been by kangaroo tendon passed through the holes bored in the ends and middle of the graft. Pegging with autobone or ivory pegs has not been so satisfactory. Silk may, in the end, cause increased ankylosis by the irritation following slight infection and expulsion, and the kangaroo has given the best results in this series of cases. Regarding the postoperative position, the Bradford frame has been used with satisfaction in all but two cases. Patients, however, are comfortable on the chest-abdomen, but do not get the same tendency toward hyperextension, reduction of the kyphosis and relief of the vertebral pressure. The addition in the back position of a well-fitted felt pad around the kyphosis adds the advantages of the chest-abdomen position.

CASE 2.—Later Report of Condition.—Following the operation, the patient's condition, probably because of the lung tuberculosis, was very uncomfortable; the pulse was fast and weak at first; later, better, and after two days the prospects were gratifying. From now on her condition could be compared favorably with that of most patients following a major operation. The absence of pain, considering the double operation and the length of incision, was noticeable. Each wound healed perfectly; the slight kyphosis is now absent, and motion is painless and increased above and below the graft, so that, standing with stiff knees, she puts the fingers of both hands on the floor (Fig. 1).

Again we must ask ourselves, is not this patient, at five months after operation carrying a bone splint which she cannot remove, and having increased fixation, a better risk than she would be with only the recumbency treatment for the same length of time?

SHORTENED PERIOD OF TREATMENT

Albee, to whom great credit is due in this work, disregards mechanical support after the operation, and gets his patients up early. This may be a safe procedure when a heavy bone splint has been deeply and securely placed between the split spinous processes, and the bodies of the vertebrae thus held apart without the aid of callus between and around them; but the trunk is so heavy, the splint so small in comparison with the weight it

Fig. 4.—Case with prominent knuckle in lower dorsal and lumbar region, best suited for the Albee method of operation.

must bear, and its leverage at such a disadvantage, that in this series of cases the splint has been looked on only as a temporary support, aiding the recumbent, brace-cast treatment, while heavy callus fills in about the vertebrae, supplying the finishing touch to ankylosis. Therefore, the patients whom I mention and whose cases I report have been given from three to ten months' support after the operation, on the supposition that if the treatment is continued for even as long as eighteen months, the time of the average non-operative treatment has been cut down about one-half.

Fig. 5.—Broken transplant (Case 6) tied with kangaroo. The kangaroo should go through holes made in each side of the split spinous process of the most prominent diseased vertebra.

SELECTION OF CASES

In this series of cases, no attempt was made to choose the good risks, operation being performed as each case was presented for treatment.

Factors which should be considered in selecting cases for operation are:

1. The general vitality of the patient.
2. The age.
3. The position of the disease in the vertebral column.
4. The amount of bone destruction.
5. The amount of callus formation.
6. The control of the patient.
7. The eagerness of patient and parent for shortened duration of treatment.

1. The more vigorous in constitution, the better the risk, but it has been surprising how nicely, with the exception of one, each patient, among those whose condition was considered hopeless, stood the operation and convalesced to an improvement in the general condition.

2. The age of the patient is considered an important factor by some who fear that the graft *per se* may not grow, and may thus tend to deform the backs of those young patients whose spinal columns will continue to grow and, therefore, bow backward when held by an unyielding, non-growing graft. Believers in this theory would select only the adult spine for operation. Experiments are in progress to determine the growth or non-growth in length, of the graft, and reports are in preparation for June, 1914.

3. The position of the disease in the vertebral column is a contra-indication to operation: (a) in those cases involving the high dorsal and low cervical region if the knuckle is so sharp and high that the occiput touches it and prevents the necessary space for work, and (b) in those cases in which the disease is in the middle and upper cervical regions.

Selection of that operation which will best lend itself to the particular region diseased.—I am inclined to

select the lower dorsal, lumbar and sacral regions as the most favorable field for the Albee operation, and the middle and upper dorsal and lower cervical regions for the Hibbs operation, on account of the advantage given by the length, thickness and angle of projection of the spinous process in the different regions. Probably the most favorable position as offered by the spinous processes for the Albee operation, as far as ease of preparing the bed and firmly placing the transplant are concerned, is that of the lumbar and twelfth; sometimes, also, the eleventh dorsal vertebra. In the Hibbs operation, when the base of a spinous process is fractured across, the end denuded and turned down into the space caused by the fractured base of the neighbor below, best results are obtained when the sixth and seventh cervical, and all the dorsal spinous processes, with the exception of the eleventh and twelfth, are selected. This is because their length and relative position to their neighbors makes them yield easily to the desired technic. When the kyphos is sharp and large, separating the spinous processes, the condition becomes more favorable for the Albee operation, even though the kyphos is in the dorsal region (Figs. 3 and 4).

4. When the amount of bone destruction is great, such help in ankylosis as comes from the graft is indicated more than in those cases in which the small bone destruction can more easily be replaced from the bodies of the vertebrae.

5. The amount of callus formation and its ankylosing effect is the greatest element in the cure of these cases, and when it is large and strong enough to hold the back firmly, there is no need for placing a bone graft.

6. The control of the patient, as already indicated in this article, should be an indication for operative treatment in those cases in which the patients themselves or through their parents persist in discontinuing treatment early, much before the callus from the bodies of the vertebrae has had time to back up the support of the bone graft.

7. Other things being favorable, the desire of the patient or the parents to cut down the mechanical treatment a year or even a year and a half should be an indication for the operation and the aid it gives in accomplishing this wish.

Fig. 6.—This patient is resting on a frame in hyperextension. The scar on the leg can be plainly seen. Michel clips were used but remained in place too long.

Among those cases, of this series of twenty-seven, which were so bad that the patients might have been denied operation were the following:

CASE 3.—A boy, aged 4, with an active tuberculous hip complicating a tuberculous spondylitis which included the fifth, sixth and seventh dorsal vertebrae, and a discharging sinus $2\frac{1}{2}$ inches to the left and on a line with the tenth dorsal vertebra, progressed splendidly. Following this operation he had only slight after-pains, no complications, clean wounds not damaged by the nearby sinus, and marked improved general condition. His kyphosis decreased in promi-

nence because of the hyperextension on the frame, but no more than would have occurred without the operation. The patient was dismissed from the hospital early, and was allowed up with no support. Nine months after the operation, the kyphosis is increased; the sinus is closed.

CASE 4.—A boy, aged 16, poorly nourished, with marked spasticity of the legs, an ulcer about as large as the palm of the hand at the middle third of the right leg, eczema of the back, and incontinence of urine, was operated on in the manner described, except that the ends of the graft were anchored with silk instead of catgut. The wounds healed nicely in the presence of the eczema; the bladder condition improved, and the spasticity appeared at first not so marked. Later, pieces of silk came away, and five months after the operation he continues a bed case; spasticity is not improved; the general condition is improved.

CASE 5.—A man, aged 35, came to the hospital with lumbar Pott's disease, and while there, fell and injured his back. Later, he developed a spasticity of both legs, but was operated on in the usual way. The wounds healed by first intention; the pain in the back has continued; the kyphosis has decreased, but has been pushed to the right, producing a scoliosis. A prominence over the left ribs, resembling the bulging which accompanies the rotation of the vertebrae in scoliosis, has developed. Diagnosis of complicating sarcoma, following the fall, has been made in this case. The patient continues confined to bed, six months after the operation, and is worse.

CASE 6.—A boy, aged 14, with a very marked kyphosis and much lateral bending, unable to stand or sit unaided, was operated on in the usual manner, with this difference: the heavy splint was accidentally sawed through at its middle point, while weakening it with saw cuts so as to bend it to fit the knuckle. The graft was long enough to allow each end to be shaped for mortising, then a hole was drilled through the split sides of the spinous processes and through the mortised ends of the graft, and fixed by kangaroo tendon (Fig. 5). A small pressure sore developed over the most prominent part of the knuckle and graft, but healed in two weeks. Three weeks after the operation a nephritis developed which cleared up nicely on a milk and lime-water diet. The kyphosis is reduced, but hardly more than would be expected from the hyperextension maintained after the operation; there is more support in the back, and the patient can sit up better than he did before the operation. Considering the loss of bone in this case, the operative support should be given its share of credit for the improved condition.

CASE 7.—A girl, aged 9, with a positive neisserian infection, and a marked knuckle of the upper dorsal vertebrae, was operated on in June, 1912. She did badly from the beginning, was unable to stay on the frame, was uncomfortable all the time in any position, and finally, two months after operation, contracted sore throat, followed by pneumonia, and died at her home, after being removed from the hospital by her parents without hospital consent. An examination was refused.

Fig. 7.—A dry transplant holding five dry vertebrae together. The saw cuts allowed some bending of the transplant. This picture gives a good idea of what we try to do in the Albee method of operation.

CASE 8.—A man aged 32, with a marked kyphosis, treated for months in recumbency at a hospital, and discharged when spasticity of the legs developed, was received as a stretcher case and operated on in the usual way. When he was turned to examine the back on the third day after the operation, a bed-sore was discovered over the sacrum. This case was not improved, six weeks after the operation.

CASE 9.—The oldest patient operated on was a man aged 49 who had been unable to work for thirteen months. He experienced remarkably slight pain, and was more comfortable after the operation than before. This patient was kept on a frame for three months, wore a cast for seven weeks, and left the hospital and returned to work, with no support, six months after the operation.

CASE 10.—The youngest patient operated on was a boy aged 2, with involvement of the upper dorsal vertebrae, the graft including the sixth and seventh cervicals. This patient is doing well, and is up and about with a cast one month after the operation. The kyphosis has decreased, and the general condition and the prospects for a firm and early ankylosis are better, in my judgment, than if the operation had not been performed.

The favorable cases, with the exception of Case 2, have not been reported.

TENTATIVE CONCLUSIONS

Experience with this group of cases justifies the following tentative conclusions:

1. The operation is not severe. It is a double operation, and the length of incision is longer than in most operations, yet the recovery from the immediate effect is quick and the suffering is small.
2. Cases with discharging sinuses near the field of operation should not be refused operative treatment on account of the sinus, unless it lies within the field of incision.
3. Complicating pulmonary tuberculosis is not a contra-indication in all cases.
4. If the ends of the transplant are loose, they absorb. If they extend beyond the spinous process, that portion not attached will absorb back to the firm attachment.
5. Loose ends and broken ends, as in the middle of a bent graft, grow, when later, through fixation by position and quiet, they become attached.
6. The three cases complicated with Pott's paraplegia have not markedly improved. The longest time elapsing since the operation of a paraplegia case has been six months. During this time the position has been good, and such as to give hyperextension.
7. The young patients do better than the adults.
8. The operation has a distinct place in the treatment of selected cases of tuberculous spondylitis.
9. Albee's method of operation is preferable in the large majority of cases on account of the location of the disease and the size of the kyphosis.
10. Operative treatment should be followed by months of perfect quiet for the transplant.

Credit is due Drs. Stoland, Packard, Boren, Callentine and Dawson for aid in histories and operations, some of which were performed by them; and to Drs. Hartung, Hubney, Van Horn and Case for painstaking work with roentgenograms.

31 North State Street.

Before Asepsis.—Sir Hector Cameron gave the following brief description of the days when Lister first went to Glasgow: "Every wound discharged pus freely, and putrefactive changes occurred in the discharges of all"; and again: "Secondary hemorrhage, tetanus, erysipelas, septicemia, pyæmia and hospital gangrene were never all absent from the hospital wards, and at times became alarmingly epidemic.—Wrench, G. T.: Lord Lister, His Life and Work.