preparation for Finkler's treatment. Changes in the compound have become necessary, which are occupying us at present.*

But it is to be questioned whether a lecithin mixture or "compound" should have any noticeable advantage over a copper salt of any high fatty acid, such as copper oleate, which has similar physical properties. In these days of immunity reactions lecithin is a *Schlagwort*; we can see no other reason for its preference over oleic acid as a vehicle for copper. There are sound physiologic reasons for expecting particular advantages for the copper salts of amino-acids, but as yet we have not been able to demonstrate any such advantages although we are conducting a systematic investigation of the properties of copper salts of various amino-acids.

We would not be understood as denying the possibility of favorable results being obtained by the use of copper compounds in tuberculosis. Our sole intention is to state that we ourselves have been unable to secure any appreciable effects in experimental animals, and that our experience is not in harmony with the experimental results described by the Countess von Linden, on which the clinical use of copper is based. We have made no clinical observations whatever.

PELLAGRA

WITH SPECIAL REFERENCE TO PATHOLOGY OF GASTRO-INTESTINAL TRACT*

H. P. MILLS, M.D.

Assistant Physician and Pathologist, State Hospital No. 2

ST. JOSEPH, MO.

The changes that occur in the body during the progress of a disease are always of interest and vital importance, and it is worthy of note that from the earliest recognition of pellagra much attention has been given to its pathology. But in spite of painstaking work by numerous investigators, our knowledge of the subject is yet in a somewhat chaotic state. Following the lead of earlier writers, those studying the subject more recently have confined their observations largely to changes in the central nervous system, while the morbid condition of the gastro-intestinal tract, it would seem, has been given insufficient attention. The extent and severity of the lesions vary with the type of the disease present, as do also the symptoms arising from their presence. With our present state of knowledge it would hardly seem possible to decide whether the gastro-intestinal lesions or the changes in the nervous system were the primary or fundamental conditions.

CHANGES IN THE NERVOUS SYSTEM

Sandwith¹ found in certain cases degenerative changes in the spinal cord, most marked in the columns of Goll and Burdach, but in many other cases no pathology was evident. This absence of cord changes was explained by the fact that either these patients had suffered from the disease only a year or less, or they were apparently not far advanced, so that in his opinion cord degeneration, as discovered by the microscope, occurred late in

the disease. From autopsies in cases in which the condition had existed for different lengths of time, cell degenerations were noticed in the brain cortex as well as in the lateral and posterior columns of the cord.

In a large number of autopsies done by Lombroso² and others, variable changes are recorded as present in brain and spinal cord. In some instances the weight of the brain was increased, in others diminished; in some hyperemia was present, in others anemia was noted. Degenerations of a fatty, pigmentary or calcareous type were present in the walls of cerebral capillaries. The pigmentation universally found in the sympathetic and spinal ganglia was considered an indication of marked nutritional changes in the central nervous system. The lesions found in the cord are described as being similar to those due to ergotism, their advance depending on severity of poisoning or lowering of resistance by intercurrent conditions.

Roberts³ states that there is atrophy of the convolutions and in a majority of cases the weight of the entire brain is reduced, but that in many instances the weight is normal or increased. Hyperemia or anemia was noted, and also degenerative changes as mentioned above. Gross cord lesions were not very evident. The microscopic changes noted were degeneration of the direct pyramidal tracts and columns of Goll and Burdach, together with pigmentary degeneration of the cells of both anterior and posterior horns.

Spiller⁴ has emphasized the point that the degeneration in the cord is not a systemic but a diffuse process, and so resembles a toxic degeneration and does not essentially differ from that in anemia, except that in pellagra the nerve-cells are more affected. He concluded that the cause must be some toxic or infectious agent acting on all parts of the central nervous system causing cellular degeneration as well as nerve fiber degeneration in the posterior and lateral columns.

CHANGES IN THE GASTRO-INTESTINAL TRACT

The changes in the gastro-intestinal tract seem to vary chiefly in degree, being more marked in the acute forms, but in all cases in which definite lesions exist, they consist of chronic catarrhal inflammation with exfoliation of epithelial cells and infiltration of stroma. Thrash⁵ was in doubt whether or not this was a terminal condition as a result of the prolonged illness. He believes that the gastro-intestinal disturbances are secondary and due to the effort to excrete poisons ingested or formed from perverted metabolism and cell destruction. Atrophy of the muscular coats of the intestinal wall has been frequently mentioned^{6, 7} as present in a large proportion of cases examined, and rarely anemia or hyperemia of gastro-intestinal mucosa, as well as chronic enteritis.

Tuczek⁷ states that definite post-mortem results are found in the acute forms, typhus pellagrosus, in the form of chronic gastro-enteritis with formation of ulcers and swelling of mesenteric glands. This is associated with marked changes in the central nervous system and with general visceral involvement. Roberts³ in studying the pathology of pellagra noted atrophy of the muscular coats of the stomach and in acute cases found a true gastritis. The changes found in the intestine seemed to depend on the duration and severity of the disease,

6. "Im Verfolg Ihrer werten Zuschrift gestatten wir uns Ihnen die Mitteilung zu machen, dass es uns gegenwärtig unmöglich ist die für das Finkler'sche Heilverfahren in Betracht kommende Präparate Ihnen zur Verfügung zu stellen. Es haben sich nämlich Änderungen der betr. Verbindung notwendig gemacht, die uns augenblicklich noch beschäftigen."

* From the Pathological Laboratory, State Hospital No. 2, St. Joseph, Mo.

1. Niles: Pellagra, pp. 153, 158.

2. Marie: Pellagra, p. 262.

3. Roberts: Pellagra, pp. 142, 145.

4. Roberts: Pellagra, p. 152.

5. Niles: Pellagra, p. 159.

6. Marie: Pellagra, p. 269.

7. Marie: Pellagra, p. 274.

8. Roberts: Pellagra, p. 119.

consisting of atrophy of muscular coats and, in acute cases, of enteritis with ulceration, the latter chiefly in the colon and rectum.

The suggestion that the gastro-intestinal lesions found at autopsy are terminal and do not correctly represent conditions existing during life loses weight when we consider the clinical importance of symptoms from this source. The gastro-intestinal symptoms are the earliest,⁹ most persistent and dangerous of all manifestations of pellagra. Their importance is emphasized by the fact that their presence together with the skin manifestation is essential to a clinical recognition of the disease, and also by the fact that the gastro-intestinal and skin lesions are sufficient for a post-mortem diagnosis. In studying the priority of symptoms in a series of cases, one is impressed by the large percentage in which gastro-intestinal disturbance is the cause of first complaint.

In the series of cases under my observation, the prominence and distressing character of symptoms indicating derangement of the gastro-intestinal tract were very marked and constant. This fact, together with the rapid course of the malady, would class these cases in the acute type, typhus pellagrosus, the chronic forms showing seasonal recurrences and less active manifestations of the disease.

Four of the fatal cases of pellagra in this institution came to necropsy. A brief report of these cases and post-mortem findings, with reference to lesions of the digestive tract, may be of interest.

CASE 1.—A woman, aged 57, had been in this institution continuously for the past nine years and for most of the time for the past twenty-two years. There was no history or evidence of a previous attack of pellagra. About June 1, 1911, characteristic skin lesions of pellagra developed on the back of both hands and wrists, together with marked stomatitis and excessive flow of saliva—evidence of acute intestinal disturbance—diarrhea and persistent abdominal pain. The case progressed rapidly and death occurred July 10, 1911. Necropsy was done twenty hours after death. Examination of the abdominal viscera showed intense congestion of omental and mesenteric vessels. The ileum, cecum and ascending colon presented a marked catarrhal inflammation, in some areas becoming hemorrhagic in character. Aside from the general hyperemia the rest of the gastro-intestinal tract appeared normal.

CASE 2.—A woman, aged 64, had been a patient in this institution for the past thirteen years, during which time her record gives no history of a previous attack. About July 10, 1911, typical lesions of pellagra appeared on the dorsal surfaces of both hands, rapidly extending over the dorsal surface of the fingers and completely encircling both wrists (Figs. 1 and 2). The border of the lesion was distinctly reddened and the line of separation from the normal skin was clearly marked. The lesions were dry with a tendency to exfoliation; about the wrists there were numerous fissures. Areas of similar character soon appeared on the forehead and on the left side of the neck (Figs. 2 and 3). The mucous membrane of the mouth and tongue developed a reddened glazed appearance and there appeared a constant, excessive flow of saliva, filling the mouth and escaping in a continuous tiny stream. There was also a diarrhea, intermittent in character and apparently uninfluenced by diet or remedies. Nausea and vomiting occasionally accompanied the frequent bowel evacuations. The stools were small, liquid, brownish in color, and gave reaction for blood pigment. No parasites were found. Urinalysis and differential blood-count showed nothing of interest. The Wassermann reaction and blood-cultures were negative. This patient showed marked mental enfeeblement, but dementia was not deepened after the development of pellagra. She was conscious and recognized friends who visited her a few hours before death on Aug. 7, 1911.

Necropsy was done five hours after death. The condition to attract attention on inspection of the abdominal viscera was the very intense engorgement of all mesenteric vessels, giving to the intestinal wall a strikingly reddened, purplish appearance. The mesenteric lymph-nodes were markedly enlarged and of a soft pinkish color, resembling inflamed lymph tissue. The mucosa of the lower quarter of the esophagus and of the entire stomach showed a catarrhal inflammation of varying degree. The duodenum was distinctly hyperemic, while the mucous membrane of the jejunum and ileum was markedly inflamed. This condition varied in degree and became more intense in the lower portion. Peyer's patches were swollen. The appendix appeared normal. The inflammatory condition extended to the cecum and for several inches along the ascending colon; areas of congestion were observed along the transverse colon. The mucosa of the descending colon, sigmoid and rectum was about normal. No ulceration was observed.

Microscopic examination of sections of the stomach wall showed superficial necrosis with extensive destruction and exfoliation of glandular epithelium, infiltration between the gland tubules, frequently hemorrhagic, with cystic dilatation of the gland tubules. The vessels in the submucosa were greatly dilated and atrophy of the muscular layers was noted. This destructive process was not so far advanced at the cardiac as at the pyloric portion, where necrosis and hemorrhagic extravasation were very marked. In the pyloric portion the deeper branching parts of the tubules retained their epithelium, but in the superficial portion this was shed off, leaving a structure composed of areolar tissue, infiltrated with round cells and showing superficial necrosis.

Sections from the duodenum showed superficial destruction of the mucosa, the villi being fused together in a structureless mass with beginning necrosis. Except for some remnants at the bottom of the crypts of Lieberkühn, all glandular epithelium was lost. Brunner's glands were not involved.

Sections from the jejunum showed similar superficial exfoliation, but the form of villus could still be seen. At the bases of the crypts there were small remnants of epithelium, but on the tips of the valvulae conniventes necrosis extended to the submucosa. The muscular coats appeared normal except for infiltration between the layers.

In the lower portions of the ileum there was almost entire destruction of the mucosa; only in areas were epithelial remnants seen, a narrow line of partially necrotic tissue being left to mark its location. The muscular layers were atrophied so that in this region the intestinal wall was extremely thin. Sections taken from the cecum and ascending colon showed a destructive process similar to that in the ileum.

CASE 3.—A woman, aged 40, was admitted to this institution in an advanced stage of pellagra. Skin lesions of pellagra extended over the dorsal surfaces of both hands and wrists, small areas were present on the back of the neck and on the left ear, and an extensive lesion involved the vulva and the entire perineal region, extending up the inner surfaces of both thighs. The mucous membrane of the mouth and tongue was universally reddened with a few superficial ulcers; the tongue was swollen, showing marked indentations from the teeth. An excessive flow of saliva filled the mouth and overflowed in a constant stream. Involvement of the digestive tract was further evidenced by abdominal pain, diarrhea and complete loss of desire for food. The diarrhea was constant and uninfluenced by treatment. Differential blood-count, Wassermann reaction and blood-culture were negative. Death occurred seventeen days after admission.

Necropsy was performed two and one-half hours after death. A hyperemic condition was noted in all the abdominal viscera. The mucosa of the esophagus was apparently normal. A catarrhal inflammation was present along the entire tract from the stomach to the cecum, most intense in the lower ileum and cecum, where there were many areas of submucous hemorrhage. The transverse and descending colon, sigmoid and rectum were about normal.

Microscopic examination of sections from the stomach wall showed superficial necrosis and exfoliation; the deeper portions of epithelium were still intact. The vessels were greatly

9. Roberts: Pellagra, p. 107

dilated and numerous small hemorrhages into the mucosa were seen. Sections taken from different areas along the jejunum and ileum showed the same loss of epithelium, with only small remnants in the bottom of the crypts. In some places the villi retained their form; in others they were fused into a structureless mass. The muscular layers of the ileum were greatly atrophied. Specimens from the colon appeared to be about normal, except that in the cecum the catarrhal inflammation was very intense.

CASE 4.—A woman, aged 59, had been a patient in this institution for the past three and one-half years, with no history or evidence of a previous attack of pellagra. About Oct. 1, 1911, there appeared on the dorsal surfaces of both hands and lower forearms areas of dermatitis resembling sunburn; other lesions appeared on the right shoulder and on the side of the neck, and, as exfoliation progressed, the dark, purplish color with brighter red border gave the typical appearance of skin lesions of pellagra. Diarrhea began in a few days and continued with only slight remissions. An almost constant

the colon presented a catarrhal inflammatory condition of varying degree, quite marked in the duodenum and jejunum but most intense in the lower ileum and cecum, becoming hemorrhagic in some areas. The lower colon and rectum were only slightly affected.

Sections from the wall of the jejunum showed fusion of the villi into a necrotic mass and only the epithelium lying deepest in the crypts, intact. There was round-cell infiltration into

Fig. 1.—Patient, showing skin lesions encircling both wrists.

abdominal pain was complained of and a burning sensation, localized to the areas of dermatitis and to the mucous membrane of the mouth, was present, the sensitiveness of the latter interfering with the taking of food. The tongue and the mucous membrane of the mouth had the characteristic red, glazed appearance and there was present the same excessive flow of saliva. The feces were constantly liquid, of a brownish color and contained some mucus and varying amounts of blood. No parasites were found. Examination of the cerebrospinal fluid, differential blood-count and the Wassermann reaction were negative. Death occurred about two months after the first appearance of the disease.

Necropsy was performed fourteen hours after death. There were skin lesions on the dorsal surfaces of both hands, over almost the entire face and ears and on the right side of the neck and right shoulder. A condition of hyperemia of the abdominal viscera similar to that noted in the previous cases was present. The gastric mucosa was about normal. The intestinal tract from the duodenum to the hepatic flexure of

Fig. 2.—Same patient, showing lesions on dorsal surface of hands and fingers and on the forehead.

Fig. 3.—Same patient, showing lesion on left side of neck.

stroma of villi and between crypts. The muscular layers and the mucosa along the ileum were atrophied. This superficial erosion was not so marked in the cecum, but there was much extravasation of blood into the mucous membrane. The lower portions of the intestinal tract were only slightly involved.

In preparing the microscopic sections, tissue was taken from numerous places in the gastro-intestinal tract, so

that the findings would correctly represent the condition present. On the whole, the changes found were those of a chronic catarrhal inflammation, but in some areas the condition was very active and closely corresponded to the description given by Coplin¹⁰ of hemorrhagic inflammation of mucous membranes. Many of the sections gave a picture very similar to his illustration of hemorrhagic colitis from colchicum poisoning.

These findings would seem to coincide with the theory that toxic products cause pellagra, whether taken with food products or of endogenous origin. In the cases observed, no probable causative factor in the food could be found.

A NEW ETHER-VAPORIZER

A PRELIMINARY REPORT ON THE TECHNIC OF INTRAPHARYNGEAL INSUFFLATION ANESTHESIA

KARL CONNELL, M.D.

Instructor in Surgery, College of Physicians and Surgeons, Columbia University; Assistant Surgeon, Roosevelt Hospital

NEW YORK

The vaporizer (Fig. 1) herein described was devised to supply certain deficiencies in preexisting types of vaporizers for insufflation anesthesia. These deficiencies are inaccurate knowledge of the amount of air passing and of the percentage of ether vaporized. The present vaporizer indicates at all times the amount of air passing and makes possible the mathematical control of the ether percentage. Only by such knowledge can control and accuracy of dosage be secured and the effect of a given dose be foretold. Otherwise the patient must be used as the primary index of the amount of ether being delivered.

The essential feature of this apparatus is a device whereby the total air delivered is at all times indicated. It consists of an inverted aluminum cup (1, Fig. 2) riding in a glass tube (2, Fig. 2) on a central rod (3, Fig. 2). This rides at increasing heights as the air-stream, flowing from below, becomes more forcible, and is comparable to the familiar glass ball riding on a water-jet in a shooting-gallery. The float is of such weight, and the air-delivery hole of such size, that the float begins to ride on a delivery of 8 liters (nearly 8½ quarts) of air per minute, and indicates with fair accuracy, by the increased height at which it rides, all passage of air up to 22 liters (about 23¼ quarts) per minute, which is the range of utility in insufflation anesthesia. The float registers by its average height not only the amount of air passing, but by its fluctuation the respiratory movement of the patient. It drops instantly on any stoppage of the air-stream, as for example by a kink in the tube.

On the glass tube which houses the float, is engraved, at convenient levels, the number of liters of air passed per minute. Also opposite each level is engraved the number of drops of ether necessary to add per minute to make the air-ether mixture 14 per cent. by weight, with the float riding opposite that particular level, and also the number of drops which would make a 1 per cent. mixture. The amount of air used is thus constantly visible, and may be controlled in volume by adjusting a pinch-cock on the air-delivery tube, while the percentage of ether may be adjusted to any amount desired by controlling the number of drops of ether added per minute.

The ether is held in a reservoir of 250 gm. (half a pound) capacity (B, Fig. 2), patterned after a standard high-pressure oil-cup, wherein the drop of ether (4, Fig. 2) may be seen and accurately controlled by a screw-valve (5, Fig. 2). The reservoir is filled from the top (6, Fig. 2), and thrown into air-circuit with the vaporizing-chamber by an air-valve (7, Fig. 2). It is mounted on the vaporizing-chamber (C, Fig. 2), a heavy glass cylinder, 4 inches by 4 inches. The ether drops onto the copper bottom of the vaporizing-chamber, which is kept at a temperature merely warm to the hand by a 4-inch electric stove (D, Fig. 2), with adjustable heat. In the absence of electricity the stove and bottom of the vaporizer may be immersed in a pan of warm water. The ether immediately vaporizes on this warm surface and diffuses into the air, passing through the chamber, the resultant being warmed to from 25 to 30 C. (77 to 86 F.)

A mercury pressure-gauge (E, Fig. 2), a thermometer (F, Fig. 2) and a safety-valve (H, Fig. 2) are mounted on the chamber.

The float has been calibrated by passing from a high-grade-test-gas-meter accurately measured amounts of air, and registering each increase of 4 liters (4¼ quarts) per minute by a horizontal line on the glass tube housing the float.

The drop has been measured by repeated count, averaging under the condition of this vaporizer 29.3 drops per cubic centimeter or 41 drops per gram.

Fig. 1.—New ether-vaporizer.

Knowledge of the amount of air which floats the aluminum cup at a particular level enables one to calculate the weight projected each minute at the cup. It is then a problem in simple arithmetic to arrive at the number of drops of ether to be added to the air-ether mixture whereby the percentage of ether-vapor by weight will be any selected amount.

Omitting the complex physical formulas and mathematical problems by which these results have been arrived at, and noting merely the results of the estimation: Dry air at 0 C. (32 F.) weighs 1.2928 gm. The conditions in the air used for calibration were those under which air would usually be delivered to the vaporizer: temperature at 22 C. (71.6 F.) under 30 mm. positive pressure and a water-vapor content of 1.5 per cent. by volume. Under these conditions the weight of air is 1.238 gm. per liter of air. If, for example, the float registers by its height the equivalent of 16 liters of air per minute, the equivalent in momentum of 19.84 gm. is projected at the float. Then to give 14 per cent. ether vapor, liquid ether running 41 drops to the gram, an amount of 114 drops of ether per minute would be added to the air-stream with the float at the 16-liter

10. Coplin: Manual of Pathology, p. 561.