The Boston Medical and Surgical Journal

TABLE OF CONTENTS

February 1, 1917

BOSTON SURGICAL SOCIETY

DIVISION OF THE URETER IN PELVIC OPERATIONS. <i>By William P. Graves, M.D., Boston</i>	149
FRACTURES OF THE EXTERNAL CONDYLE OF THE HUMERUS IN CHILDHOOD, WITH ROTATION OF THE CONDYLAR FRAGMENT. <i>By James S. Stone, M.D., F.A.C.S., Boston</i>	151
THE APPLICATION OF THE MEDIAN PATELLA INCISION FOR A KNEE ARTHROTOMY. <i>By E. G. Brackett, M.D., Boston</i>	153
THE MANAGEMENT OF OPERATIVE CASES PRESENTING URINARY BACK PRESSURE. <i>By Arthur L. Chute, M.D., Boston</i>	155
ACUTE AND SUBACUTE PERFORATIONS OF THE STOMACH AND DUODENUM AT THE MASSACHUSETTS GENERAL HOSPITAL. <i>By Edward P. Richardson, M.D., F.A.C.S., Boston</i>	158
ORIGINAL ARTICLE	
APPENDICITIS. <i>By Torr Wagner Harner, M.D., Boston</i>	165
CLINICAL DEPARTMENT	
A LABORATORY AID IN THE DIAGNOSIS OF SCARLET FEVER. <i>By D. M. Lewis, M.D., New Haven, Conn.</i>	170
CANCER OCCURRING IN ACID PARTS OF THE BODY. <i>By A. L. Benedict, A.M., M.D., Buffalo</i>	172

NEW INSTRUMENT

AN IMPROVED VENIPUNCTURE NEEDLE. <i>By Guy G. Fernald, M.D., Concord, Mass.</i>	178
---	-----

EDITORIALS

THE MEDICAL PROVISIONS OF THE YOUNG BILL.....	174
CHICAGO CONFERENCE ON MEDICAL EDUCATION.....	175
THE ELIMINATION OF THE RAT.....	176
MEDICAL NOTES.....	176

OBITUARY

EDWARD MARSHALL BUCKINGHAM, M.D.....	177
--------------------------------------	-----

MEMORIAL RESOLUTIONS

MEMORIAL RESOLUTIONS FOR DR. KEANY.....	177
---	-----

MISCELLANY

HUBONIC PLAGUE IN BOSTON.....	177
DECLINE OF POPULATION IN FRANCE.....	178
THE YOUNG INDUSTRIAL HEALTH INSURANCE BILL.....	179

CORRESPONDENCE

THE ANTI-VACCINATION CAMPAIGN. <i>Samuel B. Woodward, M.D.</i>	180
--	-----

Boston Surgical Society.

DIVISION OF THE URETER IN PELVIC OPERATIONS.*

BY WILLIAM P. GRAVES, M.D., BOSTON.

Dr. Scudder has asked me to report in greater detail the cases of division of the ureter that I mentioned in the clinical meeting last month at the Free Hospital for Women.

CASE 1. Operation October 31, 1907. The patient was a married woman of 45. The operation was for an immense fibro-adenocystoma of the uterus, which completely filled the abdomen. The tumor was densely adherent, as these rare tumors practically always are. The urine before operation was normal. The tumor was freed on the right side, the ureter being exposed along most of its course and not injured. The ureter on the left was found to enter the tumor at a point about an inch above the brim of the pelvis. On account of the length and severity of operation, it was thought best not to attempt to dissect the ureter from its bed in the tumor. The ureter was, therefore, deliberately cut above the brim of the pelvis, and again at a point near its entrance into the bladder. Inasmuch as the right kidney felt normal and the right ureter was intact, the proximal stump of the cut ureter was tied with strong silk about an inch from the end. The cut end was then sewed over with catgut. The proximal stump next to the bladder was tied with catgut. The patient made a good recovery from the operation, but had complete anuria for 36 hours. She was treated with hot packs, nitre and pilocarpin. At the end of 36 hours she began to pass urine and excreted 31 ounces in the following 24

hours. From that time she made an uninterrupted convalescence. There was no pain or tenderness of the left kidney and no apparent hydronephrosis. The opposite kidney, however, underwent a very marked compensatory hypertrophy which was noted within a few days after the operation. The patient was followed for several years, and remained in perfect health. The right kidney continued to be hypertrophied.

CASE 2. Operation October 31, 1907. The patient was a single woman of 27. The operation was for a large cervical fibroid which was tightly packed in the pelvis, extending principally into the right, broad ligament. In cutting the right uterine vessels the right ureter, which was displaced from its normal position, was completely severed. After removal of the fibroid uterus by a complete hysterectomy, the left kidney and ureter were inspected and found to be intact. The injury to the right ureter was too high to allow for convenient implantation in the bladder, so that the only alternatives were anastomosis or tying of the ureter. On account of the recent experience from the preceding case, in which the patient had had anuria for 36 hours, it was decided to perform an anastomosis. An end-to-side anastomosis of the severed ureter was performed in the classical way (illustration on black-board). The patient passed 11 ounces of urine during the 12 hours following operation, and continued to pass a normal amount thereafter. Nine days after the operation bladder symptoms appeared with rise of temperature, malaise and pus in the urine. This attack, which was evidently a pyelitis, lasted two or three days, recurred 10 days later, and again recurred 10 days after that. Each attack of pyelitis was short in duration. After discharge from the hospital, the patient continued in excellent health, her case being followed for several years.

* Read before the Boston Surgical Society, January 10, 1917.

CASE 3. Patient was a single woman of 58. Operation, November 14, 1916. The operation was for a very large pseudo-mucinous cyst filling the abdomen and densely adherent. The worst adhesions were in relation to the right ureter, which could not be definitely distinguished. Great care was exercised to avoid cutting the ureter, but when the uterus was removed, it was found that the ureter was held within two clamps and almost completely severed. In this case the injury was sufficiently near to the bladder to allow for implantation, so that the operator had a choice of implantation, anastomosis or ligature. In considering the experience of the two former cases of cut ureter, it seemed that the patient in whom the ligature had been employed had made the better convalescence. In view of the fact that the patient had undergone a long and severe operation, and in view also of the favorable reports of ligature, especially from the Mayo Clinic, it was decided to tie the ureter. In this case three separate ligatures of No. 2 chromicized catgut were placed at about one inch apart in order to secure complete closure. The purpose of using several ligatures placed at a distance was to allow for a certain amount of necrosis of the ureteral epithelial lining with the formation of plastic adhesions. This patient made an uneventful recovery. There was no palpable hydronephrosis on the right side, nor was the compensatory hypertrophy of the opposite side sufficient to be detected by examination. There was no pain or tenderness of the kidney on the side of the ligated ureter.

CASE 4. In addition to the three personal cases which I have cited, I will add one from the service of one of my colleagues. The operation was for a large cervical fibroid, during the enucleation of which one of the ureters was cut. In this case the ureter was implanted into the bladder. Soon after the operation a urinary fistula appeared per vaginam. The fistula persisted for several months, after which the patient was lost sight of.

When a ureter has been accidentally cut, the surgeon usually has the choice of several procedures. Ligating the ureter has certain advantages over anastomosis and implantation. It requires very little expenditure of time, usually an important consideration, as the operations in which the ureter is cut are generally of great severity. If the end of the ureter is properly tied there is little danger of leakage and fistula.

According to Barney's statistics, hydronephrosis occurs in 80 per cent. but the hydronephrosis is as a rule not great and gradually subsides as the kidney atrophies.

If a dangerous hydronephrosis should occur the kidney may be removed at a later operation.

Anuria appears, according to Barney, in about 1.6 per cent. after ligation of one ureter. It is probable that ureteral anastomosis is more commonly followed by fistula and infection than is the ligature operation. It is evident from the literature that most of the cases of death following ligature of the ureter have been due to shock following some extremely severe operation, rather than to the closure of the ureter.

If the choice is between anastomosis and implantation I should personally prefer an anas-

tomosis. By anastomosis only one organ is injured. If a fistula or pyelonephritis ensues the kidney may be removed and the trouble cured by one operation.

If a fistula follows implantation in the bladder the surgeon then has to deal with two injured organs. The operation for reimplantation or closure of a ureterovesical fistula is, in my opinion, less satisfactory than a simple nephrectomy, such as might be necessitated by trouble from ligature or anastomosis.

If implantation is imperative, as might be the case in the presence of injury of the other ureter, or a functional deficiency or disease of the opposite kidney, the place of choice would be, of course, in the bladder, if the proximal end of the ureter is sufficiently long.

In case of emergency the ureter may be left to drain from the vagina or carried out through the skin.

Implantation into the intestine is not desirable on account of the almost inevitable ascending infection of the kidney.

Implantation into the appendix or of one ureter into the other, and even into the Fallopian tube, are operations that have been advocated from experiments on animals and on the cadaver, but probably have no very practical value.

DISCUSSION.

DR. WILLIAM L. ESTES, S. Bethlehem, Penn.: My experience in division of the ureter has been of a very limited order. I have done anastomosis in two cases, and tied off in one. One of the anastomoses did perfectly well, and one, I think, died, rather from shock of operation than severance of the ureter.

I should like to say in regard to the remark of Dr. Graves concerning infection of ureters implanted into the intestines, that I have done an implantation into the rectum for extrophy of the bladder in a child nine years ago, and the child is to-day doing very well.

DR. COBB: I have had four cases in which I divided, unintentionally, the ureter, three were carcinoma of the cervix and one a very large fibroid in a negress. I did an anastomosis after the method just sketched by Dr. Graves; the first case died of shock largely due to the increased time taken by the anastomosis. One case of carcinoma recovered uneventfully and in two cases, one of carcinoma and one of fibroid, I had to do a nephrectomy four or five weeks after the operation.

DR. RUSHMORE: One of the major operations of gynecology has been described as tying four arteries and one ureter, and it has been said that one does not qualify as a gynecologist until one has actually cut across the ureter. However that may be, the subject of Dr. Graves' paper is of great interest and importance to everyone who does many pelvic operations.

I have had experience in three cases in which I cut the ureter. In one, in Baltimore, in which the operation was for a large myoma of the uterus, Dr. Kelly performed the anastomosis. The method was that just described by Dr. Graves: end-to-side. The patient did perfectly well.

The second case was operated on for carcinoma of the cervix with involvement of the ureter. I resected the ureter and implanted the proximal end into the bladder. The third was for carcinoma of the cervix. By accident I cut across the ureter, and then tied it. I think it would have been better to have left it alone because the patient was in such poor general condition and suffering from shock; to have left it alone would have given a little better chance.

One point in regard to the method of implantation into the bladder. I prefer some modification of the method suggested by Dr. Coffey. The incision is made first through the muscle, down to the mucosa, then the mucosa is punctured a short distance above the lower end of the incision. The ureter is best cut obliquely, not directly across, so as to give a sort of flap and that flap is drawn down into the bladder with a fine piece of catgut needled at both ends. The bladder wall is punctured by the needles from within outward and the ureter anchored by a suture, the knot of which lies outside the bladder. The bladder wall is then inclosed around the ureter and part of the muscle may be cut away to give less pressure.

The result is to approximate rather closely the valve-like opening that is naturally found between the ureter and the bladder.

In regard to results in these three cases, the first case did perfectly well. The case of carcinoma of the cervix in which the ureter was resected for about an inch and a half and implanted into the bladder, did perfectly well. A cystoscopic examination with catheterization of the ureter showed no evidence of disturbance of function. The third case died about thirty-six hours after the operation from shock.

In anastomosing ureter to ureter, a danger is the occurrence of a stricture, which predisposes to pyelitis.

DR. A. L. CHUTE: I cannot agree with Dr. Graves in the procedure of tying off a cut ureter without making an attempt to see whether the kidney cannot be saved. I feel that one should see whether something cannot be done in the way of a conservative operation, before tying off the ureter and destroying the kidney, as this means. If too much of the ureter has not been removed, it is perfectly possible to reimplant it into the bladder. I prefer this, if it can be done, rather than to make an anastomosis between the cut ends of the ureter.

I believe it is possible, if it be necessary, to free the ureter from its bed for at least one-half its length. This will allow bringing the ureter to the bladder comfortably in most cases. In women,—and of course all these cases occur in women,—the reimplantation is simpler than in men. I pass a number 10 or 12 ureter catheter to the bladder, make a small incision in the bladder wall at the place where I mean to implant the ureter and fish my catheter out through this incision. I then pass the catheter up the ureter for 5 to 8 cm., after which I introduce the ureter containing the catheter into the bladder for about a half inch, sew up my incision about my ureter, and catch the ureter to the

bladder wall at two or three points. There is little tendency to leakage in my experience, although I usually do carry a wick down to the point of suture. This procedure may be carried out rapidly. I have used it after resecting a tumor of the bladder involving the ureter, and in one case where the ureter had been injured during hysterectomy. I had an opportunity to see the latter case some months after operation, and found that she had a perfectly patent ureter, showing no particular contraction. Whether we can hope for this result in most cases or not, I do not know, but I feel we should make some effort to save the kidney in these instances where a ureter has been accidentally cut.

In all, I have seen four cases where the ureter has been tied off, severed, or wounded in doing hysterectomy. In the first case, which I saw a considerable time after operation, an abdominal urinary fistula had developed. A nephrectomy was done and the patient did well. In the second instance, both ureters had been tied off. I saw the patient about fifty-six hours after operation and did a double pyelostomy. The patient died some three days later. I have never been quite sure as to the cause of her death, since she was said to have passed a fair amount of urine following the double pyelostomy. I saw the third patient five weeks after operation. All the urine from the left side leaked through the vagina. This is the reimplantation that I have already referred to. The fourth case is one that I saw recently, and in this instance I believe the injury to the ureter was due to a needle prick. The amount of leakage has diminished; it looks to me as though this patient would require nothing in the way of operation, but that the injury would heal of itself.

DR. GRAVES: (Dr. Graves, in closing, spoke of the technic of implantation.) A very rapid method of implanting the ureter into the bladder is to split the proximal end of the ureter as one would a dandelion stem. The split end is then carried into an opening in the bladder by guide sutures. Two or three fine catgut sutures are used to make the joint more secure. (Illustrated on the blackboard.)

Another method of implantation, if the ureter is too short to reach the bladder, is that suggested by Albarran. A flap is turned up from the bladder and wrapped about the end of the ureter, the opening in the bladder being sewed up by a running suture. (Illustrated on the blackboard.)

In order to avoid stricture at the point of implantation Stoeckel suggests the following: Instead of making a slit in the bladder a round hole is cut about the end of a clamp which has been inserted through the urethra into the bladder cavity. The mucous membrane and peritoneum are sewed together before the ureter is implanted. (Illustrated on the blackboard.)

FRACTURES OF THE EXTERNAL CONDYLE OF THE HUMERUS IN CHILDHOOD, WITH ROTATION OF THE CONDYLAR FRAGMENT.*

BY JAMES S. STONE, M.D., F.A.C.S., BOSTON,
Surgeon to the Children's Hospital.

OPERATIVE intervention in fractures involving the elbow joint in children is ordinarily of very

* Read before the Boston Surgical Society, January 10, 1917.