Miscellany

Prophylactic Use of Quinin in Malaria

Surgeon H. R. Carter, in a recent issue of *Public Health Reports*, thus summarizes the results of his study of this question:

The use of quinin in small doses is an efficient method for preventing malarial fever.

This method is especially adapted for use in a farming community where it is not practicable economically to get rid of malarial mosquitoes or to properly screen against them.

In malarious states quinin should be taken in doses of from 5 to 7 grains per day by grown persons, 2 to 3 grains by children—less if small—during the malarial season, say, June to November. Somewhat smaller doses will be efficient in places where the malaria is not bad. If begun in March or April, it would prevent a certain number of relapses.

In these doses it does no injury and does not produce blackwater fever or cause malaria to be more difficult to cure by quinin.

It can be taken in these doses by almost all persons without discomfort, or with discomfort for only the first seven or eight days. In cases in which it does produce discomfort a smaller dose should be tried and, if possible, increased later.

The insoluble salts of quinin are better borne—that is, cause less discomfort—than the soluble. The tannate is the most insoluble and is said to be the best borne. Made up with chocolate and sugar into tabloids it is best suited for administration to children. As the tannate contains a smaller proportion of quinin than the sulphate, from two to two-and-a-half times as much must be taken to get the same effect.

In addition to the use of quinin, whenever it is practicable to do so, the house should be screened, brush and high weeds near it cut away and pools and wet places drained or filled. If the last is not practicable, wet places and pools should be oiled.

In a community using quinin prophylaxis there will still be a certain number of cases of malarial fever. Some of these will be failures of the method, but others, and the majority if it be used properly, will be relapses of previous malarial infections from the last season. In the Italian army in 1911 there were 304 relapses and 186 new cases to 10,000 men. A certain number of relapses are prevented by quinin prophylaxis, but not all.

The Surgery of the Ancients

A set of thirty-seven remarkable ancient Greek surgical instruments, discovered near the site of Kolophon, in Ionia, after being a considerable time in the possession of the late Mr. Alfred O. Van Lannep, Dutch vice-consul at Smyrna, has been brought to England. The instruments show a type of workmanship unequalled in any other extant specimens, and generally reveal the great progress in surgery which the ancients achieved. The date is probably the first or second century, A. D. With two exceptions, all the instruments are of bronze. The blades of the knives were originally of steel, but this metal has been almost completely destroyed by oxidation. Among the five pairs of forceps in the find is a large and beautifully made instrument, the handles of which are shaped to represent two dolphins. This is probably a pair of polypus forceps used for removing growths. The bite of the teeth is strong and close. Another interesting pair of forceps is that used for extracting arrow- and lance-heads from wounds. This pair somewhat closely resembles modern bone-forceps, and has artistically modeled handles and blades which possess strong teeth for gripping. An elevator for raising depressed bone is another interesting exhibit. The collection would seem to prove that after battle, efforts must have been made to treat surgically even the most serious wounds of the skull. Another brain instrument is the "drill-bow" for operating a skull trephine. This instrument proved a great puzzle at first, and was supposed to represent some sort of measuring appliance. Research, however, has shown that it is a bow which, when fitted in a string, was used to rotate a bone-drill or skull trephine.

The British Museum possesses the only other specimen of the drill-bow extant, and until the discovery at Kolophon was made the nature of that instrument was not known. The collection includes tenacula similar to those now in use, a number of catheters of beautiful workmanship, shaped like modern instruments of the same type, a scoop or curet for gynecologic and other work (Hippocrates speaks of such an instrument), a cautery for burning wounds, a couple of probes exactly like the modern ones, a couple of spatulas or spoons, a needle-holder, and a bronze box intended evidently as an instrument case. In addition, there is a slab of Egyptian porphyry for mixing ointment on, and a well-constructed pair of scales, still in excellent equipoise. Some cupping-vessels were also discovered. This interesting collection of instruments has been given to the Johns Hopkins University and will shortly be taken to America. Models, however, are being made and will be on view in London.

Correspondence

Minnesota Regulations on Embalming-Fluid

To the Editor:—I am surprised that Dr. Jacob Rosenbloom should not have more fully informed himself on embalming-fluids now used throughout the country before writing his communication which was published in *THE JOURNAL*, June 13, 1914, p. 1912, and am rather surprised that in the comment on this communication, more attention was not given to the condition in Minnesota, the regulations of which state, as I understand it, have been used as a basis for the regulations in several other states.

There have been regulations governing the use of certain chemicals in embalming-fluids in the state of Minnesota for some time. The present regulation (No. 33), amending some of these older ones, reads as follows:

After Sept. 1, 1909, every embalming-fluid sold or used in Minnesota must contain formaldehyd gas in the proportion of 5 per cent. by weight of the gas in every 100 parts by volume of the fluid, and must not contain any ingredient that interferes with the germicidal action of formaldehyd.

Such embalming-fluids must not contain chloral, nor shall they contain arsenic, mercury, zinc or other mineral poison.

Regulation 34 reads as follows:

After Sept. 1, 1909, a list of ingredients in every embalming-fluid sold or used in Minnesota must be on file in the office of the Minnesota State Board of Health.

These regulations have the force of law.

H. M. BRACKEN, M.D., St. Paul.

Executive Officer, Minnesota State Board of Health.

A Simple Method of Vaccination

To the Editor:—In *THE JOURNAL*, June 6, 1914, p. 1829, Dr. Shalet suggests that vaccination can be performed by dipping a sterile toothpick in glycerinated virus and abrading the arm by a rotary scarification. I have tried this method, with a sharp, chisel-shaped, hardwood toothpick, and have found it entirely satisfactory.

My serious objection to the original technic of Pirquet is the vaccination through the drop of lymph (or tuberculin). The whole point of our method is to make a circular scarification with a dry instrument on a dry skin. Then the toothpick is used to rub the virus (or tuberculin) into the abraded area. As skins are very different in thickness I find it much easier to judge the amount of pressure necessary to produce scarification if the skin and chisel are both dry. If lymph is dropped directly from a capillary tube on the scarified spots the chisel can be used for rubbing in the virus, thus eliminating the toothpick entirely. We use the toothpick because our virus comes to us in bulk and we wish to use the minimum quantity for each vaccination.

J. N. FORCE, M.D., Berkeley, Cal.

Assistant Professor of Epidemiology, University of California, Department of Hygiene.