on animals it was not demonstrated that a continuous anterior stitch tended to contract the opening, since it is applied after the jejunum has become wrinkled, such is the case on the cadaver. In trying the ligature care should be taken not to include the posterior suture within the loop. Since it was possible to include 5 cm. of the wall of the small intestine of a cat the larger human bowel will permit of the formation of an opening up to any size which may be required to side track the stomach contents.

Conclusions.—These experiments show that a successful anastomosis between the stomach and intestines, or two loops of intestine, is possible by means of the elastic ligature. The method is not applicable when an immediate opening is required.

The time of the cut-out varies, depending upon the character of the ligature, the method of application, and the resistance of the tissues in any case, but the ligature will ultimately cut out.

A knot at either end of the elastic loop increases the rapidity of the cut-out.

The pinching of the gut by the ligature seems to cause no bad symptoms.

The serous surface of the outside of the joint when completed is smooth.

The anastomosis by gross or microscopic examination shows no evidence why there should be any greater tendency to contraction of the opening than with the ordinary methods of intestinal suture.

The procedure causes a minimum operative risk, both as regards shock and infection.

Adhesions across the opening are a possible, but not a probable, complication.

Extending these conclusions to practical operative work, it is my opinion that in cases in which an immediate opening of the bowel is not imperative, and in which the avoidance of operative shock is an important factor, the elastic ligature may prove to be the operation of choice in gastro-enterostomy and lateral anastomosis.

BIBLIOGRAPHY.

- ¹ Gaston, J. M., *Atlanta Med. and Surg. Jour.*, 1884-5, Vol. i, p. 336: 385, and 1885-6, Vol. ii, p. 395: 533.
- ² Bardenheuer, F., *Exper. Beiträge z. Abd. Chir. Inaug. Diss.* 1888.
- ³ Terrier et Baudouin, *La Suture Intestinale*, p. 100.
- ⁴ McGraw, T. A., *Jour. Am. Med. Ass.*, 1891, Vol. xvi, p. 685.
- ⁵ Cordier, A. H., *Jour. Am. Med. Ass.*, 1891, Vol. xvii, p. 263.
- ⁶ Podres, A., *Arch. f. Klin. Chir.*, 1898, band, lvii s. 358.
- ⁷ Modlinski, *Verhandl. d. Deut. Gesell. f. Chir.*, 1899, Pt. i, s. 68.
- ⁸ Cutts, R. E., *Northwest Lancet*, 1901, Vol. xxi, p. 480.
- ⁹ Kakels, M. S., *Med. Rec.*, 1903, Vol. lxii, p. 556.
- ¹⁰ Maury, J. W., *D. Med. News*, 1903, Vol., lxxxiii, p. 483.

DIFFERENTIAL DIAGNOSIS OF RHEUMATISM AND THE ARTHRITES.¹

BY THOMAS F. HARRINGTON, M.D., LOWELL, MASS.

THERE is, perhaps, no class of affections more commonly met in practice than those included in the title of this paper. Notwithstanding this marked prevalence, the ancient history of its

¹ Read before The Massachusetts Medical Society, North Middlesex District, Oct. 28, 1903.

recognition, and the great progress in scientific medicine, the undisputed fact remains that there is no one term in medicine accountable for more ignorance and neglect than "rheumatism." Undefined itself, it has spread its mantle over conditions dissimilar and obscure until it has become well-nigh impossible to separate the true from the spurious. It is the hope of pointing out some of the errors and evil consequences of this classification that prompts me to present the subject at this meeting.

In the first place rheumatism and arthritis are not synonymous terms. The characteristic symptom of rheumatism is an arthritis, yet we have many arthritides which cannot be associated with rheumatism as clinically accepted. Every arthritis should be traced to its probable true cause, and no case should be considered rheumatic unless an undisputed attack of acute rheumatism can be given. Rarely is rheumatism diagnosed other than it is; the danger lies wholly in the opposite direction. As will be shown later, there is probably but one form of rheumatism, namely, acute rheumatism. Any other use of the term leads to confusion, and often to wrong treatment, with disastrous consequences.

It is with the object of avoiding such errors, as well as an endeavor to offer a classification founded upon clinical appearances, re-enforced by etiological and pathological findings, as far as known, that I would suggest the following:

A. Infective arthritis: (1) acute rheumatism; (2) gonorrheal arthritis; (3) purpuric arthritis; (4) pneumococcic arthritis; (5) scarlet fever arthritis; (6) tubercular arthritis; (7) syphilitic arthritis; (8) typhoidal arthritis; (9) influenza arthritis; (10) erysipelas arthritis; (11) spinal meningitis arthritis; (12) smallpox arthritis; (13) dysenteric arthritis; (14) glanders arthritis; (15) pyemic arthritis. B. Arthritis deformans. C. Osteo arthritis. D. Traumatic arthritis. E. Gouty arthritis. F. Hemophilic arthritis. G. Senile arthritis.

On account of the confusion and disputation existing on the question of classification I would offer the following reasons for the above grouping.

Idiopathic arthritis probably never occurs. In infective arthritis there is usually more than one agent, and the effect of the agent is modified by various conditions,¹ e. g., the resisting power of the individual, the dose of the toxin received, and by the presence of other organisms or their toxins. The synovial membrane of joints has a marked tendency to become involved in many of the acute infectious diseases.

This shows itself in four different forms²:

First.—By a transient synovitis with limited effusion, e. g., typhoid, scarlet fever, influenza.

Second.—Those with marked swelling and effusion of fluid into the joint which later becomes turbid and purulent.

Third.—Those having involvement of the peri-articular structures, e. g., gonorrheal.

Fourth.—Those cases in which the arthritis is acute and destructive from the start, suppura-

tion is early and violent, *e. g.*, pneumococcic infection.

It is very probable that all organisms which cause arthritides produce other pathological conditions as well.³ While there are some exceptions yet unexplainable, it can be stated as a general rule that all acute arthritides are secondary manifestations. In doubtful cases the probability of mixed infection must be borne in mind. The role played by the nerves in the course and development of joint affections is very great. The close relation existing between acute rheumatism and chorea, as well as the positive factors, sex, age, mental and nervous fatigue, shock, etc., are, especially in arthritis deformans, open up a wide field for investigation.

In the above classification rheumatism is placed under infective arthritis because it is now generally understood that this symptom complex is due to some form of micro-organism. Walker⁴ has isolated a micro-organism which he calls "micrococcus rheumaticus." Other factors tending to establish this relationship are⁵: its endemic and often epidemic nature depending upon climatic conditions; its many manifestations, variety of lesions, fever and sweating resembling tuberculosis and pyemia; its tendency to recur, especially upon exposure, resembling pneumonia and erysipelas; its liability to relapse like enteric fever, and to end spontaneously as do the exanthemata. Then, too, the close relationship between affections associated with follicular tonsillitis and those associated with rheumatism, namely endocarditis, pericarditis, pleurisy, chorea, erythema nodosum, as well as other cutaneous and subcutaneous effusions such as purpura, pneumonia, peritonitis and profound anemia. The occurrence of arthritis, pleurisy, etc., with puerperal sepsis also points strongly to the septic association of rheumatism.

There seems to be as little foundation for the term rheumatic diathesis, or uric acid diathesis, under such a classification as there would be to speak of a pneumonic or septicemic diathesis. The uric acid condition has been demonstrated in athletes in whom no suspicion of rheumatism or gout existed.⁶

In the classes arthritis deformans and osteoarthritis a distinction is made between those cases in which there is a true bony enlargement of the joint and those in which the deformity is due to an atrophy of the surrounding tissues, causing an apparent rather than a real enlargement. The former class is a true osteoarthritis and is so-called. The latter is a deformity due in all probability to some trophic change depending upon disturbances of the nervous system.

The clinical signs and pathological findings of each are different and characteristic. As the treatment of the two affections is very different it is essential that this distinction be recognized. It is the failure to do this that has resulted in so many permanent crippling, also salicylic acid poisoning.

Whether arthritis deformans, like some other

conditions, results later in osteoarthritis does not alter this classification. The point is that all joint affections which have true overgrowths should be called osteoarthritis, the suffix deformans being understood; the term arthritis deformans being limited to the conditions characterized by appearances quite typical and constant to be described later. Much confusion in reading textbooks and medical literature will be avoided if some such distinction is maintained, and if the term rheumatism is limited as signifying a clinical picture unlike any other joint affection, rather than an arthritis occurring in the course of many diseases. This would eliminate all such terms as muscular rheumatism, tuberculosis rheumatism; scarlatinal, gonorrheal, gouty, typhoidal, etc., with the suffix rheumatism.

DIAGNOSIS.

Infective arthritis. — There is usually a history of a primary infection to which the arthritis is secondary.

In the case of acute rheumatism, the following signs are characteristic: (1) *History.* — Often follows tonsillitis, especially the suppurative form. In the majority of cases the previous health is good and there is entire absence of any constitutional disturbance. In 270 cases occurring at Johns Hopkins Hospital, 60% occurred between ages of eleven to thirty years; 30% occurred between the ages thirty to fifty years. There was a family history of rheumatism in 25% of the cases; 45% had had previous attacks.⁷ The onset is sudden and without a chill in 94% of the cases in the Johns Hopkins series. Cold, damp weather and exposure are factors but do not cause rheumatism.

(2) *The character of the inflammation.* — Rapid development of pain, swelling and heat in one or more joints. In twenty-four hours after onset the temperature ranges from 102° to 104°, with a pulse which is more apt to be under than over 100. The temperature is not remittent, thus differing from most of the other arthritides, and it has very slight depressing action on the central nervous system.

(3) *Pain.* — As it attacks a new joint it always leaves the old one, and it is this flitting from joint to joint which often serves to distinguish it from other diseases.

(4) *Results.* — It runs an acute course, averaging three to six weeks, with a marked tendency to get well. It does not deform the joints, *i. e.*, the joint is left as uninjured as it was before the attack. The endocardium is involved in fully 60% of the cases. It tends to recur. It never suppurates. It rarely becomes chronic.

In 100 cases of alleged chronic rheumatism lately examined by the writer, only 19 gave a history of a previous acute rheumatism. In 14 of these, closer questioning or the result of the examination threw considerable doubt upon the nature of the primary attack.

The term "chronic" relates entirely to time, therefore chronic rheumatism, so-called, implies

a long continued or slowly progressing rheumatism. This we know is not the nature of infective rheumatism.

The term *recurrent* is more truly descriptive of the majority of these cases. Such cases may eventually result in deformity, but they then become osteo-arthritis, and should be so called and treated. Any case of acute arthritis which leaves the joint impaired, or which fails to yield to salicylates should excite suspicion as being something other than pure rheumatism. Such cases will usually be found to be due to some other toxin or to some blood or nerve dyscrasia.

Among the conditions most commonly called chronic rheumatism are⁸: (1) relaxed ligaments (flat foot). (2) Occupation neuroses, which are usually worse in damp weather, and which are not peculiar to any special occupation, but occur when faulty positions throw unusual amount of work upon any one set of muscles. (3) Neuritis, due to circumflex nerves sending trophic fibers into the joint. (4) Gout which is more common in this country than supposed, *i. e.*, it is three-fourths as frequent at Johns Hopkins as at St. Bartholomew Hospital, England; most common among lead workers and beer drinkers. The differential diagnosis of acute rheumatism and acute gout is very difficult. The gouty nodules in the ears, or tophi on the fingers are characteristic of gout. (5) Multiple neuritis is often mistaken for rheumatism, but the absence of knee-jerk and the alcoholic history, as well as the future course of the disease serve to differentiate them.

Chorea and acute rheumatism are closely allied. The consensus of opinion to-day seems to point to chorea being a manifestation of cerebral rheumatism in which the infective agent has acted upon the serous membrane of the joints and brain, and has also reached the cortex itself, thus producing convulsive movements.⁹ This is in contra distinction to the theory that some infective agent acted directly upon the nerves themselves. The increased reflexes noted in rheumatism are probably due to the same cause.

In gonorrheal arthritis the history is important. It is often limited to one joint, especially the left knee, and is not as painful, has less fever, and does not go from joint to joint. The inflammation is in the peri-articular surface and extends along the tendons. The effusion is rarely purulent, although suppuration of the sheaths may be found. This is important in a bacteriological examination of the joint as colonies of gonococci can be obtained in cultures taken from the outgrowths on the synovial membrane while the fluid fails to show their presence.

In ophthalmia neonatorum arthritis has occurred which yielded promptly and completely on curing the conjunctival discharge. In the multiple gonorrheal arthritis it is often hard to differentiate. It is usually confined to the lower extremities, and often attacks the sterno-clavicular, the intervertebral, the temporo-maxillary and the sacro-iliac articulations, joints rarely in-

involved in acute rheumatism. The possibility of this form of arthritis in newly married women infected by an old gleet of the husband should be borne in mind. Gonorrheal arthritis is most intractable and this often serves to distinguish it from rheumatism. It is far safer to regard atypical or doubtful arthritis as being of gonorrheal origin than due to rheumatism.

There is nothing characteristic in the blood in rheumatism, notwithstanding the marked anemia so often associated with it.

In arthritis occurring with purpura, the existence of urticarial wheals in addition to other signs of purpura is usually sufficient to distinguish it from rheumatism. Sore throat and even pericarditis may occur in this form of arthritis.

Arthritic manifestations in pneumonia indicate a general pneumococcic infection and have a mortality of 60%.¹⁰

The arthritis occurring in scarlet fever is not rheumatic. Recent investigators¹¹ ascribe the joint trouble to streptococcus pyogenes, and consider it one of the manifestations of scarlet fever, rather than a rheumatism complicating the disease. It usually attacks the wrists and fingers, or other small joints. In the monarticular form there is a greater tendency to suppuration.¹² The arthritis occurs most often at the end of the second week before desquamation sets in.

Lately French writers¹³ have described arthritis occurring in connection with lupus, as tuberculous rheumatism but which had better be called tubercular arthritis. This is painful and always slow in developing, although there is usually a history of traumatism from which the family date the trouble. The family history of tuberculosis or the presence of tuberculosis in other parts of the body is important. There is usually very little fever. The joint enlargement is soft and symmetrical. Monarticular rheumatism is more often regarded as tuberculosis than the contrary. A mixed infection of these two conditions is not uncommon. Joints in defective positions, with abscess or fistula, accompanied with muscular atrophy, justify the diagnosis tubercular arthritis. This form of arthritis closely resembles the syphilitic arthritis. The latter is usually limited to one joint and is painless, or painful only at night. A characteristic point about this class of syphilitic affections is that the amount of movement preserved by the joint is far greater than in any other joint affected by the same amount of disease. In doubtful cases the history and therapeutic tests are important aids.

Syphilis attacks the joints symmetrically and it rarely relapses, thus differing from rheumatism.

As to arthritis of the spine associated with typhoid, the conditions most likely to resemble it are myelitis, sciatica, hip-joint disease, tumors of pelvic bones, bone disease, malingering or muscular pains, and pachymeningitis.

Arthritis occurs in connection with influenza

erysipelas, smallpox, appendicitis, spinal meningitis, dysentery and glanders, but in none is there much resemblance to acute rheumatism with its sudden onset, its joint to joint evanescence, never suppurating, and always leaving perfect joints. There is, however, one form of infective arthritis which closely resembles acute rheumatism, and upon the differential diagnosis of which serious consequences depend. I refer to *acute osteo-myelitis*.

There is, perhaps, no more serious condition in which rheumatism should be positively excluded than in acute-osteo myelitis. We must be able to recognize this disease in the first forty-eight hours if we expect the best aid from the surgeon. There is a pocket of pus in the bone and if it is not immediately drained, absorption, high temperature, delirium and septicemia follow, resulting often in death, or at the very best in destruction of the bone.

The symptoms that distinguish acute osteo-myelitis from acute rheumatism are classical. Chill, acute pain followed by high temperature, sepsis with its symptoms of absorption, delirium and death. The pain is usually in the extremities of long bones. Usually, especially in children, there is not an effusion into the joint near the focus in the first forty-eight hours. This is different from rheumatism in which effusion occurs in first few hours. The greater intensity of local symptoms, the involvement of the epiphysis rather than the joint, and the more constitutional disturbances are all characteristic.

In the first forty-eight hours pain and sepsis increase but there is no local sensitiveness. Deep and continued pressure for two minutes elicits excruciating pain and frequently indicates the seat of the infection.¹⁴ There is a board-like infiltration of skin and muscles with subsequent involvement of joint and suppuration. Most of the cases occur between puberty and twenty-five years. The favorite seats are head of humerus, lower end of femur, and upper end of tibia. Monarticular pain in children should suggest osteo-myelitis. If there are scars of former operation over the bone the pain is probably a second attack of osteo-myelitis. In many cases there is an existing furuncle, traumatism without external lesions, skin wounds, or one of the infective diseases.¹⁵

In doubtful cases aspiration is justifiable. The disease may simulate typhoid fever. Blood examination may be helpful in the diagnosis. There is a decided leucocytosis in the early stages of osteo-myelitis, thus differing from typhoid and rheumatism.

No acute arthritis occurring in connection with a possible pyogenic infection should be looked upon as rheumatism or gouty until osteo-myelitis has been positively excluded.

The chronic form of osteo-myelitis is often diagnosed chronic rheumatism, but the localization of the pain in the bone rather than in the joint, and the evidence of necrosis will often serve to differentiate them.

Arthritis Deformans.

This is a good descriptive term and has the advantage in eliminating all reference to rheumatism and gout. It is probably due to some nerve disturbance, and is an apparent enlargement of the joints due to the atrophy of the muscles and inter-articular cartilages. It closely resembles rheumatism in its acute stage, and may occur in combination with acute rheumatism, but it is not caused by rheumatism. It is non-inflammatory and may occur without pain. There is seldom redness or tenderness. It is rarely monarticular, and it never suppurates. There are no true bony enlargements. The characteristics of arthritis deformans are: the spindle-shaped appearance of the joints, its soft boggy feeling, its symmetrical development, beginning in the small joints and spreading without leaving the first joint, its stationary character, never flitting from joint to joint, and always leaving the joint deformed. It is less painful than rheumatism and has a tendency to attack young and middle aged women, especially when there has been any wear and tear of the nervous system. It often comes on insidiously, lasts months, and does not involve the heart. The process stops grating and ankylosis, with resulting deformity, is left. The deformity is usually one of extension or flexion, seldom lateral. In the polyarticular form of any arthritis which does not leave the joint uninjured, especially if the pain and disturbance are more annoying when the patient is worried, or under a nervous strain, we should suspect this form of arthritis. Gout is excluded by the absence of deposits of urates. The so-called Charcot joint is in all probability one of this class. It is readily recognized by the history of tabes as well as the increased lateral movement of the joint without pain. Intermittent hydrops,¹⁶ characterized by painless swelling of sudden onset, without heat and redness, is probably due to some vasomotor neurosis.

The form of arthritis described by English observers,¹⁷ as acute rheumatoid arthritis, confounds acute rheumatism and arthritis deformans more often than any other affection. In this disease occurring in young women a distinction is made in the swelling beginning in the proximal phalangeal joints, extending beyond the affected joints, and becoming fusiform in shape. Other joints become affected until the patient is helpless. There is much muscular wasting or atrophy. It runs a definite course, lasting about three weeks. It attacks the fingers first and spreads to other joints, involving the temporo maxillary and spinal joints, and it leaves the joints immobile and enlarged, points which clearly eliminate acute rheumatism. Its association with influenza, parotitis, and the like, would suggest a microbic origin, acting no doubt through the nervous system, producing trophic changes in the joints, resulting in arthritis deformans. It is neither rheumatic nor osteo-arthritic.

There are seldom any adhesions in arthritis

deformans. Contractions and stiffening from faulty posture, and extensive degenerative erosions of the cartilages may result. These must not be mistaken for true bony growths which are the characteristics of osteo-arthritis.

Osteo-Arthritis may result from many causes. It may start without any known cause. The disease is essentially a chronic disease, the so-called acute painful period being due in all probability to a double infection. At first there is very slight deformity, and often the person is unaware of any enlargement on account of the gradual development of the disease. The pain is more often at night and is considered rheumatic. It usually affects the large joints (knee), and is often unilateral. It is a true bony growth, "ring bone," and locks the joint mechanically by the formation of osteophytes at the ends of the bones. Change of position often aggravates the pain, especially in the spinal form, and the person walks with a characteristic gait. This serves to distinguish this form from lumbago. The disease is more common in men, especially old men, but it does occur in young people. The true nature of the disease is obscure. Cold and damp may be factors in the etiology, but they do not cause the disease. There is less atrophy of muscles and less synovial swelling than in arthritis deformans. The enlargements are bony and not cartilaginous. The joint capsule is relaxed. The deformity is due to the hypertrophied tissue, as well as to enlarged bone. These hypertrophies have a tendency to form on the dorsal side of the joint, causing a lateral deformity or flexion, thus differing from the deformity of arthritis deformans which is usually one of hyperextension.¹⁸

In osteo-arthritis due to the presence of Heberden's nodes a very slight injury will often set up an active process resulting in increased deformity and a permanent disability far in excess of that expected from a light blow. The presence of Heberden's nodes in other joints will often confirm this fact. Such nodes suggest strongly the presence of osteo-arthritis, rather than arthritis deformans, and are valuable in deciding the diagnosis, prognosis and treatment in a given deformity.

In arthritis deformans there is practically a false ankylosis which is amendable to proper manipulations, while in osteo-arthritis any manipulations of the joint would most likely break off some of the osteophytes with increased irritation in the joint.

A form of osteo-arthritis occurring in the spine, *spondylitis*, is far more common than supposed.

There may be little or no subjective symptoms referable to the back, and there may be much overgrowth of bone without any marked deformity. The deformity is greater in the slowly developing cases than in the more active process, and its nature depends upon the surface of the bone affected. These cases are usually diagnosed lumbago or rheumatism. The different motions of vertebral articulations should

be tested in every case of backache, especially when such is aggravated by change of position, and associated with neuralgic pain, or sensory disturbances in the legs. Even in the early stages there is a "spot" in the spine at which the pain is localized. It can be differentiated from Pott's disease by the pain being local, rather than referred or abdominal, as in Pott's. In doubtful cases the use of tuberculin and the Rontgen rays are helpful. If paralysis has supervened it is peripheral in spondylitis while that due to Pott's disease is paraplegic.¹⁹

In spondylitis the paralysis is seldom equal on both sides. There is often an associated ankylosis and fixation of the thorax. As the disease advances the tissues shrink and pressure on the cord and nerve roots is released. This results in the abatement of the nerve symptoms, and is often mistaken for an improvement in the osteo-arthritis with an unfortunate relaxation in following the prescribed treatment.

Traumatic Arthritis. — The history of the case is important, care being taken that the trauma is not the exciting cause of an osteo-myelitis or other infectious arthritis, rather than the cause of a simple arthritis. There is usually water in the joint in traumatic cases. Joint mouse or floating cartilage belongs to this class.

Gouty Arthritis. — Many arguments can be offered in support of the theory that gout is due to the destruction of leucocytes in their battle with toxins rather than to uric acid.²⁰ This brings it nearer to the infective group of arthritides. The characteristics of gouty arthritis are sudden onset, especially at night, often disappearing in day time, also the presence of deposits in the ear (tophi). It attacks usually single small joints. A distinction should be made between a scanty urine with high specific gravity which is permanently so, and one with the same characteristic dependent upon exercise, diet, etc. In chronic gout ulcerations often occur about the finger joints, from which urate of sodium crystals can be collected.

Hemophilic Arthritis. — This should be kept in mind especially in arthritis suggesting surgical interference. A recent writer reports 252 cases.²¹ Traumatism does not seem to be an aetiological factor. The joint becomes distended with blood. There is a feeling of tension and pain, but no tenderness or symptoms of inflammation. Echymosis in a few days followed with apparently full recovery. This condition soon recurs and results in thickened and roughened synovial membrane, structural changes in the joint, muscular atrophy and resulting deformity resembling osteo-arthritis. The disease occurs usually between the fourth and sixth years and most often in males. The joints affected are knee 50%, elbow 25%. The mortality in hemophilics is less among those having arthritis than among the others. It is 60% in the latter class.

In *Senile Arthritis* there is a thickening of the capsule, but not of the bones. It has no relation to rheumatism and is more stiffness than

soreness. It does not involve the heart and grows better as the day progresses.

Rheumatism in children differs somewhat from that of adults. The tonsils are the most common source of suspected infection. The arthritic symptoms are not as well marked as in adults, and are more apt to come on insidiously. Recurrent tonsillitis, chorea, erythema, or nose bleed, are often associated with arthritis. Continued fever with slight remissions suggests endocarditis.²² There is not the same successive involvement of joints seen in adults. The family history, or a history of "growing pains," especially if accompanied with even slight swelling (fibrous nodules) suggests rheumatism.

The distinction between rheumatism in adults and that of children is best told by Cheadle, in a single sentence:

"The various manifestations of rheumatism massed together in the case of adults tend to become isolated in the case of children, so that the whole phenomena are distributed over years, instead of weeks or months, and the history of a rheumatism may be the history of a whole childhood."²³

REFERENCES.

- ¹ Bradshaw Lecture, Lancet, Dec., 1902.
- ² Marsh, British Medical Journal, Dec. 13, 1902.
- ³ Webster, Journal American Medical Association, Jan. 10, 1903, p. 82.
- ⁴ Walker, Guy's Hospital Gazette, London, August, 1903.
- ⁵ The Hospital, London, June 27, 1903.
- ⁶ Lambert, American Medicine, June 6, 1903.
- ⁷ The Journal American Medical Association, Jan. 24, 1903.
- ⁸ Walsh, Ibid, Aug. 29, 1903.
- ⁹ British Medical Journal, Aug. 29, 1903.
- ¹⁰ Slaughter, American Medicine, April 18, 1903.
- ¹¹ Monti, Kinderheilkunde, Vol. ii, p. 571.
- ¹² Bunce, The Philadelphia Polyclinic, 1897.
- ¹³ Bonveyron, Revue de Chirurgie, Paris, September, 1903.
- ¹⁴ Murphy, Journal American Medical Association, July 25, 1903.
- ¹⁵ Gonser, Jahrbuch für Kinderheilkunde, 1902, Vol. lvi.
- ¹⁶ Brackett and Cotton, Boston Med. and Surg. Journ., 1901, Vol. cxlv.
- ¹⁷ W. Hale White, Guy's Hospital Report, 1902, Vol. lvii.
- ¹⁸ Goldthwait, Boston Med. and Surg. Journ., Jan. 28, 1897.
- ¹⁹ Goldthwait, Ibid, Aug. 10, 1899.
- ²⁰ Woods Hutchings, Lancet, Jan. 31, 1903, p. 288.
- ²¹ Corless quoting Poillet of Lyons, Practitioner, 1903, Vol. lxx, p. 85.
- ²² Freedlander, Cleveland Medical Journal, July, 1903.
- ²³ Quotation Crandell, Transactions of the American Pediatric Society, 1902, Vol. xiv, p. 59.

Clinical Department.

REPORT OF CASES OF DECAPSULATION OF THE KIDNEY.¹

BY F. G. BALCH, M.D., BOSTON.

THE following cases are reported in the endeavor to add a little to our knowledge of the results obtained by various surgeons in their efforts to cure a disease, which until recently has been considered purely medical, by surgical interference.

CHRONIC DIFFUSE NEPHRITIS. DECAPSULATION.

CASE I. Ellen W., thirty-three, married. Laundress. Entered Massachusetts General Hospital, Nov. 5, 1901.

Personal history.— Unimportant, excepting she has for several years drunk one-half gallon of beer daily. Subject to headaches and vomiting spells.

¹ Read at the annual meeting of the Suffolk District Medical Society, October 31, 1903.

Present illness.— For past two or three weeks severe headache, vomiting and swelling of feet and abdomen. Urine scanty. Diarrhea persistent.

Examination.— Well developed but obese. Heart and lungs negative, except for few râles in back. Much edema of feet, legs, thighs, back and face, with marked ascites. Pulse 70, small volume, low tension. *Urine.*— Acid, 1,009, albumin 2½%. Sediment, blood, chiefly abnormal, many fatty renal cells, and hyaline, granular and brown granular casts. Quantity in first three days, less than 10 oz.

With daily hot air baths, pilocarpin and purgative treatment she improved slowly during first two months. Amount of urine rising to from 25 to 40 oz. per day. Albumin falling to ½% on average. Casts and blood corpuscles fewer in number.

During winter very slight improvement. Ascites required tapping.

March 24, 1902. Less edema and ascites.

Occasional headache and vomiting. *Urine* 1,012, smoky. Albumen ½%. Sediment as before, with occasional waxy cast. In spite of this apparent improvement patient's general condition became so much worse that operation seemed advisable.

Operation, March 28 on both kidneys. Tissues edematous, kidneys small, dark, granular in appearance; patient stood operation well. During first three days after operation twenty-four-hour amount of urine rose from 20 to 40 oz. Improvement after operation slow, but less headache and vomiting.

May 1. Less edema, and in legs and feet only, but ascites continues; tapped and 8 qts. of fluid drawn. *Urine* shows fewer casts, hyaline and granular, with fat; fewer renal cells and blood corpuscles; albumen ¼%.

June 1. Patient stronger, appetite improved; still occasional headache and vomiting. *Urine* pale, about 1,010, average 30 oz. in twenty-four hours. Albumen, large trace. Casts same as before, with renal and fatty renal cells, few blood corpuscles. Returned to medical ward June 6. Hot air baths renewed and diuretics, which raised amount of urine to an average of 40 oz. Abdomen requires tapping about once in three weeks. Discharged July 20, 1902, when she felt well, but vomited every few days; had no edema and very little ascites. *Urine* pale, 1,011; albumin 1¼%. Occasional hyaline casts; renal cells, with fat; occasional blood corpuscles.

Patient returned to hospital in October, 1902. Had felt well since operation and had been doing housework daily, and said that she was better than she had been for a long time.

Urine then showed ¼% of albumen, with a sediment of a few waxy, many granular casts, with fat adherent; few red corpuscles, and compound granule cells. Patient lost sight of and subsequent history unknown excepting that she died in May, 1903, after considerable excess in alcohol.

CASE II. Mary B. L. Twelve years old.

Etiology.— She had measles, mumps and scarlet fever as a child. She has always been well and strong. Last February caught cold and was at home a week. Two days later her feet swelled and she was edematous below her eyes. She had no headache or disturbance of vision. She had been under treatment since April 16.

Examination.— The tension of the pulse was somewhat increased. There was slight dullness in the flanks shifting on change of position. There was edema of the feet and legs and under the eyes. *Urine.* In the medical wards before transfer she had been secreting from 8 to 10 oz. of urine a day. On the