cusses whether in the present status of our knowledge it is possible to determine the seat of the sensations of hunger and thirst, replying to his question in the affirmative. He explains that hunger and thirst are phenomena common to the cells of every living being as manifestations of the lack of chemical elements indispensable for the life of the cells. These trophic sensibilities of the cells may be unconscious in the lower animals but they become conscious phenomena as the nervous system of vertebrates becomes more highly developed. In man, hunger is manifested by painful contractions of the stomach, and thirst by dryness of the mouth and throat. These phenomena are purely reflex, the principal centers for hunger and thirst being in the medulla oblongata, but Auerbach's plexus is also involved, especially in relation to hunger. He discusses in detail the phenomena from lack of lime, etc., in man, animals and birds, and the way in which amphibious animals change their diet as they live in water or air.

Smith, M. I. AUTONOMIC DRUGS AND THE STOMACH. [Am. Jl. Phys., 46, 1918, 232.]

Pilocarpine causes contraction of all parts of the surviving stomach. *Atropine* antagonizes pilocarpine and relaxes. *Nicotine* causes contraction of all parts except the fundus and the cardiac sphincter of the cat and the antrum and body of the rabbit's stomach. *Adrenaline* causes mostly relaxation, but in the guinea-pig, rabbit, and dog, some parts may be contracted. The sphincters are always contracted by it. *Ergotoxine* causes those parts of the dog's stomach which are contracted by adrenaline to relax. The conclusion is drawn that the sympathetic innervation is wholly inhibitory in cat and man; inhibitory for certain regions, augmentator for others, in guinea-pig, rabbit, and dog. The sympathetic innervation of the sphincters is augmentor. [W. M. B., Phys. Abst.]

Bañuelos, M. CELIAC PLEXUS REFLEX. [Rev. Med. y Cirugía, March 14, 1919.]

Pressure on the celiac plexus, the author claims, modifies the rhythm of the heart and raises the arterial blood pressure. The respiration, pupils, and likewise the tone of the peripheral vessels are stimulated. The respiration is increased by five or ten inspirations per minute and the pupils dilate. The latter occurs more strikingly in sympathicotonic individuals. Vagotonics are not affected. This reaction is marked in hyperchondria, pylorospasm and gastric ulcer, and other vagotonic states.

Pagniez, P., Valléry-Radot, P., and Nast, A. TREATMENT OF MIGRAINE. [Presse Méd., April 3, 1919.]

These authors had noticed two years ago, in studying a case of giant urticaria, that ingestion one hour before meals of a small amount of the