THE ORIGIN OF THE RAMMELSBERG ORE DEPOSIT.

WALDEMAR LINDGREN AND JOHN DUER IRVING.

INTRODUCTION.

In the course of a brief visit to the great copper deposit of Rammelsberg on the northern slopes of the Harz Mountains the authors had an opportunity to study the occurrence and mineralogy of its ores. The results of such a short investigation are, of course, entirely insufficient as a base for a detailed description. Nevertheless, it appeared to them as if certain relations not previously emphasized might help to elucidate the difficult and much discussed question of the genesis of that ore deposit. For kind permission to visit the mine the writers are indebted to Oberberghauptman Krümmner in Clausthal and to Bergrath Wolff, of Goslar. For our guidance in the mine we are under obligations to Obersteiger Voigt, who directs the exploitation of the deposit.

The Rammelsberg deposit, as shown on the accompanying map, Fig. 47, is situated on the northern slope of the Harz Mountains in close vicinity to the town of Goslar. As well known, it has been worked for copper ores since ancient times, the first records dating back to the tenth century. Its geological structure has been investigated by a number of authors, but its complete and detailed description is as yet a problem of the future. The most diverse explanations have been offered as to its mode of origin. By some, perhaps by a majority, including such well-known geologists as Bergeat and Klockmann, it has been considered as a sedimentary deposit contemporaneous with the surrounding sedimentary rocks. Others, like Professor Vogt, following Freiesleben and Lossen, explain it as a deposit from solutions immediately derived from igneous magmas. Still others, like Professor Beck, occupy a non-committal standpoint.

Geology and Structural Features.—The deposit is enclosed, apparently conformably, in Devonian rocks, which at Goslar appear as an overturned anticlinal and dip towards the north. It lies in the so-called Goslar slates of the Middle Devonian; these slates being again overlain by a thick series of Lower Devonian *Spirifer* sandstone, which makes up the summit of the Rammelsberg Mountain, at the foot of which the mine is located. The following cut (Fig. 48) after Klockmann will show these relations.

The slates have suffered considerable contortion and the ore

FIG. 47. Sketch map showing location of Rammelsberg.

body apparently follows these contortions more or less clearly. The underground developments extend over a horizontal distance of about 2,000 meters and have attained a vertical depth from the surface of 380 meters from the level of the Richtschacht. It will be seen from this that mining has not yet penetrated to great depths in spite of the fact that the deposit has been worked for nearly a thousand years. The ore body is divided horizontally into two parts; respectively referred to as the old and the new

beds; they are connected by a narrow and contorted seam showing, however, beyond doubt that the two are really parts of one deposit.

The following sketch, after Klockmann shows the relation of these two ore-masses as seen in plan.

FIG. 48. Cross-section of Rammelsberg ore deposit showing overturned anticline and manner in which the deposit is believed by earlier writers to be conformable to the lamination of the enclosing slates. (After Klockmann.²)

The thickness of the ore bed, as it is generally referred to, varies considerably; in places it swells to dimensions of as much as 30 meters, but this is rather due to folding and local enlargement. In most places where we observed it in the workings the thickness did not extend over 2 or 3 meters and was sometimes

FIG. 49. Plan of the Rammelsberg ore body showing the relation of the old and new ore body. (After Klockmann.)

only 0.5 to 1 meter. The dip is vertical uniformly 45° to the southeast. In depth the ore in general appears to become a little richer in zinc and barite. At present the ores are sorted by hand at the surface and transported by wagons to the railroad, about

one mile distant. The older methods of working are now being replaced by more modern ways of exploitation. One of the main difficulties in the treatment of the ore lies in the fine grained intergrowth of chalcopyrite, zinc-blende and galena, which is eminently characteristic of the deposit.

It is stated, even in the modern descriptions of the Rammelsberg, such as that of Professor Beck,¹ in the third edition of his book on ore deposits, that the bedding or schistosity absolutely conforms with the outline of the ore and with its banded structure. The ore is said to take part in all the folding and contortions of the slate.

This idea of a mass of ore conformable to the schistosity of the slates, which has for so long been accepted in regard to this ore body, is shown in the well-known cross-section (Fig. 48) of the older of the two ore lenses. The slaty lamination is here shown curving around the irregular offshoot or apophysis which is figured in this section.

In view of the widely circulated opinion of this conformity the writers were surprised to find that, while true in places, it is certainly not a general characteristic of the deposit. In part, as was first noted, and called to their attention by Bergrath Wolff, it is absolutely unconformable to the stratification of the slate.

The ore *banding* follows in all cases with great faithfulness the outlines of the sulphide mass whether these be long smooth borders or small detailed irregularities of outline. For much of the distance above the third level the *edge* of the new ore body, and consequently the banding of the ore also, is indeed parallel to the lamination of the enclosing rock. On the third level, however, the ore mass flattens out and crosses the lamination at an angle of only a few degree, to again turn down parallel to it after intersecting the laminae for a very considerable distance. In this portion of the mass the banding of the ore follows the edge of the sulphide mass and makes therefore the same angle with the lamination of the slates as the outline of the ore mass itself.

¹ Beck, "Lehre von den Erzlagernstätten," 3d ed., Vol. 2, pp. 134-138.

The relations are shown in the sketches I and II of Fig. 50 below:

Where the ore crosses the lamination of the slates it is about five feet in width and shows no differences in either material or structure from that portion which is parallel to the structure of the enclosing rock. The outline of this ore body is not a plane but a warped surface which shows broad gentle undulations of large radius. Occasionally a sharp inequality is present and the ore veers off in a projection or apophysis only a few inches in

FIG. 50. Sketch of new ore body, 3d level, Rammelsberg. I shows manner in which the ore body, which is conformable with the lamination of the enclosing slates crosses this lamination at the point *A*. II is an enlarged sketch of the transition zone *A* showing how the banding of the ore truncates the lamination of the schists and how this lamination terminates abruptly at the ore and continues in the same direction on the other side.

cross section for a distance of 8 to 10 inches into the slate. One of these is shown in Fig. 51.

In this case the gneissoid banding of the ore still follows the outline of the periphery and runs into the apophysis with a series of embayment shaped lines, which exactly parallel in their arrangement the lines drawn on a map to indicate bodies of water which follow faithfully the outline of the shore absolutely irrespective of the structure or nature of the material of which the land may be composed.

In another place on the third level the new bed is folded into the form of an S as shown in Fig. 52. Here also the gneissoid

FIG. 51. Small embayments about eight inches wide, showing how banding of ore follows smaller inequalities in the enclosing rock.

banding follows with perfect 'faithfulness' the periphery of the mass while the schistosity is abruptly truncated by it.

FIG. 52. Sketch of S-shaped fold in new ore body, 3d level, Rammelsberg mine. Shows how banding of ore truncates the lamination of the enclosing rock.

In all cases where the ore bodies are set at an angle to the bedding the laminae of the slates extend up to and terminate abruptly against the sulphide mass and thus continue in the same

direction on the opposite side. They do not enter it nor are they in the least distorted against it.

The Ores.—A number of varieties of ores are recognized. The principal minerals are zinc-blende, chalcopyrite, galena, pyrite and arsenopyrite; the quantity is approximately proportional to the place of the minerals as enumerated. The gangue is almost entirely barite, but rarely appears in large quantities,

FIG. 53. Lenticular augen, Rammelsberg Ore. Shows also fine banding and intimate intergrowth of minerals.

and often is entirely inconspicuous. Masses and veinlets of calcite are often found in the surrounding slate, but rarely contain ore. On the whole the limits of the "ore bed" are sharply defined and the ores themselves are entirely or predominantly composed of sulphides. An alteration of the enclosing slates is rarely observed. At most there is a slight im-

pregnation of pyrite. The so-called *kupferkniest*, which adjoins the ore in certain places, is a siliceous argillite which contains disseminated pyrite and chalcopyrite. It has often been con-

FIG. 54. Lenticular augen of pyrite in Rammelsberg Ore. A A A A A = Pyrite lenses, B = —.

sidered as an impregnation, later than the ore itself, but, as far as we could observe, there is no sufficient basis for such an interpretation.

The structure of the ore varies with the locality, and the composition also varies distinctly so that in one place the "ore bed" will contain predominant copper values, while in other places there is scarcely any copper, but only zinc-blende with a little galena.

The intergrowths of these minerals are very fine-grained and extremely intimate. By far the most common structure is that of

FIG. 55. A large nodule of barite and sphalerite in Rammelsberg Ore, showing banding of the ore and gneissoid structure.

the so-called Melir-Erze, which are intimately banded, in most cases consisting of dominant zinc-blende with narrow and gently curved streaks of chalcopyrite and galena. The appearance of the ore is reproduced in Figures 53 and 54. In places the ore contains rounded nodules, generally of pyrite, around which the fine-grained streaks of zinc-blende and chalcopyrite bend in regular curves. This is shown in Fig. 54, which also brings out the fact that the pyrite is in part drawn out into long streaks. In general, however, it shows a notable resistance to such deformation. Not uncommonly one finds rounded nodules consisting of

zinc-blende and barite in granular form, coarser than that of the ordinary ore. Fig. 55 illustrates an occurrence of this kind, which clearly shows how the streaks of other sulphides surround and envelop these nodules. The pyrite nodules have often been noted, but without satisfactory explanation. Some observers have held them for fossil remains, in which the pyrite has replaced the shell of the organism. A goniatite has actually been found in the slates forming the easterly continuation of the ore body, and, according to K. Andree,¹ the interior of the shell contained pyrite, quartz, barite and calcite.

Origin.—The structural relations of the ore body indicate, according to our view, without any doubt, that the deposit is a bedded vein; that is, a fissure vein lying in part conformable to the surrounding slates. The distribution and structure of the ore itself is inconsistent with the theory of sedimentary deposition. As far as our experience goes the structure is unique in ore deposits, but as to its interpretation there can be no reasonable doubt. The sulphides do not occur with their primary texture. The structure is that of a dynamo-metamorphic rock, in which all of the constituents, except pyrite, have been drawn out into streaks which are intricately mingled. The appearance shown in the figures could be easily duplicated from any fine-grained gneiss area resulting by pressure from an original granular rock. The different constituents have acted under pressure as plastic material and are thoroughly mashed and squeezed. It is difficult to avoid the conclusion that the sulphide mass has moved almost like a plastic clay and that the present structure is due to this movement. There is no evidence, in our opinion, to justify the assumption of sedimentary deposition.

At first glance it seems strange that the pyrite has acted so differently from the other constituents. The explanation of this behavior is easily found in some interesting experiments recently undertaken by Professor F. D. Adams.² In this memoir Pro-

¹ Andree, K., *Z. f. prakt. Geol.*, 1908, p. 166.

² Adams, F. D., "An Experimental Investigation into the Action of Differential Pressure on Certain Minerals and Rocks," *Jour. of Geol.*, 18, 1910, pp. 480-535.

fessor Adams shows that plasticity is a function of hardness, so that the minerals become less plastic as they become harder. He finds that the limit of easily produced plasticity lies in the vicinity of five or six in the scale of hardness. Pyrite in his experiments was crushed under a load attaining 43,000 pounds without showing any trace of plastic deformation. Minerals of lower hardness presented decided evidence of plastic flow. The pyrite nodules in the Rammelsberg ores are then simply residual parts of the original granular deposit which have been less deformed than the other sulphides.

In conclusion then, we regard the Rammelsberg deposit as one of epigenetic origin, in which the structure of the ore has been profoundly changed by dynamo-metamorphism. While the surrounding slates are soft they evidently behaved quite differently from the sulphide mass, which seems to have flowed almost like a thick mush between the slate walls. The reason for this difference in behavior is not as yet fully explained, but we believe the question could be settled experimentally by a method similar to that of Professor Adams.

The association of minerals with barite as predominating gangue tends to show, in our opinion, that the deposit is not of the deep seated type, but was formed at a medium depth below the original surface, probably, within a few thousand feet of it.

Regarding the source of the ores, the only suggestion we would like to offer is that the deposit may have been formed by ascending solutions derived from the neighboring batholith of granite, which is only three kilometers distant from the mine.