ous purposes, such as fitness for the open-air classes, mental retardation and throat and teeth examination. Without burdening you with many figures, some few may be of interest concerning the retarded pupils. These children had all been in the same grade more than a year, and it was thought best to see if there were any marked and readily found physical causes which might be a factor in the mental backwardness. One hundred and fifty-eight pupils were examined.

Forty-one, or 25.3%, had defective vision; 2, or 1.2%, had defective hearing; 122, or 77.2%, had chronically enlarged tonsils; 127, or 80.3%, had defective or carious teeth; 34, or 21.5%, had marked pallor (the Hb not estimated); 3, or 1.8%, were markedly under-sized; 1, or 0.6%, was extremely nervous without definite signs of chorea.

Practically all of these children have been sent to the proper sources for treatment of these various conditions. It is of interest to compare the result of the examination of the teeth and throat of these children with that of 212 children of normal mental caliber looked over last year.² Seventy-two per cent of these children showed chronic enlarged tonsils; 50% showed carious teeth. Comparing, we have 212 average children; 72% had chronic tonsils; 158 retarded mental development; 77% had chronic tonsils. Two hundred and twelve average children, 50% had carious teeth; 158 backward children, 80.3% had defective or carious teeth. It would seem from these statistics that chronically enlarged tonsils and the generally accompanying adenoids had little to do with the mental backwardness of these children, but that carious teeth might play a considerable rôle in the causation of this condition.

In conclusion, no one is more alive to the inadequacies of the present system of school inspection than the school physicians themselves. At the same time, a large amount of work of direct benefit to the children has been done by school inspection.

Dr. Durgin and those directly associated with him in the Board of Health are the only ones who have an intimate knowledge of the truth of this statement. The general public is absolutely ignorant of the true state of the case and the public press seemingly willfully ignorant of it.

When the public is better educated as to the necessity and importance of school inspection, better results may confidently be expected.

THE State Health Department has had made a list of the lepers now in the state of Texas, who number 30. Of these, 11 are reported married and 9 single; 10 are severe cases, 2 medium and 6 mild; 20 are said to be Americans and 2 Mexicans; 16 are male and 6 female. As to location, 11 are in Galveston, 8 in San Antonio, 3 in Dallas, 3 in Sourlake and 1 each in Liberty, High Island, Del Rio, Comfort and Austin. They vary in age from thirteen to sixty-eight years. — *Jour. Am. Med. Asso.*

² Coates, W. B.: The Results of the Clinical Throat Examination of Two Hundred and Twelve School Children. *BOSTON MED. AND SURG. JOUR.*, vol. cxlix, p. 215.

TUBERCULOUS MENINGITIS, WITH LEUCOCYTOSIS AND A PURULENT EXUDATION CONTAINING MANY BACILLI. A CASE REPORT.

BY I. CHANDLER WALKER, M.D.,

Medical Intern, Carney Hospital, Boston, Mass.

(From the Medical Clinic of the Carney Hospital, Dr. Henry A. Christian, Physician-in-Chief.)

If the spinal fluid obtained by lumbar puncture is turbid and if this turbidity is found to be due to polymorphonuclear leucocytes, one is naturally inclined to the diagnosis of acute meningitis caused by some one of the pyogenic cocci. In some cases it is very difficult to find typical cocci in meningeal fluids, so that the absence of cocci in the smears might not lead one to suspect some other bacteria not easily stained by the ordinary methylene blue stains. In this way it is quite possible to overlook a case of tuberculous meningitis presenting a turbid polymorphonuclear leucocytic exudate, especially since most of the textbook and system discussions of tuberculous meningitis emphasize strongly the non-purulent character of the spinal fluid and the prominence of lymphocytes in the fluid. That cases, however, do occur showing the above described type of spinal fluid caused by the tubercle bacillus is emphasized by the following case.

P. B., age nineteen, admitted to the Carney Hospital Oct. 6, 1910.

History (obtained from patient's mother). — The patient had been perfectly well, mentally normal and had worked as a plumber's apprentice up to the middle of July, 1910. At this time he became ill with "typhoid pneumonia." He was confined to his bed for five weeks. During this illness the patient had no abnormal mental symptoms. He appeared to have perfectly recovered from the illness and worked the following three weeks. On Sept. 3, 1910, he became ill with "pneumonia." In the beginning he had pain in his left chest, and a few days later he complained of severe headaches unaccompanied by vomiting. On Oct. 3, the patient failed to recognize his mother and became violently delirious. During the next three days there was no change in his condition, and he was brought to the Carney Hospital. On admission, Oct. 6, 1910, his temperature was 101.5, his pulse 110, his respirations 36 to the minute.

Physical examination showed an emaciated but fairly well-developed young man lying in bed with his head slightly retracted and turned towards the right side, his legs acutely flexed on his thighs and trunk, and breathing rapidly though not laboredly through a widely opened mouth. There are some fibrillary twitchings of the muscles of the arms and legs and the arms are constantly being moved about in a semi-purposeful manner. The patient seems to vaguely notice his surroundings, but he makes no response when addressed.

Eyes. — Both eyes are kept constantly rolled up under the lids with a slight tendency to direct the pupils upward and to the left. There is no evident strabismus. The pupils are moderately dilated, both are irregular, the left possibly is slightly larger than the right. No definite reaction to light is obtained, although both pupils are constantly altering their size.

Ears reveal no discharge, and there is no mastoid tenderness.

¹ Diagnosis as stated by the patient's mother.

Lips, teeth and tongue are covered with dry sordes. Tongue lies in the middle of the mouth, is protruded in the mid-line.

Throat is negative.

Neck is held rigidly and slightly retracted. Attempts to bring the head forward appear to elicit pain.

Glands, cervical, axillary, right epitrochlear and inguinal, are palpable, not greatly enlarged.

Thorax is long, narrow and symmetrical. Expansion is good and equal on the two sides.

Lungs.—Everywhere in front there is resonance, and no râles are heard. In the left back below the level of the spine of the scapula there is dullness, and a few scattered fine moist râles are heard. In the right back below the angle of the scapula the percussion note is flat, there is a decreased transmission of breath sounds and many fine and consonating râles are heard.

Heart.—The apex beat is palpable in the fifth interspace $3\frac{1}{2}$ inches from the mid-sternal line. On percussion the left border of the heart corresponds with this position of the apex beat, while the right border is one inch to the right of the mid-sternal line, and the upper border is at the lower margin of the third rib. The heart action is regular, the sounds are of a weak and poor quality, unaccompanied by murmurs.

Abdomen is moderately distended, tympanitic throughout except in the right flank. There is no shifting dullness, no palpable masses.

Liver dullness extends from the sixth rib to the costal margin, edge not palpable.

Spleen not enlarged to percussion, edge not palpable.

Genitalia are small, no palpable masses or areas of tenderness.

Extremities.—Marked atrophy of the calves and the thighs. Muscles of the arms and legs are spastic, but the spasticity is easily overcome. Positive Kernig sign is elicited on both sides.

Reflexes.—Arm reflexes cannot be elicited. Right knee-jerk is present, the left absent. Both Achilles reflexes are present, but the left one is sluggish. There is a positive Babinski sign on the left side; on the right the plantar reflex is normal. There is no ankle clonus, no cremasteric and no abdominal reflex present.

Skin shows no rash. There is some redness over the bony prominences, but no decubitus.

Blood.—Hemoglobin 85%, white blood cells 22,400. Smear shows a polymorphonuclear leucocytosis, no other abnormality. Widal reaction negative.

Oct. 7. Patient is quite irrational, but will protrude the tongue and close the eyes when requested. The legs are held flexed and are immediately returned to this position after forcible extension. The patient is incontinent. His temperature is 99.6, pulse 120, respiration 60 during the forenoon. Under chloroform anesthesia a lumbar puncture was performed and 40 ccm. of turbid fluid was removed under a moderate increase of pressure. Thirty cubic centimeters of Flexner's anti-meningococci serum was injected. The patient's condition remained unchanged during the day, and the evening temperature was 101.2, pulse 140, respiration 60.

Oct. 8. The patient is gradually failing. He is in a state of stupor. Respirations are Cheyne-Stokes in type, temperature 102, pulse 134, respirations 60, with periods of apnea lasting for twenty seconds.

Oct. 9. The patient is distinctly worse, temperature is 104, pulse 150, respiration 70, still of Cheyne-Stokes type. The patient became increasingly stuporous and died four hours later.

Examination of the spinal fluid.—Amount, 40 ccm.; turbid, not blood stained. Specimen was centrifuged in a sterilized tube. Smears from this sediment showed a great predominance of polymorphonu-

clear leucocytes, there being 98% of polymorphonuclear and 2% of mononuclear cells. There were also present very many bacilli, acid-fast, and not decolorized by the Gram's method. These bacilli resisted decolorization by 10% nitric acid after carbolfuchsin staining for fifteen minutes.

Two guinea pigs were injected, each with $\frac{1}{2}$ ccm. of the spinal fluid, one subcutaneously, the other intraperitoneally. The guinea pig injected subcutaneously died three days after the injection with a subcutaneous abscess at the site of injection. Smears made from this abscess showed an acid-fast bacillus. Material from this abscess was rubbed up with sterile water and injected subcutaneously into a guinea pig. This guinea pig killed three weeks later showed the same acid-fast bacillus present in caseous areas in the spleen. The guinea pig which was injected intraperitoneally with spinal fluid was killed three weeks later, and at autopsy the peritoneum was found to be studded with small tubercles, and in the liver, spleen and omentum there were small foci of caseous material. Sections from these areas showed under the microscope miliary tubercles with giant cells and caseation. In these sections an acid-fast bacillus was found.

SUMMARY.

A young adult with physical signs of meningitis, fever and leucocytosis shows on lumbar puncture spinal fluid turbid, containing 98% of polymorphonuclear leucocytes and very many bacilli, with the morphological and tinctorial peculiarities of the tubercle bacilli, yielding by animal inoculation the typical lesions of tuberculosis. The case is reported to emphasize the fallacy of diagnosing meningitis as due to pyogenic cocci from obtaining a turbid spinal fluid containing a large excess of polymorphonuclear leucocytes.

THE PUBLIC HEALTH LABORATORY.*

BY B. L. ARMS, M.D.,

Director of the Bacteriological Laboratory of the Boston Board of Health.

THE first public health laboratory was established in 1888 by Dr. Gardner T. Swarts, of Providence, R. I., but since that time laboratories have become such a necessity that they are found in nearly all large cities, and the states with but few exceptions also have laboratories in order that they may do the needed work for the smaller towns and cities which would otherwise be unable to have that service and also to do the larger or state problems,—as water and sewage work, food and drug analysis and medico-legal examinations. Besides the large cities and states, a constantly increasing number of smaller cities and towns have municipal laboratories and keep constant supervision over contagious diseases and the milk supply.

At the outset, the aid of the laboratory was most frequently asked in the diagnosis of diphtheria, but at the present time there are many other ways in which they may aid the physician.

In order to render the best service there should

* Read at a meeting of the New England Pediatric Society, Feb. 25, 1911.