This article was downloaded by: [130.132.123.28]

On: 03 January 2015, At: 16:26

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer
Street, London W1T 3JH, UK

Annals and Magazine of Natural History: Series 5

Publication details, including instructions
for authors and subscription information:

<http://www.tandfonline.com/loi/tnah11>

XI.—Classification of the Dinosauria

Prof. O.C. Marsh

Published online: 07 Oct 2009.

To cite this article: Prof. O.C. Marsh (1882) XI.—Classification of the
Dinosauria , Annals and Magazine of Natural History: Series 5, 9:50, 79-84,
DOI: [10.1080/00222938209458999](https://doi.org/10.1080/00222938209458999)

To link to this article: <http://dx.doi.org/10.1080/00222938209458999>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of
all the information (the “Content”) contained in the publications
on our platform. However, Taylor & Francis, our agents, and our
licensors make no representations or warranties whatsoever as
to the accuracy, completeness, or suitability for any purpose of
the Content. Any opinions and views expressed in this publication
are the opinions and views of the authors, and are not the views
of or endorsed by Taylor & Francis. The accuracy of the Content
should not be relied upon and should be independently verified
with primary sources of information. Taylor and Francis shall not be
liable for any losses, actions, claims, proceedings, demands, costs,
expenses, damages, and other liabilities whatsoever or howsoever

caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

XI.—*Classification of the Dinosauria*.

By Prof. O. C. MARSH*.

IN the May number of the 'American Journal of Science' (p. 423), I presented an outline of a classification of the Jurassic Dinosaurian reptiles of this country which I had personally examined. The series then investigated is deposited in the museum of Yale College, and consists of several hundred individuals, many of them well preserved, and representing numerous genera and species. To ascertain how far the classification proposed would apply to the material gathered from wider fields, I have since examined various Dinosaurian remains from other formations of this country, and likewise, during the past summer, have visited most of the museums of Europe that contain important specimens of this group. Although the investigation is not yet complete, I have thought the result already attained of sufficient interest to present to the Academy at this time.

In previous classifications, which were based upon very limited material compared with what is now available, the Dinosaurs were very generally regarded as an order. Various characters were assigned to the group by Von Meyer, who applied to it the term Pachypoda; by Owen, who subsequently gave the name Dinosauria, now in general use; and also by Huxley, who more recently proposed the name Ornithoscelida, and who first appreciated the great importance of the group, and the close relation it bears to birds. The researches of Leidy and Cope in this country, and of Hulke, Seeley, and others in Europe, have likewise added much to our knowledge of the subject.

An examination of any considerable portion of the Dinosaurian remains now known will make it evident to any one familiar with reptiles, recent or extinct, that this group should be regarded, not as an order, but as a subclass; and this rank is given to it in the present communication. The great number of subordinate divisions in the group, and the remarkable diversity among those already discovered, indicate that many new forms will yet be found. Even among those now known there is a much greater difference in size and in osseous structure than in any other subclass of vertebrates, with the single exception of the placental Mammals. Compared with the Marsupials, living and extinct, the Dinosauria show an equal diversity of structure, and variations in size from by

* Communicated by the Author, having been read before the National Academy of Sciences, at the Philadelphia meeting, November 14, 1881.

far the largest land animals known (50 or 60 feet long) down to some of the smallest, a few inches only in length.

According to present evidence, the Dinosaurs were confined entirely to the Mesozoic age. They were abundant in the Triassic, culminated in the Jurassic, and continued in diminished numbers to the end of the Cretaceous period, when they became extinct. The great variety of forms that flourished in the Triassic render it more than probable that some members of the group existed in the Permian period; and their remains may at any time be brought to light.

The Triassic Dinosaurs, although so very numerous, are known today mainly from footprints and fragmentary osseous remains. Not more than half a dozen skeletons at all complete have been secured from deposits of this period; hence many of the remains described cannot at present be referred to their appropriate divisions in the group.

From the Jurassic period, however, during which Dinosaurian reptiles reached their zenith in size and numbers, representatives of no less than four well-marked orders are now so well known that different families and genera can be very accurately determined, and almost the entire osseous structure of typical examples, at least, be made out with certainty. The main difficulty at present with the Jurassic Dinosaurs is in ascertaining the affinities of the diminutive forms which appear to approach birds so closely. These forms were not rare; but their remains hitherto found are mostly fragmentary, and can with difficulty be distinguished from those of birds, which occur in the same beds. Future discoveries will, without doubt, throw much light upon this point.

Comparatively little is yet known of Cretaceous Dinosaurs, although many have been described from incomplete specimens. All of these appear to have been of large size, but much inferior in this respect to the gigantic forms of the previous period. The remains best preserved show that, before extinction, some members of the group became quite highly specialized.

Regarding the Dinosaurs as a subclass of the REPTILIA, the forms best known at present may be classified as follows:—

Subclass DINOSAURIA.

Premaxillary bones separate; upper and lower temporal arches; rami of lower jaw united in front by cartilage only; no teeth on palate. Neural arches of vertebræ united to centra by suture; cervical vertebræ numerous; sacral vertebræ co-

ossified. Cervical ribs united to vertebræ by suture or ankylosis; thoracic ribs double-headed. Pelvic bones separated from each other and from sacrum; ilium prolonged in front of acetabulum; acetabulum formed in part by pubis; ischia meet distally on median line. Fore and hind limbs present, the latter ambulatory and larger than those in front; head of femur at right angles to condyles; tibia with procnemial crest; fibula complete. First row of tarsals composed of astragalus and calcaneum only, which together form the upper portion of ankle-joint.

(1.) Order SAUROPODA (Lizard-foot). Herbivorous.

Feet plantigrade, ungulate; five digits in manus and pes; second row of carpals and tarsals unossified. Pubes projecting in front, and united distally by cartilage; no postpubis. Precaudal vertebræ hollow. Fore and hind limbs nearly equal; limb-bones solid. Sternal bones parial. Premaxillaries with teeth.

(1.) Family *Atlantosauridæ*. Anterior vertebræ opisthocœlian. Ischia directed downward, with extremities meeting on median line.

Genera: *Atlantosaurus*, *Apatosaurus*, *Brontosaurus*, *Diplodocus*, ? *Camarasaurus* (*Amphicælias*), ? *Dystrophæus*.

(2.) Family *Morosauridæ*. Anterior vertebræ opisthocœlian. Ischia directed backward, with sides meeting on median line.

Genus *Morosaurus*.

European forms of this order: *Bothriospondylus*, *Ceteosaurus*, *Chondrosteosaurus*, *Eucamerotus*, *Ornithopsis*, *Pelorosaurus*.

(2.) Order STEGOSAURIA (Plated lizard). Herbivorous.

Feet plantigrade, ungulate; five digits in manus and pes; second row of carpals unossified. Pubes projecting free in front; postpubis present. Fore limbs very small; locomotion mainly on hind limbs. Vertebræ and limb-bones solid. Osseous dermal armour.

(1.) Family *Stegosauridæ*. Vertebræ biconcave. Neural canal in sacrum expanded into large chamber; ischia directed backward, with sides meeting on median line. Astragalus coossified with tibia; metapodials very short.

Genera: *Stegosaurus* (*Hypsirhophus*), *Diracodon*, and in Europe *Omosaurus*, Owen.

(2.) Family *Scelidosauridæ*. Astragalus not coossified with tibia; metatarsals elongated; four functional digits in pes. Known forms all European.

Genera: *Scelidosaurus*, *Acanthopholis*, *Cratæomus*, *Hylæosaurus*, *Polacanthus*.

(3.) Order ORNITHOPODA (Bird-foot). Herbivorous.

Feet digitigrade, five functional digits in manus, and three in pes. Pubes projecting free in front; postpubis present. Vertebrae solid. Fore limbs small; limb-bones hollow. Premaxillaries edentulous in front.

(1.) Family *Camptonotidae*. Clavicles wanting; postpubis complete.

Genera: *Camptonotus*, *Laosaurus*, *Nanosaurus*, and in Europe *Hypsilophodon*.

(2.) Family *Iguanodontidae*. Clavicles present; postpubis incomplete. Premaxillaries edentulous. Known forms all European.

Genera: *Iguanodon*, *Vectisaurus*.

(3.) Family *Hadrosauridae*. Teeth in several rows, forming with use a tessellated grinding surface. Anterior vertebrae opisthocelcian.

Genera: *Hadrosaurus*, ? *Agathaumas*, *Cionodon*.

(4.) Order THEROPODA (Beast-foot). Carnivorous.

Feet digitigrade; digits with prehensile claws. Pubes projecting downward, and coossified distally. Vertebrae more or less cavernous. Fore limbs very small; limb-bones hollow. Premaxillaries with teeth.

(1.) Family *Megalosauridae*. Vertebrae biconcave. Pubes slender, and united distally. Astragalus with ascending process. Five digits in manus, and four in pes.

Genera: *Megalosaurus* (*Poikilopleuron*), from Europe; *Allosaurus*, *Cælosaurus*, *Creosaurus*, *Dryptosaurus* (*Laelaps*).

(2.) Family *Zanclodontidae*. Vertebrae biconcave. Pubes broad elongate plates, with anterior margins united. Astragalus without ascending process; five digits in manus and pes. Known forms European.

Genera: *Zanclodon*, ? *Teratosaurus*.

(3.) Family *Amphisauridae*. Vertebrae biconcave. Pubes rodlike; five digits in manus, and three in pes.

Genera: *Amphisaurus* (*Megadactylus*), ? *Bathygnathus*, ? *Clepsysaurus*, and in Europe *Palæosaurus*, *Thecodontosaurus*.

(4.) Family *Labrosauridae*. Anterior vertebrae strongly opisthocelcian and cavernous. Metatarsals much elongated. Pubes slender, with anterior margins united.

Genus *Labrosaurus*.

Suborder CÆLURIA (Hollow-tail).

(5.) Family *Cœluridæ*. Bones of skeleton pneumatic or hollow. Anterior cervical vertebræ opisthocœlian, remainder biconcave. Metatarsals very long and slender.

Genus *Cœlurus*.

Suborder COMPSOGNATHA.

(6.) Family *Compsognathidæ*. Anterior vertebræ opisthocœlian. Three functional digits in manus and pes. Ischia with long symphysis on median line. Only known specimen European.

Genus *Compsognathus*.

DINOSAURIA ?

(5.) Order HALLOPODA (Leaping-foot). Carnivorous ?

Feet digitigrade, unguiculate ; three digits in pes ; metatarsals greatly elongated ; calcaneum much produced backward. Fore limbs very small. Vertebræ and limb-bones hollow. Vertebræ biconcave.

Family *Hallopodidæ*.

Genus *Hallopus*.

The five orders defined above, which I had previously established for the reception of the American Jurassic Dinosaurs, appear to be all natural groups, well marked in general from each other. The European Dinosaurs from deposits of corresponding age fall readily into the same divisions, and, in some cases, admirably supplement the series indicated by the American forms. The more important remains from other formations in this country and in Europe, so far as their characters have been made out, may likewise be referred with tolerable certainty to the same orders.

The three orders of herbivorous Dinosaurs, although widely different in their typical forms, show, as might be expected, indications of approximation in some of their aberrant genera. The Sauropoda, for example, with *Atlantosaurus* and *Brontosaurus* of gigantic size for their most characteristic members, have in *Morosaurus* a branch leading toward the Stegosauria. The latter order, likewise, although its type genus is in many respects the most strongly marked division of the Dinosaurs, has in *Scelidosaurus* a form with some features pointing strongly towards the Ornithopoda.

The Carnivorous Dinosauria now best known may all be placed at present in a single order ; and this is widely sepa-

rated from those that include the herbivorous forms. The two suborders defined include very aberrant forms, which show many points of resemblance to Mesozoic birds. Among the more fragmentary remains belonging to this order, but not included in the present classification, this resemblance appears to be carried much further.

The order Hallopoda, which I have here referred to the Dinosauria, with doubt, differs from all the known members of that group in having the hind feet especially adapted for leaping, the metatarsals being half as long as the tibia, and the calcaneum produced far backward. This difference in the tarsus, however, is not greater than may be found in a single order of Mammals, and is no more than might be expected in a subclass of Reptiles.

Among the families included in the present classification, I have retained three named by Huxley (*Scelidosauridæ*, *Iguanodontidæ*, and *Megalosauridæ**), although their limits as here defined are somewhat different from those first given. The suborder Compsognatha also was established by that author in the same memoir, which contains all the more important facts then known in regard to the Dinosauria. With the exception of the *Hadrosauridæ*, named by Cope, the other families above described were established by the writer.

The *Amphisauridæ* and the *Zanclodontidæ*, the most generalized families of the Dinosauria, are only known from the Trias. The genus *Dystrophæus*, referred provisionally to the Sauropoda, is likewise from deposits of that age. The typical genera, however, of all the orders and suborders are Jurassic forms; and on these especially the present classification is based. The *Hadrosauridæ* are the only family confined to the Cretaceous. Above this formation there appears to be at present no satisfactory evidence of the existence of any Dinosauria.

XII.—*On a small Collection of Lepidoptera from Melbourne.*
By ARTHUR G. BUTLER, F.L.S., F.Z.S., &c.

THE present series consists of ninety-one examples, some of them unfortunately in very poor condition, forwarded to us from Australia by Dr. T. P. Lucas. Notwithstanding that not a few of the specimens are more or less worn or broken,

* 'Quarterly Journal Geological Society of London,' vol. xxvi. p. 34, 1870.