New York had examined her and pronounced her deafness incurable. This result suggests that the rays may be of value in some forms of deafness.

In conclusion, I am convinced that:

1. The *x*-rays are of undoubted value in the treatment of certain cases of both superficial and deep-seated carcinoma and tuberculosis.

2. The more a case has been tampered with, the less likely it is to yield to the influence of the *x*-rays, *e. g.*, Case 6.

3. Daily treatments carefully and properly given will produce the best results, *e. g.*, Cases 1 and 4.

4. We should try never to produce a dermatitis beyond a simple erythema.

5. There are idiosyncrasies in certain people which render them most susceptible to the *x*-rays, and in these people deeper burns may occur in spite of the most careful treatment, *e. g.*, Case 3.

6. Epithelioma involving the mucous membrane is much less likely to yield to the effect of the *x*-rays than when it simply involves the skin, *e. g.*, Case 6.

7. There is not likely to be any interference with the sense of sight, even though the *x*-rays are used directly over the eye, *e. g.*, Case 4.

8. Tuberculosis, whether of the skin or of the glands, will yield at least in certain cases to the effects of the *x*-rays, *e. g.*, Cases 7, 8 and 9.

9. The *x*-rays may be of value in certain cases of deafness.

10. The *x*-rays will give better cosmetic results than any other form of treatment in simple epithelioma of the face, *e. g.*, Case 5.

11. Epithelioma of the mucous membrane should be removed as early as possible by the knife, and this followed by *x*-ray treatment.

12. Operable cases should be operated on and in each instance followed by a course of *x*-ray treatment, and *x*-ray treatment should be given at the first sign of a recurrence.

13. Inoperable cases should be given a trial with the *x*-rays, since even hopeless cases sometimes yield to this form of treatment, *e. g.*, Case 4.

14. It is desirable to produce a distinct reaction in the treatment of lupus, *e. g.*, Case 9.

Clinical Reports.

TWO CASES OF TRANSITORY COMPLETE BLINDNESS OF BOTH EYES.*

CHARLES J. KIPP, M.D.
NEWARK, N. J.

IDIOPATHIC OPTIC NEURITIS.

CASE 1.—Mrs. K. S., aged 32, has been married eleven years, and is apparently in robust health. She has given birth to two children, both of whom are living and in good health; the younger is 4 years old. Patient has always been in good health. No history of syphilis. For some weeks she has suffered more or less from frontal headache.

She came to the Newark Eye and Ear Infirmary for the first time Dec. 22, 1901, complaining of loss of vision of left eye and impairment of vision of right eye. The examination made at this time showed that the exterior parts of the eye were perfectly normal in appearance, and that the movement of the eyes were unimpeded in any direction. On closer examination it was found that the left eye was blind; this eye could not distinguish between light and darkness. The pupil of this eye was larger than that of the right eye and it did not react

to light. It did, however, respond to light thrown in the right eye. The ophthalmoscope showed clear media and a well-marked papillitis. The optic disc was swollen and apparently enlarged. The retinal veins were full and tortuous, while the arteries were of about normal caliber. The retina immediately surrounding the disc was hazy, but otherwise it was normal. The right eye also showed swelling and enlargement of the optic disc, but not quite so marked as that of the other eye. The retina was transparent except very close to the optic disc. The pupil of the right eye was of normal size and responded readily to light. The sight of this eye was reduced to 5/12. An examination of the urine showed it to be free from albumin and sugar. Close questioning failed to reveal disease in any other part of the body.

She was given iodid of potassium in small doses. She was seen again the next day, when it was found that both eyes were now totally blind. She could not recognize the light of a candle held close to her eyes. The swelling of the optic discs had increased. Their surfaces were best seen with a +3 dioptric glass. The pupil of the right eye was now as large as that of the left eye, and neither pupil reacted to light.

She remained in this condition until December 28, when she could distinguish between light and darkness. December 30 it was found that the pupil of the right eye responded very sluggishly to light, but contracted in convergence. The pupil of the left eye responded even more sluggishly than the right eye to light, but also contracted in convergence. She can see movements of hands in front of her eyes, but can not yet count fingers close to her eyes. She was at this time taking fifteen grains of the iodid of potassium three times daily.

Jan. 8, 1892, it was found that the pupils of both eyes reacted to light in convergence fairly well. The vision of both eyes remained about the same as at last, January 14. Can now count fingers with either eye at four or five feet. She still has frontal headache. The optic discs are less swollen and the retina adjoining the disc is more transparent.

January 22: Her vision is now 5/18 in the right eye and 5/36 in the left. The visual field is intact in both eyes. She is taking twenty grains of the iodid three times daily. After this time her vision gradually improved so that on February 5 it was 5/9 in either eye. The discs were now almost flat and quite white. The headache had subsided.

March 15 her vision was about normal in both eyes and the visual field intact in both. During the following five years I saw the patient at intervals of several months and always found her in good health. Her vision when I last examined her five years after the attack was normal in both eyes and so were the visual fields. Her health was excellent. I have been trying to find her since, but have not succeeded in so doing.

The absence of the pupillary light reflex for a while made it probable that the seat of the lesion was in the visual tract, this side of the corpora quadrigemina.

The visual centers may also have been involved, but this can not be proved, as the conduction of visual impressions was interrupted by disease of the tract. The ophthalmoscope showed the existence of a marked optic neuritis in both eyes. Cases in which an optic neuritis has caused total blindness of both eyes and in which the pupillary light reflex was absent for a time, with subsequent restoration of normal vision, and very seldom encountered. With the exception of a frontal headache my patient was in excellent good health at the time the optic neuritis developed, and as she remained in good health during the next five years it seems reasonable to assume that the eye disease was not the result of a general affection which was latent when she became blind.

AMAUROSIS, PROBABLY FROM AUTOINTOXICATION.

CASE 2.—Mrs. S., aged 52, stout, of typical bilious temperament, is subject to frequent attacks of headache; is constipated most of the time; is fond of rich food and indulges freely in eating.

July 27, in the evening, she was attacked with a sudden intense pain in the head, with nausea and vomiting. The attack was so severe that her attending physician was obliged to give her ¼ grain of morphin hypodermically. She was re-

* Read at the one hundred and thirty-seventh annual meeting of the Medical Society of New Jersey.

lieved in about half an hour. Calomel and Warburg's tincture were given next morning, but did not act well. In the evening of this day, at about the same hour as on the previous day, another severe attack of headache came on and another dose of morphin was given her, which gave her some relief.

On the morning of the 29th she was given one ounce of Warburg's tincture, and in the evening she received ten grains of quinin. She had no attack of headache on this day.

On the 30th, in the evening, she had another very severe attack of headache. In the morning of this day she had taken ten grains of quinin and ten more in the evening. During the night of the 30th she noticed that her vision was becoming impaired, and on the morning of the 31st she was totally blind in both eyes. Pupils reacted but feebly to light. I saw her for the first time shortly after noon on this day. She acted like a person who was absolutely blind and was quite self-possessed. I found that the movements of the eyeballs were unimpeded in any direction. She was unable to recognize the position of a lighted candle held but a short distance from her eyes. The pupils of both eyes responded readily to light and in convergence. The ophthalmoscope showed absolutely normal eye grounds. In one of her eyes she had a patch of opaque nerve fibers such as is frequently seen in eyes with good vision. There was absolutely nothing hysterical about the woman. I feel very confident that she did not exaggerate her impairment of vision. The hearing was normal. I made a favorable prognosis. An examination of her urine showed it to have a specific gravity of 1030; it contained a trace of sugar, but no albumin. The quantity of urine passed during the day preceding the blindness is not accurately known, but it was said to have been of the usual amount. The bowels had not moved freely for some days, although sulphate of magnesia in large doses had been administered at frequent intervals. A high enema was given shortly after I saw her, which brought away a quantity of fecal matter, in which, on examination, were found seeds of berries eaten two weeks before. The total blindness continued for thirty-six hours, and then gradually her vision returned in both eyes. There was no rise in temperature during the attack, but on the day following her temperature rose to 102. A week after the attack her vision was as good as before the attack, and the visual fields were of normal extent for white and colors in all directions.

All our efforts to find a cause for the sudden blindness was fruitless. She has no disease of the kidneys. Habitual constipation, for which she always was taking cathartic mineral waters, was about the only abnormal condition which we could discover at the time. She has been in her usual good health ever since.

That this was a case of amaurosis from some kind of intoxication there can be no doubt. But as to whether it was due to a poison introduced into the body from without or whether the poisoning was the result of an inadequate physiologic elimination of the product of the tissue metamorphosis or of the formation of a poison in the intestinal tract due to abnormal processes of decomposition or fermentation, must remain in doubt. That the poison was not introduced from without seems, however, pretty certain. Neither the food of which she had partaken nor the drugs which had been administered can be held responsible for the eye affection. We know that quinin in comparatively small doses will occasionally produce blindness, but both the clinical features and the absence of all change in the backgrounds of the eyes, as well as the subsequent history of the case, show that this drug was not the cause of the blindness. The absence of disease of the kidneys probably eliminates uremia as the cause of the amaurosis, though it must be admitted that this attack closely resembles in all its features the blindness known to occur in this affection. In one of the examinations of the urine a trace of sugar was found, but only once, and none has been found since the attack. With the exception of habitual constipation, there is no evidence of the presence of a morbid affection of the digestive tract, and yet the most plausible explanation seems to me to be that the toxin which caused the amaurosis was formed here through putrefactive changes of the contents of the intestinal tract. As to the part of the visual apparatus

affected in this case, there can be no doubt. The blindness with prompt pupillary reaction to light established the fact that the optical centers were the seat of the lesion. But as to whether it was an edema of the meninges or an affection of the brain itself no one can tell. There are very few cases on record similar to the one here described, and in most of these the absence of albumin in the urine was not positively excluded. I am indebted to Dr. Cladek of Rahway for notes of the previous history of the case.

To recapitulate: In both of the cases above reported the blindness was due to an affection of the nervous visual apparatus. In the first the seat was in the conducting apparatus, as was manifest from the fact that the pupillary light reflex was absent during a part of the time that the blindness existed. In the second the seat of the lesion must be looked for in the optic centers, as the pupils reacted promptly to light while total blindness was present. In both cases the vision was completely restored. It is doubtful that the treatment adopted had anything to do with the restoration of the sight in either case.

I. PROLONGED INTUBATION. II. ESOPHAGOTOMY FOR A DENTAL PLATE.

III. LARYNGECTOMY FOR CARCINOMA.*

GEORGE F. COTT, M.D.

CLINICAL PROFESSOR OF OTOTOLOGY, UNIVERSITY OF BUFFALO
BUFFALO, N. Y.

I.—A CASE OF PROLONGED INTUBATION.

One of the rarest accidents following intubation for diphtheria is stenosis, which results from pressure ulceration of the subglottic region and is very difficult to remedy. There are, perhaps, half a dozen cases on record; several made complete recoveries. One of these is as follows:

Intubation.—Rosa McGlynn, aged 6, was taken sick with diphtheria in December, 1901, and in a very few days required intubation. Being otherwise engaged, I sent my assistant, Dr. James E. King, to perform the operation. A week later the tube was removed, but had to be replaced immediately. Several days thereafter I removed the tube, but within a very few minutes had to replace it. This occurred frequently for several weeks. In order to keep better watch of the child I had her removed to the Riverside Hospital, and there the tube was removed and replaced off and on for several months.

Excision of Stricture.—Being satisfied that intubation would do no further good, especially since it had become necessary to use a one-year tube, I split the larynx completely and found a fibrous stricture below the vocal cords in the subglottic tissue. This was cut out and considerable granulation tissue removed with a sharp spoon, and a silver tube passed up from the tracheal wound between the vocal cords and the larynx, and a tracheal tube downward through which the child breathed. These tubes were left in for some time, when intubation was again resorted to, but gradually the lumen was again diminished until it became very difficult to insert a one-year tube. The larynx was opened a second time and the stricture removed as before. The upper or laryngeal tube was not replaced, but the lower or tracheal tube was allowed to remain for several weeks. Intubation was intermittently performed during this time.

Course.—After a while the tracheal tube was removed and the child allowed to go several days without a tube, but occasionally she had to be intubated, on very short notice. The interne was instructed to be on his guard either to intubate or to do tracheotomy on a moment's notice. About 2 o'clock one morning he was suddenly summoned and found the child suffocating. Dr. Weed hurriedly opened the trachea and the danger was tided over once more. After that I opened the larynx for the third time, removed the stricture again and

* Read at the Fifty-fourth Annual Session of the American Medical Association, in the Section on Laryngology and Otology, and approved for publication by the Executive Committee: Drs. George L. Richards, O. Joachim and G. V. Woollen.