plasms and inflammatory exudates. Large doses are often required and tolerated. A case of total ophthalmoplegia with involvement of the upper extremities, respiration and deglutition, necessitating feeding by the stomach tube for eight weeks, was restored to perfect health after taking one ounce of iodides a day for three months. The iodides irritate the larynx and cause distressing dyspnea in some cases, and the potash in combination is a poison to the myocardium, and of course very depressing. Mercury is indicated in all stages of syphilis, but this can not be said of the iodides. As a precautionary measure in the intensive treatment of syphilis with mercury and Salvarsan, the urine should be examined frequently for evidence of kidney irritation. Mercury may cause albuminuria or casts, and the fatalities of Salvarsan have been traced to renal insufficiency.

DISCUSSION

Dr. E. H. Cary, Dallas, Tex.—This is the greatest subject which confronts you men, or rather the greatest which confronts humanity, and at all times the greatest with which the oculist has to deal. I remember during my service in Bellevue Hospital that it was not long before I came to the conclusion that everybody was more or less infected with syphilis. And just as Dr. Osler stated, I believed that no man understood disease until he understood syphilis. I had occasion to watch for the symptoms which might occur from this infection and it was not long before I felt, from a pretty general observation, that humanity was badly involved.

In my service at the New York Eye and Ear Infirmary I found that syphilis was the one great diagnosis and I believe we all treated as syphilis every case that we were not able to make out and differentiate. I don't at this moment wish to say that syphilis has particularly changed so far as involving humanity is concerned, but I do believe we are more willing to let some other causes prevail in our mind and seek for those causes. And I do not think we are willing to go on record now as often as we used to that this or that may be due to syphilis without trying to differentiate symptoms and seek for cause. The Wassermann reaction has been of valuable assistance, but can only be regarded in the light of an aid. You can not place all your faith in it. I have frequently found it necessary to watch results which would make it possible to conclude from treatment that the Wassermann was wrong. This has been true in both positive and negative reports. After all, none of the laboratory findings, known to be of great assistance, can take the place of your ability to put two and two together. That is an individual preference and your skill is based upon not only your individual ideas, but your knowledge and ability to corroborate all these things

when given to you. Of course, those are generalities which are self-evident. I once felt that practically 80% of my iritic cases were syphilitic, but I have changed my opinion that they are in any such proportion. In fact, you can drop it down to 25%, or something of that kind, and you would be more correct as to the cause of iritis. That, of course, has been pretty nearly always treated as syphilis. If we could not come to some other conclusion at once, syphilis was the cause.

There are so many conditions occurring in and out of the eye which have been classified as syphilis that I simply want to call your attention to the fact that there are a great many ways of arriving at different conclusions; but we must always take into consideration that while syphilis is not the whole thing it is a great disease that we have to consider.

Not long ago a very interesting case of a little child came here who had a fistula from the ethmoidal sinus. The orbital contents were well pushed forward, the child being very weak, anemic and toxic. The differential diagnosis had to be made of course as to whether it was syphilitic involvement of the sinuses or sarcoma of the orbit. It was a suppurative process. The child was five years old. The Wassermann was positive, but in going back in the histories of the families they were as pure as you have ever seen. Both parents were negative with the Wassermann. Again going back into its history, I found that the child had been nursed by an old negro "mammy" who had lost an eye from syphilis; and as one of the doctors from east Texas said to me, "Yes, I know her. She still has syphilis and she is still nursing children for different families." Although it looked as if the child might die, accurate treatment was given and it recovered sufficiently for us to remove later the necrotic bone, which hastened cure.

Syphilis involves so many different parts in which we are interested that it can not be lost sight of. The doctor said he did not think the ear was much involved. However, I have been struck in the last few years with a certain type of cases that were losing their hearing, and I found that some of them would respond actively to syphilitic treatment. I am sure there are a greater number of cases in which ear trouble is due to syphilis than we are usually inclined to think because we are more likely to connect syphilis with the eye than the ear. Many of you have seen syphilitic mastoids, but I am sure to discuss this question properly now, a man would tire you to death.

THE INJECTION OF THE GANGLION OF GASSER THROUGH THE FORAMEN OVALE.*

By H. H. MARTIN, M.D., F.A.C.S.,
Savannah, Ga.

The injection of the ganglion of Gasser

*Read in Eye, Ear, Nose and Throat Section, Southern Medical Association, Ninth Annual Meeting, Dallas, Tex., Nov. 8-11, 1915.

through the foramen ovale for the relief of trigeminal neuralgia and for conduction anesthesia has passed beyond the experimental stage and will eventually supersede surgical extirpation of the ganglion, division of the sensory root, peripheral and nerve trunk injection, and other more or less unsatisfactory surgical procedures. The operation is, of course, a very delicate one and requires in addition to a thorough knowledge of the anatomy of the parts, some surgical skill and dexterity; the needle will necessarily pass very near to important blood vessels, principally the internal maxillary artery, which lies in close proximity to the inferior maxillary nerve in the sphenomaxillary fossa and sends a small meningeal branch through the foramen ovale.

If improperly directed, the needle may enter the foramen spinosum, the jugular foramen, or an anomalous foramen transmitting an emissary vein known as the foramen Civinini, occasionally found just below the foramen ovale. If the needle be passed too far through the foramen ovale, the cavernous sinus or even the internal carotid, both of which are in intimate relation with the ganglion, might possibly be injured. But all such accidents can be safely guarded against; and the procedure, while a formidable one, is not so formidable as the surgical extirpation of the ganglion, while the desirable results are equally good and the undesirable results, such as paralysis of other cranial nerves, weakness of the muscles of mastication, and neuroparalytic keratitis, are never so bad.

The history of this operation, like many others, is one of evolution and dates back to 1903, when Schlosser, an ophthalmologist of Munich, discovered by chance that subcutaneous injections of alcohol in or about an offending nerve afforded prompt relief from pain.

This idea has been taken up and elaborated upon by many investigators in France, England, Germany and America, including ophthalmologists, rhinologists, neurologists, general surgeons and others, until there has been evolved a very simple method of injecting the ganglion through the foramen ovale, which can be acquired by any one who will give sufficient time to experiments on the cadaver.

In a paper presented at the Richmond meeting of the Southern Medical Association, the essayist gave detailed description of the technic of this operation and made preliminary reports on the injection in three living subjects. Since that time three additional cases of genuine, old and persistent "tic" have been treated by the speaker by this method, and I desire in this paper to give more detailed reports of these six cases with some personal observations and comments. I hope also to be able to demonstrate the operation at some of the clinics arranged for this meeting.

My experimental work in conduction anesthesia or nerve blocking was interrupted by an accident which has temporarily disabled me, but I have to report two intra-nasal operations and one tonsillectomy done under this method of conduction anesthesia.

My first injection in the living subject was a woman, twenty-eight years old, complaining of pain in right side of head and face. Careful examination revealed nothing more serious than alveolar abscess on that side. I explained to her that I could open this abscess without pain if she would let me do a preliminary injection. She consented. The injection was done under light ether anesthesia October 19, 1914, as follows: The needle entered the foramen while patient was in primary anesthesia; ten minims of 2% novocaine solution was injected into the ganglion; she was allowed to regain consciousness with the needle in situ, and she was then found to have complete analgesia of the entire area supplied by the fifth nerve. One cc. of 95% alcohol was then injected into the ganglion without the patient's complaining of the slightest sensation. The alveolar abscess was opened the next day without pain or the use of any anesthetic. The patient was kept under observation for five days in order to protect the eye. When she was discharged October 28th following she still had analgesia over the entire area and partial anesthesia as well. This patient was seen six weeks after the injection. At that time there was analgesia over the entire area supplied by the fifth nerve, with total anesthesia of the cornea, the skin over the forehead, and half the tongue.

My second case, male, age thirty-five, was admitted in the acute eruptive stage of syphilis, with violent irido-cyclitis in both eyes, the left eye showing a large bulging mass arising from the ciliary body above, and showing evidence of suppuration. The patient complained of severe pain all over left side of head and face. He consented to the injection preliminary to opening the abscess. It was done November 4, 1914, as follows: Skin and subcutaneous tissues were injected with novocaine; after ten minutes the needle was introduced until the inferior maxil-

lary nerve was reached (evidenced by expression of pain); novocaine was injected at this point, and after ten minutes needle was introduced through foramen ovale (patient evidencing great pain); novocaine injected at this point; and, as there was not complete analgesia after ten minutes, the injection was repeated. After ten minutes more the conclusion was reached that there was something wrong with the novocaine solution. One cc. of alcohol was then slowly injected into the ganglion, patient evidencing great pain during the injection. Ten minutes after the injection of alcohol patient was free from all pain; cornea on injected side could be lightly touched without his knowledge, and there was analgesia over the entire distribution of the nerve, but anesthesia did not extend to the eye.

Twenty-four hours after injection the same condition obtained, although he would flinch from slightest pressure over the ciliary abscess. Forty-eight hours after injection the patient was still free from all pain; there was both analgesia and partial anesthesia of the area supplied by the two lower branches of the nerve; but the eye was still sensitive to pressure. The ciliary abscess, however, was freely opened at this time without anesthetic and with little pain. I believe in this case the injection was not a perfect one, and when the eye was enucleated, some two weeks after the injection, ether was administered. When he left the hospital he had a very incomplete analgesia and no anesthesia.

My third case was one of genuine unilateral neuralgia, left side, male twenty-three years old, had existed for one and one-half to two years. The operation was classical; a few minims of novocaine solution in the skin and subcutaneous tissues produced perfect local anesthesia, and the needle passed through the cheek and sphenomaxillary fossa without pain. When the inferior maxillary nerve was reached sharp pain was complained of, which was almost immediately relieved by a few minims of novocaine. The needle was then advanced one c. m., and when pain was complained of novocaine was again injected; when the needle was advanced another half c. m. and novocaine injected, the patient at once remarked that he had a funny feeling in the eye, face and tongue. One c. c. alcohol was then injected without pain and patient was found to have complete anesthesia of cornea, skin and mucous membranes supplied by the fifth nerve. The patient developed a neuro-paralytic ulcer in the lower temporal quadrant of the cornea, which persisted for weeks. He would not remain in the hospital, but was seen once a week for two months. When discharged the corneal ulcer had healed with only a slight opacity, as the ulceration had been very superficial, apparently no deeper than the corneal epithelium and gave him no trouble at all. When last seen, three months after injection, there was practically total anesthesia of the entire area supplied by the fifth nerve, and no return of tic.

My fourth case came into Dr. Horsley's service at the Memorial Hospital in Richmond on the day following adjournment of the Southern Medical Association.

She had been a victim of trigeminal neuralgia for years, had had numerous peripheral opera-

tions, including resections and nerve trunk injections. She complained that all of her injections had been exceedingly painful, and of little benefit. She was suffering intensely when I saw her first and would cry out if even a motion were made to touch her face. The entire distribution of the nerve was apparently involved, but the mandibular division seemed to be the most affected, this area being exquisitely sensitive. She would cry out if the floor were jarred or the bed brushed against, and in short presented a typical trigeminal neuralgia. Incidentally she had become a morphine habitue. The injection was done under ether anesthesia, and the patient was seen the next morning sitting up eating breakfast, free from pain, and has remained so ever since, so far as I am informed. Furthermore, she was freed from the morphine habit and returned to her home a very happy woman.

My fifth case, a woman thirty years of age, of a highly neurasthenic type, had suffered from tic on left side for years, with occasional attacks on the right side. Nasal examination revealed constant contact of both middle turbinates with the septum. This was corrected without relief. She then visited Sluder and had an alcohol injection in the sphenopalatine ganglion on the left side without relief. Previous to this she had division of the supra-orbital and of the mandibular through their respective foramina. In the latter part of November, 1914, the left ganglion was injected with one cc. 95% alcohol with immediate relief from pain, lasting four months. She had total anesthesia of the entire area with abducent paralysis on left side, which persisted through the second injection of two cc. alcohol four months after the first one. The relief from the second injection lasted about the same length of time; and then severe pain recurred in the mandibular branch and she is still suffering from that condition.

My sixth case was a woman, about sixty, who was brought into the hospital suffering from *tic couloureaux* on both sides, with spasmodic tic involving the muscles of mastication on the right side, the pain on this side being very great. Her history was that of long-standing trigeminal disorder on both sides. The ganglion on the left side had been extirpated some years before by Cushing with considerable relief, but not a complete cure. On the right side numerous peripheral operations had been done without relief. The right ganglion was injected with two c.c. 95% alcohol under ether anesthesia in the latter part of May of this year, with prompt and, so far, permanent relief from both spasmodic and painful tic, and no complications. She stated that Cushing for some reason had declined to operate on the right side. This injection was done while I myself was confined to the hospital. I introduced the needle with the left hand and Dr. Barrow, of Savannah, completed the injection for me.

As stated above, my experience with nerve blocking by this method is limited to three cases. In these cases one-half cc. of 5% novocaine was injected into the ganglion. The anesthesia was instan-

taneous and complete in all branches. In the first case a middle turbinate was removed without the slightest evidence of pain. In the second case ethmoidectomy was done by Reaves' method. And in the third case an old tonsil stump was dissected out—all without pain.

In closing, I beg to apologize for the incompleteness of this case report and for quoting abstracts from a former paper. I have, as you know, been seriously handicapped for some months and could not devote so much time and work to the preparation of this paper as I should like to have done.

A CASE OF DERMOID OF THE CORNEO-SCLERAL MARGIN.*

By ADOLPH O. PFINGST, M.D.,

Professor of Ophthalmology and Otology
University of Louisville, Medical
Department,
Louisville, Ky.

Among the benign extra-ocular tumors of the eye we find the different kinds of dermoid, a not altogether uncommon class of growths.

To illustrate, I have had appear before you this case of dermoid of the corneo-scleral margin of the left eye.

This girl is, as you see, well developed and apparently in splendid health. She is 24 years of age and gives a history of never having had an illness of any kind. She has a birth mark on her left eye. The growth which was noticed soon after her birth has grown somewhat in breadth and height, and recently has developed hairs on the surface. It has never caused her trouble in the way of pain and has never caused the eye to become inflamed.

You will notice at the outer margin of the cornea of the left eye, low down, an irregular hemispherical prominence of firm consistency about 3-16 of an inch in diameter. It is pink or faintly yellowish-pink and resembles a half pearl. You will notice that although it springs from the conjunctiva it slightly overrides the cornea.

The surface is smooth and contains several good sized hairs. Vision in both eyes is normal.

The appearance of this growth, with the history that it is congenital, makes the diagnosis of dermoid positive.

Dermoids are the most common of the teratoid tumors—cysts being far more common than dermoid tumors in the strict sense of the term. They are made up usually of fatty and white fibrous tissue and occasionally contain portions of hyaline cartilage. They have an investing membrane of epithelium. Their structure is characterized usually by the presence of all structures which enter into the make-up of normal skin, viz., hairs and hair follicles, sebaceous and sudoriferous glands, nonstriated fibers and adipose tissue.

The name dermoid was given these growths by Ryba in 1853, when he described the first case of the kind on record. They are congenital defects, and by V. Duyse are supposed to be remnants of amniotic membrane implanted upon the eye during the first two or three months of intrauterine life, while the lids are yet apart.

Remah and others advanced the theory that they are due to fetal invagination of the external blastoderm. At all events, it is evident that embryonic skin tissue must remain on the surface of the eye ball in order for the dermoid to develop.

Von Hippel states that 65% of these cases are associated with other congenital malformations, principally with coloboma of the lids, iris or choroid, but also with cleft palate and harelip.

In cases of congenital defect of the lids, the tumors could develop even after closure of the lips, or, in other words, after the second or third month.

The most frequent seat of dermoid tumors of the eye is at the outer and inferior margin of the cornea. They usually spring from the conjunctiva, but encroach upon the superficial corneal and scleral tissue. Exceptionally they spring from the caruncle and extend from there to the cornea. They are also found in the transition folds of the conjunctiva, most frequently on the eye ball between the superior and external rectus muscle. The growths in this locality contain more fat

*Reported before the Medico-Chirurgical Society of Louisville, Ky., Feb. 23, 1916.