removing the window and allowing the bladder to empty. Turn on the water, replace the window and resume the examination. This may be repeated many times in an ulcerated, contracted bladder, or in one which is freely bleeding, and makes possible a satisfactory examination of any bladder.

Three years' use of the above technic, with air inflating cystoscopes, justifies its recommendation as a safe, simple and practical method.

BONE CYSTS.

A CONSIDERATION OF THE BENIGN AND ADAMANTINE DENTIGEROUS CYSTS OF THE JAW AND BENIGN CYSTS OF THE LONG PIPE BONES.*

JOSEPH C. BLOODGOOD.

BALTIMORE.

Recent accumulated experience, in the ultimate results after amputation for sarcoma of the long pipe bones, and complete resection for sarcoma of the upper and lower jaw, has demonstrated that local recurrence is unusual, but death from internal metastasis is common in a certain number of cases. When these cases are studied pathologically it is found that the patients who have remained well suffered from special types of sarcoma essentially different from the tumors removed from the patients who ultimately succumbed to internal metastasis. In other words, we were not accomplishing a cure in the latter group of cases even after amputation at the highest joint, because internal metastasis takes place early and is present at the time the patient seeks surgical treatment, and we were subjecting the first group of patients to an unnecessarily extensive operation.

In 1899¹ I discussed the literature and the experience of the surgical clinic at the Johns Hopkins Hospital in regard to the different relative malignancy in sarcoma of bone, and that in certain varieties much less extensive operations would accomplish a cure with as great a certainty as amputation at the highest joint.

Experience has demonstrated that, in some cases, curetting is sufficient; for example, the benign bone cysts of the long pipe bones, dentigerous cysts of the jaw and medullary giant-cell sarcoma. The latter was advocated many years ago by Koenig. In other cases resection, the extent of which is indicated by the local infiltration of the disease. For example, the various forms of epulides of the upper and lower jaw; the periosteal and medullary giant-cell sarcoma; the periosteal fibrosarcoma and osteosarcoma; the myxochondrosarcoma and a special tumor of the jaw—the cystic adamantinoma epithelioma. Such local resections as against amputations have been advocated by von Mikulicz, Weisinger, Morton, Karewski, Hinds.²

Amputation is indicated in these varieties of sarcoma only when the necessary resection would result in a useless limb. Infiltration of muscle is not a positive indication for amputation. In this group of cases amputation at the highest joint, except due to the position of the tumor, is never indicated.

In April, 1901, in a discussion before the Philadelphia Academy of Surgery,³ I advocated this more con-

servative procedure. Since then the further experience of Dr. Halsted's clinic and my own and the reading of the literature have accumulated additional facts justifying the more conservative operation in this group of sarcoma of bone, of relatively low malignancy.⁴

In December, 1902, I removed with the curette a large medullary giant-cell sarcoma filling and expanding the upper third of the tibia.⁵ This patient has no evidence of recurrence and a limb with unimpaired function.

The experience of the surgical clinic in tumors of the jaw and long pipe bones can be expressed in a few words.

In Group 1 the patients have remained well since operation, the time varying from six months to twelve years. There has been a slight operative mortality, and a few cases in which, on account of the size and position of the tumor, the disease was considered inoperable.

Benign Cysts of Long Pipe Bones.—Three cases; one died after operation (Fig. 2); 2 cases are well.

Benign Dentigerous Cysts.—Ten cases (upper jaw, 4; lower jaw, 4; ethmoid, 2 cases). Inoperable, no cases. Death after operation, 2 cases (Figs. 5 and 6). In these two instances a complete resection was performed. Well, 8 cases. In these 8 cases the operation consisted in partial removal of the bony wall, curetting and packing.

Adamantine Epithelioma.—Twelve cases; inoperable, one case (Fig. 8). Death after operation, 1 case (Fig. 9). Well, 10 cases. In one instance there was a second operation for local recurrence; this patient has remained well eight years since the second operation.

PERIOSTEAL SARCOMA.

Epulis.—Twenty-three cases; upper jaw, 13; lower jaw, 10. One patient died of pneumonia; in this case, an extensive giant-cell tumor of the lower jaw, it was necessary to do tracheotomy and perform a complete resection of the jaw. Well, 22 cases. In all of these cases the operation consisted of removal of the tumor with the alveolar border of the jaw. In only one case was there a local recurrence, and this patient has remained well since the second operation.

Spindle-cell Fibrosarcoma or Myxosarcoma.—Eight cases; antrum, 3, all well; lower jaw, 3, 2 well, 1 died of pneumonia after complete resection; orbit and antrum, 2 cases; 1, a young child, died after an extensive operation; the other, also a child, has remained well since the less extensive operation, two years.

Osteosarcoma.—(I employ the term osteosarcoma only in those periosteal tumors in which new bone formation predominates.) Eight cases. Lower jaw, 4; 2 well, 2 refused operation. Upper jaw, 2, both well. Long pipe bones (humerus and fibula), 2; both well.

Giant-cell Sarcoma.—Those situated on the jaw and called epulis have already been considered. It is a rare tumor of the long pipe bones. We have observed three cases: Ulna, 1 case, resection, well twelve years. Tibia, upper third, 2 cases, both well; in both the tumor was recurrent; in one the tumor was excised without destroying the continuity of the tibia, in the other the limb was amputated at the thigh.

MEDULLARY SARCOMA.

Giant-cell Tumors (Myeloma).—Eight cases, all well; lower jaw, 1; long pipe bones, 7.

Myxochondrosarcoma.—Three cases; one involving the sacrum, inoperable; one involving the upper third of the humerus, well; one of the femur, death two years after operation from tuberculosis of the lungs.

Seventy-eight bone tumors in this group are either benign or of relatively low malignancy. Sixty-nine cases are well, two inoperable; two refused operation; seven died after operation. In six of these cases the

* Read at the Fifty-fifth Annual Session of the American Medical Association, in the Section on Pathology and Physiology, and approved for publication by the Executive Committee: Drs. V. C. Vaughan, Frank B. Wynn and Joseph McFarland.

1. *Progressive Medicine*, December, 1899, p. 234.

2. *Progressive Medicine*, December, 1899, pp. 38-42.

3. *Annals of Surgery*, 1901, vol. xxxiv, p. 94.

4. *Progressive Medicine*, 1902, pp. 151-186.

5. *Johns Hopkins Hospital Bulletin*, May, 1903.

tumor involved the upper or lower jaw. In these six cases, I believe, a less extensive operation could have been performed which would have reduced the dangers of the operation, but not the probabilities of an ultimate cure.

Group 2 includes patients who have not been cured, either because the condition was inoperable when they presented themselves at the clinic, or because of death from internal metastasis after operation.

PERIOSTEAL TUMORS.

Spindle and Round-cell Sarcoma.—Six cases; lower jaw, 2; long bones, 4.

Perithelial Angiosarcoma.—Long bones, 2 cases.

MEDULLARY TUMORS.

Spindle and Round-cell Sarcoma.—Long bones, 4 cases.

Perithelial Angiosarcoma.—Long bones, 2 cases.

In these 14 cases a complete operation was performed. In the 12 cases of the long pipe bones a high amputation; in the 2 jaw cases, an extensive resection. In every case death has taken place, usually within a year after the first symptom of the tumor. In only 1 case was the duration of life longer than two years.

Sarcoma of the Upper Jaw Involving the Antrum.—Clinical diagnosis: Six cases, all inoperable.

Carcinoma of Upper Jaw.—Twenty-one cases; inoperable, 16; 1 death of pneumonia; remainder not cured.

We have, therefore, observed 41 cases of tumors involving bone of a relatively high malignancy, none of which has been cured as compared with 78 of a relatively low malignancy, of which 69 are well.

These facts demonstrate the hopelessness of the more malignant varieties of bone tumors.

Surgeons who take the view advocated in this paper must educate themselves to recognize clinically, or through the Roentgen negative, or at the exploratory incision, the different varieties of sarcoma of bone, and govern the extent of the operative procedure by the relative malignancy and extent of the tumor.

In this paper I shall discuss only the benign bone cysts of the long pipe bones, the dentigerous cysts of the jaw and the cystic adamantinoma epithelioma.

BENIGN BONE CYSTS.

The benign bone cysts of the long pipe bones are rare tumors. Up to the present time we have observed but 3 cases in the surgical clinic.

CASE 1.—Bone cyst of the humerus. White girl, aged 7, tumor one year, pathologic fracture. Operation, curetting and drainage, June, 1904, ten months, well.* I saw this patient in August, 1903. The parents gave the following history:

History.—The apparently healthy child one year ago fractured the upper third of the humerus of the right arm after a slight fall. The fracture was treated by their family physician and united. After the dressing was removed a swelling was observed which has never disappeared. If the swelling had been present before the fracture, it had not attracted the attention of the parents or the physician who treated the fracture. For a year there were absolutely no symptoms except swelling. Three days ago, after a very slight fall, the child refused to use the arm because of pain, and for this reason was brought to the surgical clinic.

Examination.—On examination there was a fairly uniform expansion of the upper third of the humerus, easily seen, greatest toward the surgical neck. On palpation the soft parts were normal, but one could feel the normal shaft of the humerus expanding into a thin shell of bone. The surface of this shell of bone was not smooth like the normal shaft, but irregular. In a few places one could elicit definite parchment crepitation, first

described by Dupuytren, or what I have called "ping-pong ball" crepitation. The arm was very tender.

Clinically, a diagnosis of bone cyst was suggested, because so far in my experience every sarcoma of the more malignant type in children at this age had caused death by internal metastasis within a year.

The only medullary tumors of the long pipe bones which, in their growth, expand the bone and produce a definite shell, are the bone cysts, the myxochondrosarcoma and the giant-cell sarcoma. Of the latter two we had no observations in the clinic, in patients at this age, and so far I have been able to find none in the literature. The Roentgen negative is illustrated in Figure 1.

Operation.—At the exploratory operation the tissues were normal until the periosteum was separated. The bone beneath was irregular and varied in thickness from 1 to 4 mm. On removing a piece of the shell of bone a cavity was exposed filled with blood; there was no connective tissue lining, and no evidence of cartilage, but as only sufficient bone was removed to allow curetting of the cavity, one can not exclude the possibility that cartilage was present in some part of the wall. After curetting, the cavity was partially packed with gauze, the remainder allowed to fill with blood-clot. Eight weeks after operation an x-ray picture demonstrated that the cavity was almost completely filled with new bone.

The origin of these bone cysts has been demonstrated by Virchow, Zeronie, Schlange, Koenig and others to be due to liquefaction of misplaced islands of epiphyseal cartilage, and in the majority of cases cartilage has been found in some parts of the wall.⁷ In the two other cases observed in Dr. Halsted's clinic cartilage was demonstrated in the wall of the cyst.

CASE 2.—Colored woman, aged 37. Expansion of the lower end of the right femur, four years.

History.—The swelling shown in Figure 2 reached its greatest height about one year after the onset, and during the last three years there has been little or no increase in size. This patient refused amputation. Five years and eight months later she returned to the clinic.

Examination.—The tumor had increased in size, but had not changed its characteristics. The shell of bone was rough, similar to Case 1, but thicker, and one could not elicit parchment crepitation. The x-ray showed a shadow somewhat similar to Case 1. The diagnosis lay between a bone cyst and a myxochondroma. The long duration and the preservation of the shell of bone excluded a malignant bone tumor.

Operation.—The findings at the operation by Dr. Follis, the resident surgeon, were similar to the case just discussed, except that cartilage was present within the shell of bone in many parts of the wall.

CASE 3.—The age, clinical history, appearance and x-ray shadow were almost identical in this patient with Case 1, except that the first pathologic fracture had been five years instead of one year before the patient came under observation. The expansion of the upper two-thirds of the femur in this case was produced in the upper portion by cartilage, in the central portion by a cyst filled with blood, and in the lower portion by a fibromyxomatous connective tissue which extended down the medullary cavity of the femur some distance below the point of its expansion (Fig. 3).

Dr. Halsted exhibited this patient and the specimen at a recent meeting of the Johns Hopkins Hospital Medical Society. Later a detailed report, with the interesting histologic findings, will appear. The extensive formation of fibromyxomatous connective tissue histologically like ositis fibrosa, described by von Recklinghausen, is, I think, a unique finding in both cysts.

Koch⁸ gives the most complete résumé on the subject

G. Progressive Medicine, December, 1903, p. 191.

7. Progressive Medicine, December, 1899, p. 236.

8. Archiv f. klin. Chir., 1902, vol. lxxviii, p. 976.

of bone cysts, collecting from the literature, in addition to his one observation, 22 cases.

Heineke⁹ has recently published the first case of multiple bone cysts in which x-ray negatives were made. He is inclined to the conclusion that the cystic degeneration is part of a general osteomalacia. I have recently learned that Dr. Goldthwaite in Boston has under observation, confirmed by x-ray studies, a similar case of multiple bone cysts, in which there is no doubt clinically as to the presence of osteomalacia.

The etiology of these multiple cysts is apparently entirely different from the single cyst, but in every case numerous x-rays should be taken to exclude multiple cysts. Codman¹⁰ in Boston reports a very interesting tumor in the digital phalanx which undoubtedly represents the cartilage stage of the benign bone cyst.

The benign cyst of the long pipe bones can not always be recognized clinically, nor does the x-ray negative differentiate it from medullary giant-cell sarcoma or the myxochondrosarcoma. As a rule, these three tumors can be distinguished from the more malignant and rapidly growing medullary sarcoma. However, if there is any doubt, one should never proceed with an amputation without excluding these tumors of less malignancy by an exploratory incision.

BENIGN DENTIGEROUS CYSTS.

Ten cases. Upper jaw, 4 cases; lower jaw, 4 cases; ethmoid, 2 cases. The age of the patients varied from 6 to 30 years; 4 were under 15 years of age, 6 between 20 and 30. The duration of the tumor varied from three months to thirteen years. Whether the tumor is situated in the upper or lower jaw, it is of slow growth and usually painless. There is a slow expansion of the jaw, and on palpation one can feel a smooth, thin shell of bone. Usually there is parchment crepitation. At the exploratory incision the periosteum is normal. The outer shell of bone is smooth; lining the bone there is a thin, vascular connective tissue membrane. The contents of the cyst is usually a blood-stained serum.

Microscopically, one finds frequently cholesterol crystals, blood corpuscles and degenerated cells, which suggest epithelium. Histologically, however, I have never been able to demonstrate an epithelial lining.

Usually the cyst is single; now and then there are thin partitions. In a few cases the cysts are multiple. Complete resection is unnecessary. Partial resection with curetting and drainage will accomplish a cure. In three of our cases a non-erupted tooth was found in a recess of the cyst. These dentigerous cysts are apparently due to the distension of the connective tissue capsule of a non-erupted tooth.

CASE 4.—Figure 4. The patient was a white boy, aged 15.

History.—The swelling of the body of the lower jaw was of three months' duration. Because of the painless and uniform expansion, the distinct shell of bone and parchment crepitation, I made the diagnosis of a dentigerous cyst.

Operation.—The exploratory incision revealed the pathologic findings already described. The outer expansion was cut away with the chisel without destroying the continuity of the lower jaw. The connective tissue membrane and tooth were removed. The bone cavity was then curetted, allowed to fill with a blood-clot and the skin incision closed. The wound healed *per primam*.

Result.—June, 1904, six years after operation, there is a slight depression in the jaw at the site of the scar, but no other deformity.

CASE 5.—Figure 5. The patient was a white girl, 8 years of age, the tumor of some years' duration.

Operation.—Complete resection was done. The child's hemoglobin was but 52 per cent. at the time of the operation, and although there was no loss of blood the patient died of shock. The tumor was a very large one and extended from the zygoma almost to the symphysis of the jaw. I believe incision and curetting would have been sufficient in this case.

CASE 6.—Figure 6. The patient was a colored girl, 19 years of age, the tumor of thirteen years' duration. Complete resection was performed. (Fig. 7.) The patient died at the end of the third week from abscess of the lung.

In the remaining five cases of dentigerous cysts, two of the upper jaw and three of the lower, the operation was similar to that followed in Case 4. The patients recovered and have remained well for from two to eight years since operation.

The two cases of cysts of the ethmoid bone presented themselves clinically with a tumor projecting from the angle between the nose and the supraorbital ridge, producing slight exophthalmos. On palpation the tumor was smooth and presented a thin shell bone giving parchment crepitation. In both, partial excision with curetting and drainage was performed. The patients have remained well, one four years and the other eighteen months since operation.

ADAMANTINE EPITHELIOMA.

Of these there were 12 cases. In 4 the tumor projected from the alveolar border of the jaw (3 lower, 1 upper). The tumor was covered with normal mucous membrane and did not invade the bone. In 8 cases the epithelial tumor was situated within the body of the jaw (1 upper and 7 lower), and in its growth produced an irregular expansion very similar to a dentigerous cyst. The age of onset varied from 18 to 61 years. The majority of cases were 20 and 35 years of age; the duration of the tumor from seven months to twenty-nine years, the majority from six to twenty years. In one case the condition was considered inoperable. (Fig. 8.) In the remaining 11 cases a complete resection of the diseased area was made. Nine cases have remained well for from one to twelve years. In one instance there was a local recurrence, but this patient has remained well eight years since the second operation. One patient (Fig. 9) died after complete excision of a huge tumor involving both upper jaws.

This epithelial tumor is apparently of a very low grade of malignancy. In none of our cases was there metastasis to the glands of the neck. The tumor can be differentiated from the more malignant neoplasms of the jaw by its very slow growth. The adamantine epithelioma involving the alveolar border can not always be differentiated clinically from the connective tissue epulis. The epulis is more apt to be associated with ulceration of the mucous membrane.

When the adamantine epithelioma originates in the body of the jaw and produces expansion, with the formation of a thin shell of bone, it can not be differentiated from a benign dentigerous cyst until the exploratory incision is made. Then, when the shell of bone is incised, we do not find a cavity, but a white, finely granular tumor containing connective tissue trabeculae, and usually many small and large cystic cavities. The gross appearance is well illustrated in Figure 10. This patient (Fig. 11) was a colored man, 42 years of age, tumor of eight years' duration. The patient has remained well three years since operation.

The microscopic appearance is well illustrated in Figure 12.

9. Beiträge z. klin. Chir., 1903, vol. xl, p. 481.

10. Boston Med. and Surg. Journal, vol. cl, No. 8, p. 211, Feb. 25, 1904.

CASE 6.—Figure 13. The patient was a white woman, aged 52.

History.—Four years ago, when 48 years of age, she observed a tumor like a gumboil on the outer side of the alveolar border of the left lower jaw, opposite the canine and first molar teeth. At the end of two years, when it had reached the size of a hickory nut, it was removed. A local recurrence took place within six months.

Fig. 1.—Characteristic shadow thrown by any medullary tumor which in its growth produces a shell of bone. A pathologic fracture is also seen.

Fig. 2.—Bone cyst of lower end of femur of four years' duration.

Fig. 3.—Bone cyst of upper portion of femur. Dr. Halsted's case.

Examination.—The recurrent tumor is the size of an egg and involves the alveolar border of the left lower jaw from symphysis to within one centimeter of the angle. The tumor is present on both sides of the alveolar border. It is distinctly circumscribed. The mucous membrane at one point has ulcerated, exposing a small cyst.

Operation.—The tumor was removed by turning back flaps of mucous membrane which were not adherent to the tumor. Then the tumor and the alveolar border of the jaw were removed in one piece. The patient has remained well since the operation, a period of ten years.

Appearance of Tumor.—The fresh appearance was quite typical: white, friable, granular alveoli of various sizes and cysts in a definite fibrous stroma.

Fig. 4.—Sketch of reconstructed lower jaw in a case of dentigerous cyst. The expansion of the body, chiefly on the outer side, is well shown and the recess containing the non-erupted wisdom tooth.

Fig. 5.—Photograph of the alcohol specimen of an excised lower jaw, in which the dentigerous cysts are multiple.

Fig. 6.—This illustrates the huge size which a dentigerous cyst of the upper jaw may reach.

Under the microscope (see Fig. 13) one sees the normal mucous membrane of the gum, then a zone of connective tissue, beneath which is the circumscribed tumor. The tumor is composed of branching epithelial alveoli in a connective tissue stroma. Some of the alveoli are cysts lined by the typical basal adamantine epithelium. Other alveoli are solid, with cells showing the various morphologic changes of the adamantine epithelium.

DISCUSSION.

DR. F. J. HALL, Kansas City, Mo.—It would be of interest to know what distinction, if any, Dr. Bloodgood makes between the cystic bone tumors of the long bones; in particular, two varieties of pathologic bone condition: one is the myeloma

Fig. 7.—Photograph of the alcohol specimen of Figure 6 after removal.

Fig. 8.—Colored man, aged 54; tumor of twenty years' duration.

Fig. 9.—Dentigerous cyst of upper jaw. Death followed complete excision.

and the other is the traumatic myositis ossificans. Several cases of both these conditions have occurred in my own territory, and three of the cases of myeloma—so-called central sarcoma of the bone—have been operated on, and none has recurred. Three cases also of myositis ossificans have occurred wherein the

central portions of the tumors were occupied by these cysts, and I should like to know if Dr. Bloodgood has had any experience with these other two varieties of conditions, and what relation they bear to the conditions which he has described.

DR. JOSEPH C. BLOODGOOD, Baltimore—I am inclined to think that what you mean by myeloma is a giant-cell tumor.

DR. HALL—Yes, sir.

DR. JOSEPH C. BLOODGOOD—Dr. Hall, in his question in regard to myeloma, undoubtedly means the medullary giant-cell sarcoma. This tumor is one of a relatively low grade of malignancy. We have observed about 10 cases; all have remained well since operation. I have reported them in the Johns Hopkins Hospital Bulletin for May, 1903. In regard to the question as to the origin of the blood cysts in the condition called

Fig. 10.—Photograph of resected lower jaw; a, large smooth-walled cyst; b, bony capsule; c, smaller cysts; x, friable, white, adamantine epithelial tissue; t, teeth and symphysis of jaw.

Fig. 11.—Patient from whom tumor shown in Figure 10 was removed.

myositis ossificans, they are probably due to hemorrhage. Recently there has appeared in the literature a number of interesting articles and reports of cases on this subject. These I have reviewed in *Progressive Medicine* for December, 1903. With or without a history of trauma an indurated mass associated with some pain and tenderness is observed by the patient in one of the large muscles, most frequently the thigh. The tumor rapidly becomes bony in hardness. In some cases the x-ray demonstrates a zone of normal tissue between the shaft of the bone and the osteoid tissue in the muscle. In other cases the two bony shadows are in contact. This has given rise to two views as to the etiology of the new bone

formation in the connective tissue between the muscle bundles. A number of authorities conclude that the bone is a product of detached pieces of periosteum, others that it rises from the connective tissue cells between the muscle bundles. Clinically the condition is not difficult to recognize, especially with the aid of the x-ray. However, when the new shadow rests directly on the shaft of the neighboring bone it will be difficult to differentiate the ossifying myositis from a condition called

Fig. 12.—Microphotograph of a section of an adamantine epithelioma, involving chiefly the alveolar border of the lower jaw. In this section the basal columnar cell and its various morphologic changes toward the center of the alveolus are well shown. (Sent me by Dr. Steensland.)

Fig. 13.—Microphotograph of a section of an adamantine epithelioma of the gum.

traumatic exostosis or ossifying periostitis. Blood cysts have been observed only in the ossifying myositis. Extensive operations are not necessary. One should remove as much of the new bone production as possible without destroying function. Slight local recurrences of the bone formation are to be expected. Fortunately the amount is never great, gives little or no discomfort, and second operations are rarely necessary. I

do not think there is any relation between the hemorrhagic cysts observed in myositis ossificans and the cysts of the long pipe bones.

As to the question of the origin of the giant cells, recent investigation would indicate that the periosteal or medullary giant-cell tumor is probably an angioma or an angiosarcoma, and that the giant cells are due to budding of the endothelial cells of the vessels in these very vascular tumors. The most interesting publication on this subject is by Friedländer.* Further investigation, however, should be made of this most interesting tumor.

EXPERIMENTALLY PRODUCED GENUINE EPITHELIAL METAPLASIA IN THE STOMACH,

AND THE RELATIONS OF EPITHELIAL METAPLASIA TO CARCINOMA, AS DEMONSTRATED BY CASES REPORTED IN THE LITERATURE.

GUSTAV FÜTTERER, M.D.

CHICAGO.

As the observations on epithelial metaplasia to be reported in this paper were made in a series of experiments on the stomach of rabbits, undertaken for the purpose of studying the relations between mechanical irritation and development of carcinoma, I think it necessary to state this fact, merely to show under what circumstances the findings were made. My views concerning the rôle which mechanical irritations play in the etiology of carcinoma, and how those views have been formed, have been fully explained in former writings, and I can, therefore, refer the reader to them, in order to prevent repetitions, which would make this article too extensive.¹

Some observations of importance have been made which I do not report, because they need corroboration, but they certainly encourage further efforts along the same lines.

In a series of experiments on the stomachs of 94 rabbits, a genuine epithelial metaplasia was found in five cases, only two of which will be described in detail, as a description of the others would cause an unnecessary repetition, for while the findings differ somewhat in all the cases, they are the same in principle. It may also be mentioned that I have specimens of epithelial metaplasia from a former experiment, bringing the total of cases to six.

In each case a piece of mucous membrane of the stomach was resected and hypodermic injections of a solution of pyrogallie acid were made, to reduce the percentage of hemoglobin, and thus prevent the healing of the defect, the edges of which were then exposed to friction by solid food. At first, as in former experiments, circular resections of mucous membrane around the pylorus were made, but the feeding of the animals was not carried out according to orders given, and they all died, postmortem examination revealing very much dilated stomachs filled with solid food, while only liquid food was to be given during the first four days. Not being able to get better care of the animals at the time, I had, in order to reduce the mortality, to resect pieces of mucous membrane near or further away from the pylorus, and later, when Dr W. H. English, to whom I am very much indebted, kindly took it on himself to supervise the keeping of the animals, my results became very much better.

* Archiv f. klin. Chir., 1902, vol. lxxvii, p. 202.

1. Fütterer: Aet. d. Carc., Bergmann, Wiesbaden, 1901, and THE JOURNAL A. M. A., March 15, 1902.