prominent, punctures larger than on rest of face, partly confluent; face covered with short black hair; labrum with a distinct median depression; mandibles black, tips rufo-testaceous, notch one-fifth of length from the blunt tip, strongly grooved without, space between eyes and base of mandibles not as great as width of latter; antennæ short, black, flagellum deep brown beneath, reaching to line of tegulæ; cheeks sparsely fringed with short, black hair; dorsum of the thorax with short, sooty hair, some black hair intermixed in spots, disc shiny, sparsely pubescent; pleura with black hair; thorax quite evenly sparsely punctured, post-scutellum more finely punctured, base of metathorax with transverse series of pits, triangle shining, not smooth; tegulæ shining, distinctly piceous; wings hyaline, nervures and stigma testaceous, marginal cell very dark, second submarginal narrow at top, third not narrowed as much as usual in one specimen; legs black with sooty pubescence, tarsi reddish with rufo-testaceous hair, first joint very dark; abdomen black, punctured, first two segments shining, white hair-bands on segments 1-5, on 1 and 2 interrupted, otherwise sparsely pubescent with black hair, venter with very short black hair.

Described from two females: Ft. Collins, Colo., 6-viii.-96 (Gillette), and Horsetooth Mt., Colo., 22-vi.-99, on *Potentilla*. This species differs from known Colorado species by the black hair on the face and pleura and the sooty hair on thorax. Prof. Cockerell writes that it is closely related to *C. pascoensis*, Ckll., from Washington; but differs by its smaller size and by possessing hair-bands.

I wish to acknowledge the kindness of Prof. Cockerell and Mr. Robertson for favours shown me in revising portions of my manuscript, and for the general help they have given me.

NOTES ON SOME NORTH AMERICAN SPECIES OF TINEIDÆ.

BY HARRISON G. DYAR, WASHINGTON, D. C.

About the time that Lord Walsingham's valuable paper on *Acrolophus* and *Anaphora* appeared (Trans. Ent. Soc., Lond., 1887, 137-173), Mr. Beutenmüller was working on the same group; but neither author has since attempted to recognize the species named by the other, so far as I am aware. In Prof. Smith's List Lep. Bor. Amer., 1891, the group is recognized as a family—*Anaphoridæ*—but this can hardly stand. The genera will fall in the *Tineidæ*, in the more restricted sense (see Walsing-

ham, Proc. Zool. Soc., London, 1897, 139-175). The following synopsis of genera is from Walsingham, with addition of two genera described since the original publication :

ANAPHORINÆ.

Palpi erect or slightly recurved.

Veins 8 and 9 of fore wing stalked.

Palpi erect.....*Eulepiste*.

Palpi appressed to head.....*Neolophus*.

Veins 8 and 9 separate.

Antennæ bipectinate.....*Ankistrophorus*.

Antennæ simple or serrate towards apex.

Tarsal joints of hind legs strongly fringed above...*Thysanoskelis*.

Tarsal joints not fringed above.

Palpi erect, with separate tufts on each joint.....*Ortholophus*.

Palpi slightly recurved, uniformly hirsute.....*Pseudanaphora*.

Palpi strongly recurved.

Antennæ bipectinate.....*Felderia*.

Antennæ serrate throughout.

Veins 8 and 9 of fore wings stalked.....*Cænogenes*.

Veins 8 and 9 of fore wings separate.....*Anaphora*.

Antennæ simple, compressed, or slightly serrate at ends.

Veins 7 and 8 of fore wings stalked.....*Atopocera*.

Veins 8 and 9 of fore wings stalked.

Head with an erect crest.....*Urbara*.

Head without an erect crest.....*Hypoclopus*.

No veins of fore wings stalked.

Palpi roughly clothed throughout.

An erect fringe along lower margin of cell on hind

wings.....*Pilanaphora*.

No such erect fringe on hind wings.....*Acrolophus*.

Palpi smooth, the last joint only tufted.....*Stæberhinus*.

Genus EULEPISTE, Walsingham.

Wals., Trans. Am. Ent. Soc., X., 169, 1882 ; Trans. Ent. Soc., Lond., 1887, 142.

Synopsis of Species.

Uncus single, the opposing lower limb not half as long as the upper limb.

Harpes concave, rounded at the ends.....*Cressoni*.

Harpes concave, obliquely truncate above.....*maculifer*.

Uncus single, its opposing lower limb nearly as long as the upper and stouter.

Harpes nearly flat, the ends bent inward, rounded. *Cockerelli*.

Eulepiste Cockerelli, n. sp.

Palpi upturned in front of the head, free, as high as the vertex ; male antennæ simple, slightly serrated toward the ends ; veins 8 and 9 of fore wing stalked ; thorax and fore wings dark brown-gray, somewhat grizzled or mottled with darker and with an obscure darker spot at the end of the cell. Hind wings dark brown ; abdomen gray-brown ; expanse 16 mm. One male, Mesilla Park, New Mexico, at light, July 8th (T. D. A. Cockerell) ; U. S. Nat. Mus., Type No. 4417.

Genus HYPOCLOPUS, Walsingham.

Wals., Trans. Ent. Soc., Lond., 1887, 141.

Synopsis of Species.

Uncus single, down curved, broad at base, tip pointed ; harpes broad, narrow at base, squarely truncate. *griseus*.

Uncus double, the two points separate, sharply down curved, the ends bent outward, enlarged and rounded ; harpes long, narrow and uniform *mortipennellus*.

Hypoclopus griseus, Walsingham.

Wals., Trans. Ent. Soc., Lond., 1887, 144.

Arizona (Morrison, from Lord Walsingham, through C. V. Riley) ; San Diego, Texas, May 9 (E. A. Schwarz) ; Oracle, Arizona, June 28 (E. A. Schwarz) ; Brownsville, Texas, Apr. 27 (C. H. T. Townsend) ; Washington, D. C., July 20 (A. Busck).

The Texas specimens are pale, the ground colour an ashy white, on which the dark specks and streaks show plainly. The specimen from Oracle, Ariz., is very dark, the black markings predominating.

Hypoclopus mortipennellus, Grote.

Grote, CAN. ENT., IV., 137, 1872 ; XVIII., 199, 1886. Wals., Trans. Am. Ent. Soc., X., 167, 1882 ; Trans. Ent. Soc., Lond., 1887, 150 ; *quadripunctellus*, Beut. (ined.), Smith's List Lep. Bor. Amer., No. 5057, 1891.

This species, described as *Anaphora* and placed by Lord Walsingham in *Acrolophus*, may be removed to *Hypoclopus*, as a majority of the specimens have veins 7 and 8 of fore wings stalked, at least on one side. Of twelve specimens before me, seven have these veins stalked on

Florida (coll. Beutenmüller, type of *angustipenellus*); Georgia (coll. Beutenmüller); Orange Co., Florida (coll. Beutenmüller); Texas (Boll. coll. C. V. Riley, identified by Walsingham); Columbus, Texas, June (E. A. Schwarz).

Acrolophus violaceellus, Beutenmüller.

Beut., Ent. Amer., III., 139, 1887.

The genitalia are not very different from those of *Arizonellus*, Wals., though the harpes may be somewhat slenderer. However, the moth differs in its uniform, unspotted, purplish colour.

North Carolina (Beutenmüller's types); Iowa (from Dept. Agriculture).

Acrolophus Arizonellus, Walsingham.

Wals., Trans. Ent. Soc., Lond., 1887, 153.

Arizona (Morrison, from Lord Walsingham, through C. V. Riley); Mesilla, New Mexico, June 25 and July 1 (T. D. A. Cockerell); Tucson, Arizona, July 19, 20 and 21 (E. A. Schwarz).

Genus ANAPHORA, Clemens.

Clem., Proc. Ac. Nat. Sci., Phil., 1859, 261; Wals., Trans. Ent. Soc., Lond., 1887, 155.

Anaphora popeanella, Clemens.

Clem., Proc. Ac. Nat. Sci., Phil., 1859, 261; Wals., Trans. Ent. Soc., Lond., 1887, 161 (references and synonymy); Riley, Smith's List Lep. Bor. Am., No. 5061, 1891; *confusellus*, Beut. (ined.), Smith's List Lep. Bor. Am., No. 5056, 1891.

The form *confusellus* is smaller than the usual form, the ground colour lighter and more purplish, the dark marks strongly relieved. In genitalia there is no marked difference. The form differs from *popeanella* about as much as *Acrolophus cervinus*, Wals., does from *A. plumifrontellus*, Clem. U. S. Nat. Mus., type No. 405.

Georgia (Beutenmüller's type); Kirkwood, Missouri? (labelled only "148 M," *i. e.*, Murtfeldt); Georgia (A. Oemler, labelled "Anaphora, n. sp., Wism., '86"); Georgia (labelled "Anaphora, n. sp., doubtless = *plumifrontellus*, C. V. R., '86, with Wism."); eight specimens, badly worn (labelled only "667, Aug., '80, coll. C. V. Riley").

Genus FELDERIA, Walsingham.

Wals. Trans. Ent. Soc., Lond., 1887, 165.

Felderia filicornis, Walsingham.

Wals., Trans. Ent. Soc., Lond., 1887, 165; *Mexicanellus*, Beut., Ent. Amer., IV., 29, 1888.

Arizona (Morrison, from Lord Walsingham, through C. V. Riley); City of Mexico (Beutenmüller's type of *Mexicanellus*); Oracle, Arizona, July 12 (E. A. Schwarz); Fort Grant, Arizona, July 20 (H. G. Hubbard); Brownsville, Texas, June 10 (C. H. T. Townsend).

Genus ORTHOLOPHUS, Walsingham.

Ortholophus variabilis, Walsingham.

Wals., Trans. Ent. Soc., Lond., 1887, 169.

Arizona (Morrison, from Lord Walsingham, through C. V. Riley); Arizona (coll. Beutenmüller); Arizona (yellow labels 5, 7 and 8 marked "prob. undescribed," Wals., 1886); Oracle, Arizona, July 8, 10, 12, 16 and 24 (E. A. Schwarz); Fort Grant, Arizona, July 19, 20 and 22 (H. G. Hubbard); Tucson, Arizona, July 21 (E. A. Schwarz); Chiricahua Mts., Arizona, July 4 (H. G. Hubbard); Mesilla Park, New Mexico, July 8 and Aug. 13 (T. D. A. Cockerell); Texas (coll. Beutenmüller); Sharpsburg, Texas, May 11 (E. A. Schwarz); Glenwood Springs, Colorado (W. Barnes); Nevada.

Genus PSEUDANAPHORA, Walsingham.

Wals., Trans. Ent. Soc., Lond., 1887, 170; *Eutheca*, Grote, Bull. Geog. Surv. Terr., VI., 257, 1881; Dyar, CAN. ENT., XXVII., 15, 1895; *Sapinella*, Kirby, Cat. Lep. Het., 524, 1892.

Synopsis of Species.

Uncus single, long and slender.....*Davisellus*.

Uncus double, two spines projecting from a rounded plate....*arcanella*.

Pseudanaphora Davisellus, Beutenmüller.

Beut., Ent. Amer., III., 139, 1887.

Though described as an *Acrolophus*, this is obviously referable to *Pseudanaphora*, from the short erect palpi.

Arizona (Beutenmüller's type); Fort Grant, Arizona, July 20 (H. G. Hubbard).

Pseudanaphora arcanella, Clemens.

Clem., Proc. Acad. Nat. Sci., Phil., 1859, 262; Wals., Trans. Ent. Soc., Lond., 1887, 170 (references); Beut., Ent. Amer., IV., 29, 1888; Forbes, 16th Rept., Ill., 98, 1890; *mora*, Grote, Bull. U. S. Geol. Surv., VI., 257, 1881; Kirby, Cat. Lep. Het., 524, 1892; Dyar, CAN. ENT., XXVII., 15, 1895.

Rhinebeck, New York (Dyar); Rhinebeck, N. Y., July 5 (Miss L. J. Hoff); Fordham, N. Y. (G. Gade); Staten Island, N. Y., June 25 and July 16 (coll. Beutenmüller); District of Columbia, July 18 (coll. C. V. Riley); Washington, D. C., October 10 (A. Busck); St. Louis, Missouri, issued July 3 (C. V. Riley, breeding No. 2563); Texas (coll. Beutenmüller). The specimen taken in October is very dark in colour, blackish, the markings being only faintly indicated.

NEW COCCIDÆ FROM CALIFORNIA.

BY EDW. M. EHRHORN, MOUNTAIN VIEW, CAL.

Xylococcus quercus, n. sp. (Plate 7, figs. 1 and 2.)

Egg quite large, of a light orange colour.

Young larvæ dark orange-red, active, body broadly oval, about $\frac{2}{3}$ mm. long. Legs and antennæ light brown, well developed. Antennæ short, 6-jointed. Joint 1 stoutest, joint 6 longest, and joint 4 shortest. Formula: 651234. Joints 2 and 5 with three bristles. Joint 6 with numerous long stout bristles. Legs moderately long, with femur quite swollen. Tarsus longer than tibia. Digitules of tarsus fine hairs; those of claw long stout clubs curved upwards. Each segment of abdomen bears a backward directed short stout spine. On each side of anal tube is a long fine bristle. Anal tube large, with numerous stout spines. Stigmatal tubes well developed.

♀ second stage, body crimson, shiny, nearly spherical, about $1\frac{1}{2}$ mm. long, 1 mm. broad, surrounded by cottony and waxy secretion. Antennæ and legs wanting. Anal tube well developed, producing a glassy rod, like a stout white hair, rather brittle. Last segment of body dark brown. When cleared in K. H. O., surface of body finely granulated, more so near caudal end. Stigmatal tubes are large and well defined. There are numerous spines and gland openings scattered over the body.

♀ third and fourth stages very similar to second stage, but larger in each case from the preceding, and varying in the further development of stigmatal and anal tubes, glands, spines, etc.

Adult ♀ head, thorax, legs and antennæ reddish-brown, abdomen blackish-brown, segmentation distinct. There is a distinct constriction between the thorax and abdomen. Length of body about $5\frac{1}{2}$ mm., breadth $2\frac{1}{2}$ mm., quite convex above. Ventral side of abdomen concave, with revolute margins. Insect quite active. When ready to