time that this is really the explanation of the poor reactions in certain cases. In our opinion it is more likely that the secondary invading organism had begun to develop its lesion in the lung and that a new and subsequently fatal factor was added. In this district it is almost entirely the pneumococcus. From the data at hand, we cannot state that there is a noteworthy difference among donors, in regard to the clinical results produced by the injection of their blood. Those having a leukopenia or a leukocytosis and those who pursued a mild or severe clinical course seemed about the same.

The leukocyte count followed very closely a general rule. A leukopenia of from 3,000 to 5,000 was present at the onset of the influenza in a vast majority of cases. With the onset of pneumonia a very moderate leukocytosis, one of from 9,000 to 12,000, occurred in the majority. At times it was higher. We had, however, but eight counts between 20,000 and 30,000 in the 150 counts estimated. This finding is in marked contrast to that observed in the lobar pneumonia.

In some, particularly the very severe pneumonia cases, the leukopenia persisted throughout the course. A drop in leukocytes was a serious prognostic sign during the course of the pneumonia. We have very slight evidence of an influence on the leukocyte count by the blood transfusion. In two or three cases the increase was well marked, from 5,000 to 10,000, but in the remainder there was very little change, whether we began with a leukopenia or a moderate leukocytosis.

SUMMARY

The use of citrated blood from convalescent influenzal patients administered to patients when pneumonia followed the influenza appeared to have a decided beneficial effect, particularly if given early. Beyond the third day of the pneumonia it was of little value.

That the citrated blood establishes an immunity to *B. influenzae* is inferred but not proved. Many additional factors may play an important rôle. Extensive work along these other lines was conducted by the University of Pittsburgh during the recent epidemic, a report of which will appear later.

Suggestion by the Patient.—We are often misled by our patient's interpretations of his own case. In other words, we are human and subject to the power of suggestion. Often we are informed that they are suffering from heart disease, kidney or liver disease, or appendicitis, constipation or diarrhea, and we listen to their story and become hypnotized ourselves and follow along the line of the patient's suggestions. They are anxious to tell somebody their symptoms and if we are good listeners they are willing to pay us for our trouble and time for listening.—President's Address, Minnesota State Medical Society.

Clinical Notes, Suggestions, and New Instruments

ONE MAN WITH FIVE TESTES: REPORT OF CASE

GEORGE H. DAY, M.D. (LOUISVILLE)

Major, M. C., U. S. Army; Camp Venereal Officer.

CAMP PIKE, LITTLE ROCK, ARK.

This case was referred to the urologic department, July 1, 1918, for confirmation by Major H. C. Loos, chief of the surgical service at Camp Bowie.

The man in question was discovered by Major Loos at a local garage, where he was employed as a mechanic. After questioning the man and hearing his history, Major Loos was interested enough to have him report to the base hospital for examination. The case has since been presented to the Clinical Society of the Base Hospital, Camp Bowie, for examination.

G. E. M., aged 19, of San Benito, Texas, presented a negative family and personal history other than that his father had died of typhoid ten years before.

He had had gonorrhea twice, the first time at 17 years, and again six months before we saw him. The first time he knew he was other than normal was two years before, when on attempting to enlist in the service, he was rejected at Brownsville, Texas, by the recruiting officer of the Twenty-Sixth Infantry. At that time he was told that he was ineligible on account of severe varicocele. He attempted to enlist again at Oklahoma City one year later with the same result.

Three months before he again attempted enlistment at Fort Worth, Texas, and was again rejected with a diagnosis of varicocele and hydrocele.

The man was well nourished, and aside from being overdeveloped, with the contour of the arms rather feminine and the pelvis and

hips somewhat large, he appeared to be normal.

It was immediately noticed that the scrotum was much larger than usual. The penis was about twice the size found in boys of this age.

On retracting the scrotum it was noticed immediately that the young man had another normal scrotum and perhaps a third, which was rudimentary in origin, to the right of the normal one. On further examination four normal testes were discovered, two being in each of the two scrota and another atrophied one being in the redundant tissue posterior to the second pair. The third or rudimentary scrotum in the groin (right) contained no trace of the sixth, and may possibly be intra-abdominal. The line of demarcation in the tissue of the scrotum was most marked and showed forcibly the fact of the scrota.

When he assumed the knee-chest position, the posterior view strikingly brought out the line of demarcation between the scrota. Aside from an overabundant sexual desire he said he had never noticed anything abnormal in his sexual life.

We have not had the opportunity of looking up the literature on the subject and are presenting it simply as an anatomic freak. Certainly the man is possessed with four normal

Scrota with five testes; white arrow points to line of demarcation between scrota; black arrow, rudimentary scrotum; 1, 2, 3, 4, 5, testes.

testes and cords, a fifth atrophied testicle and perhaps a sixth which is undescended.

It would be interesting to see a section to note where the attachment of the second set joins the vas of the first, and it is not at all improbable that he has also multiple seminal vesicles, although only one prostate is palpable.

A SHORT METHOD OF MAKING NOGUCHI ACETONE-INSOLUBLE ANTIGEN

G. A. MAGNUSSON, M.D., SEATTLE

In order to gain the most information from the Wassermann test, more than one antigen should be used in the "set up." One of the antigens so used should undoubtedly be the

"Set up" used in titration.

Noguchi acetone-insoluble, as it will give positive reactions with a higher percentage of syphilitic serums than the alcoholic extracts, while it is less likely to give false positive reactions than the more sensitive cholesterinized extracts. But its preparation requires so much time that this becomes quite a factor in a laboratory where considerable quantities are used. With a view of hastening the process of extraction and evaporation we tried this at higher temperatures and find that boiling temperature in no way interferes. In this way we can prepare and titrate an antigen all in one day.

PROCEDURE

To one part of finely minced beef heart, nine parts of commercial ethyl alcohol are added. (We have recently tried the extraction with denatured alcohol and this seems to do as well. As alcohol is denatured according to varying formulas, some may not be usable.)

This is boiled in a water bath for from ten to fifteen minutes, and filtered while hot.

The filtrate is evaporated over a boiling water bath until the alcohol and water are removed.

The residue is extracted with a small amount of ether.

The ethereal extract is poured into a centrifuge tube and centrifuged until the supernatant liquid is perfectly clear. This is added to ten volumes of chemically pure acetone.

The precipitate is allowed to settle. The greater part of the acetone is decanted, and the rest is poured into a clean graduated centrifuge tube and centrifuged for five minutes at moderate speed, the acetone poured off, and drained thoroughly.

To the precipitate, ten volumes of chemically pure methyl alcohol are added, and the mixture is stirred with a glass rod until as much as will dissolve has gone into solution and then centrifuged until the supernatant liquid is perfectly clear.

This is poured off and kept in a clean, well stoppered bottle.

TITRATION OF ANTIGEN

An emulsion is made with one part of this stock alcoholic solution and nine parts of physiologic sodium chlorid solution.

Titration is performed for antigenic, anticomplementary and hemolytic properties.

The accompanying chart illustrates the "set up" used in titration.

To each tube, physiologic sodium chlorid solution is added to bring the volume to 1.5 c.c.

Incubation is performed for one hour.

To each tube 0.5 c.c. of sensitized sheep cell suspension (5 per cent. sheep blood cells and 2 units of amboceptor) is added.

Incubation is performed for one hour.

Most of the antigen preparations we have made in this way have had an antigenic unit of 0.02 c.c. of the 1:10 emulsion, while a full cubic centimeter was neither hemolytic nor anticomplementary.

TREATMENT OF CHRONIC GONORRHEAL SKENITIS WITH THE ELECTRIC CAUTERY*

CHESTER B. MOORE, B.S., M.D., SAN FRANCISCO

A not uncommon pelvic focus for the lodgment of the gonococcus is the gland of Skene. In this minute epithelial pocket such an infection may persist indefinitely in spite of local applications. Even if the ducts are exposed and medicinal injections directed into them with a hypodermic needle, the infection may continue without interruption. Often it is possible to express a few or several drops of pus week after week, and month after month, along with such oft-repeated treatment. Furthermore, this minute gland may remain in a chronic state of suppuration without a local symptom, the focus of infection being discovered only in the course of a general pelvic examination. Unless one makes an attempt to express the contents of these ducts, this focus may be entirely overlooked. In this type of gonorrheal infection of Skene's gland nothing less than the complete destruction of the gland will terminate the disease in this locality. This destruction may be easily attained by the use of the thermocautery.

The operation is quite easy and simple, provided one can get a good exposure. For this purpose I take a Kelly endoscope that has been cut off at the proper length so that it will not extend into the bladder, close it with the olive tip, and cut out a small window as shown. I have found that a No. 8 endoscope serves the purpose very well. This makes an instrument, a skenoscope, with a handle that is directed upward out of the way; the entrance is expanded to hold back and protect the labia; the tubal portion protects the urethra except at the point of attack, the gland, the ducts of which protrude into the window cut for the purpose. Under local anesthesia, it is possible to pass a small wire thermocautery

Front, side and top views of instrument devised for use in the treatment of chronic gonorrheal skenitis with the electric cautery.

into the gland without fear of cauterizing any other place than that desired. In order to obtain good light, I have had a small post so placed on the handle that it will carry an E. S. I. light directed toward the aperture. However, any means for obtaining good light will do.

To cauterize the gland, as much discharge as possible is expressed from the ducts. The urethral entrance is cocainized with a 10 per cent. solution for about ten minutes. The instrument is passed, handle upward, until a good exposure is obtained. The entrance of the ducts is sought with a large size hypodermic needle, from which the point has been filed, or with the wire contained in the tubal portion of the instrument. The needle or wire is passed along the duct to the

* From the Division of Obstetrics and Gynecology, Leland Stanford Junior University School of Medicine.