DISCUSSION.

Marcus Feingold, New Orleans. It is unfortunate that in the case reported the skin condition was not seen by the dermatologist, though no reason exists to dispute the diagnosis of Dr. Stark. Yet it is possible that the eye condition may have had another interpretation. The following case is of interest in connection therewith. A young woman of low mentality, aged about 25 years, appeared at the Touro Infirmary Out-Patient Clinic with a chalazion on the left upper lid, telling that she had been taking some medicine and using a lid ointment prescribed by her physician. On everting the lower lid a white opaque area was found in the conjunctiva of the lower lid and fornix near the inner canthus. The suspicion at once arose that we were dealing with a burn of the conjunctiva occasioned by the use of mercury while the patient was taking iodid of potash. Acting upon this, only bland treatment was used, but the condition remained unchanged during the next few days, possibly the area getting somewhat smaller.

A few days later to our great surprise the opaque necrotic area appeared much larger, the balance of the conjunctiva, at the same time, being remarkably free from symptoms of inflammation. Soon a similar smaller spot was discovered in the other eye in the identical location. The former diagnosis was now abandoned and the strong suspicion of induced conjunctivitis arose. Pharyngeal reflex was found entirely absent and the neurologist reported the patient suffering from dementia praecox. During the

next four or five weeks the condition remained practically unchanged but for some slight exacerbations and remissions first in one eye, then in the other one. Admitted to the hospital for observation and close watching, and without any local treatment, the conjunctiva returned to normal within a few days and remained so during the whole stay of six weeks in the hospital. In the case of Dr. Stark it is possible that the condition began as described; but that the child being made the center of attention in the family, may have kept up the irritation to maintain this interesting position.

Harold Gifford, Omaha, Nebraska. The fact that Dr. Stark's case extended to the upper tarsus would be against the idea of malingering.

H. H. Stark, El Paso, Texas. My experience of malingering is that the disturbance is more apt to be on the temporal than on the nasal side. I think it is absolutely impossible for malingering to have occurred in this case. Fuchs calls attention to the fact that the condition may exist over many years. The child is the daughter of our mayor, and her mother is a very intelligent woman. The child had not had any attention called to her eyes for four years, all treatment having been omitted.

Dr. Gifford asked whether Dr. Stark had tried staining with fluorescein after removing the membrane. Dr. Gifford had found fluorescein as useful in the conjunctiva and in the fornix as on the cornea.

H. H. Stark (closing). No. There were no granulations, but there might perhaps have been some staining.

CORNEO-SCLERAL TREPHINING.

C. A. CLAPP, M. D., F. A. C. S.

BALTIMORE, MD.

A report of five cases subjected to this operation which resulted in as many disappointments.

While the operation of corneo-scleral trephining is theoretically ideal, and in many hands seems to elicit only the highest terms of commendation, in others, there have been noted subsequent complications which have caused disappointing end results. I think our experience has been unusual in that all cases operated upon have proven in the long run to be disappointing. While I say *all* cases and then show that this is only five in number, still there are certain features about some of them that make them of considerable interest.

Some of the unsatisfactory conditions that have been reported are: Late infection, latent iritis, fulminating glaucoma, and hypotony with poor vision; all of which have occurred in this series of cases.

Following is a brief report of case histories with comments:

Case I.—W. S. Colored. Age 55. Presented himself October 7, 1912, with an acute glaucoma in his right eye following a blow upon the eye, left eye showed absolute glaucoma. Myotics failing to reduce tension, even in ab-

sence of light perception, it was decided to trephine. The operation was performed February 3, 1913, with a small iridectomy. Healing was uneventful, but the trephine opening seemed to fill with organized exudate, and tension to increase until March 3, 1913, when a tension of plus three, (fingers), was noted. No pain being experienced, nothing further was done, and the patient failed to report thereafter.

Case II.—P. R. Hebrew. Age 25. January 27, 1913, gave history of poor vision in left eye for one month, with pain for the past week. Examination shows deep anterior chamber, floating opacities in vitreous, myopic crescent about each nerve, left nerve decidedly cupped and pale.

Myotics ordered.

$$V = \text{O.D.} - 16 = 15/40$$

$$V = \text{O.S.} - 14 = 15/40$$

$$\text{Tension O.D. 15 mm.}$$

$$\text{Tension O.S. 40 mm.}$$

Tension remaining high; on February 13, trephining of left performed. There was only moderate reaction.

On February 26,

$$\text{Tension O.D. 18 mm. Hg.}$$

$$\text{Tension O.S. 18 mm. Hg.}$$

On March 24,

$$\text{Tension O.D. 30 mm. Hg.}$$

$$\text{Tension O.S. 40 mm. Hg.}$$

April 14, left pupil dilated. Tension 50 mm. Hg. Tension remained high even under myotics. $V =$ in O.S. on January 28, 1914, was 2/200, nerve becoming more atrophic.

March 16, 1914, patient returned with a violent attack of acute conjunctivitis. March 18, 1914, the marked bleb over trephine opening had broken with flattening, and extensive fibro-purulent exudate filled anterior chamber. Under treatment in the hospital the eye cleared up so that on April 1, 1914, it showed clear anterior chamber, large floating opacities in vitreous, tension reduced, nerve white, and vision hardly more than fingers. We have been unable to get the patient to report for further examination.

Case III.—E. K. White, widow. Age 62. On March 3, 1913, gave history that left eye was operated on twelve years ago for failing vision without re-

sult. That vision in right eye has been failing for six months, and she has been using drops from a hospital for the same. Examination showed operative coloboma upward in left eye with absolute glaucoma. Right eye showed simple chronic glaucoma with pitting, but normal tension. Continuance of myotics ordered. Tension 20 mm. March 19, 1913, tension O.D. 50 mm. Blood pressure 220. Patient ordered stronger myotics and blood pressure treatment, tension still remained above 40 mm. Hg., and in view of the unsuccessful iridectomy on left eye trephining was advised for right, which was performed May 26, 1913, using $1\frac{1}{2}$ mm. trephine. Eye did well for two days, when fulminating glaucoma developed and eye was lost, to absolute glaucoma.

Case IV.—M. O'N. White man of 56. July 19, 1913. Gave history that left eye had been failing for several years and right for six months. Had been under care of a recognized oculist. Examination showed simple chronic glaucoma in each eye with tension of O.D. 37 mm. Hg., O.S. 40 mm. Hg.

$$V = \text{O.D.} + 1.75 = 20/20$$

$$V = \text{O.S. Fingers in temporal fields.}$$

Patient was treated by eserine and pilocarpin until December 4, 1913, with no loss of vision and tension below 35 mm. Hg. Upon the advice of Colonel Elliot a bilateral corneo-scleral trephining was performed by the Colonel. There resulted very little reaction and tension soft.

On February 6, 1914, $V = \text{O.D. } 20/100$, O.S. Motion.

On May 4, 1914, tension was O.D. 13 mm. Hg., O.S. 13 mm. Hg.

March 24, 1915, $V = \text{O.D. } 20/100$, O.S. Motion.

May 20, 1917, patient returned with small areas of infiltration in upper part of right cornea. No change in vision.

Case V.—K. K. White, female, single. Age 37. July 3, 1913. History of failing vision in each eye for some time.

$$V = \text{O.D. } 15/200 \quad 2.25 = 15/40$$

$$V = \text{O.S. } 15/200 \quad 2.25 = 15/40$$

Diagnosis: Simple chronic glau-

coma with cupping and contracted fields, worse in right.

Tension O.D. 37 mm. Hg.
O.S. 37 mm. Hg.

Left field showed greater contraction and also less effect from myotics and so left eye was trephined on February 23, 1913.

March 10, 1914, tension finger normal with vision in O.S. of 20/50 plus.

April 8, 1914,

Tension O.D. 25 mm. Hg.
O.S. 9 mm. Hg.

Operation upon right eye was advised and thinking it gave a good opportunity to note the relative merits of trephining against iridectomy the older operation of iridectomy was performed on right eye May 12, 1914.

June 11, 1914, with correction,

V = O.D. 20/80
O.S. 20/30

On July 20, 1914, patient returned complaining of some pain in left eye, which showed some ciliary congestion. Patient did not report again until August 7, 1914, when well-marked case of iritis was present in left and atropin ordered.

March 10, 1914,

Tension O.D. 20 mm. Hg. with al-
O.S. 17 mm. Hg.

most complete posterior synechia and beginning opacity in lens of left eye. Atropin viii gr. ordered. March 29, 1915, left lens completely opaque.

July 20, 1915, removed lens by incision above, slight loss of vitreous. Cornea started to become opaque from above and has continued until most of tissue is opaque and vision of light perception only remaining.

April 16, 1917,

V = O.D. with correction 20/100++
O.S. Light perception.

July 19, 1917, left cornea steamy. Tension plus. Enucleation advised.

COMMENTS.

Case I. This was our first experience with the operation of trephining and not a very favorable case. The operation was performed to see if the ten-

sion would remain lowered without expecting restoration of vision. But neither was there permanent lowering of tension nor restoration of vision.

Case II. This case showed satisfactory lowering of tension for the first month; but it did not remain permanent even with the presence of a patent opening, with marked bleb over opening; and later showed a typical late infection, just about one year after the primary operation. Although the anterior chamber infection cleared up the vision was not restored.

Case III. This case one would have thought to have presented proper indications for the corneo-scleral operation, as iridectomy had failed on the left, and myotics failed to keep tension down in the right. Later writings and our own experience tend to make us think we erred in using the smaller trephine, although we believe that this represents one of those fulminating cases where all procedures are without results.

Case IV. The first three cases did not have the same skill in execution that this case had and yet the filtering scars were all present and could be demonstrated by pitting. This case had a vision of 20/20 in the O. D. and the tension was kept down with myotics prior to operation. Since operation there has been even hypotension but the vision has never been better than 20/100. Would not this case have done better under prolonged myotics? Does not the beginning corneal infiltration result from the hypotension? Certainly we can find no fault with the operative technic.

Case V. This case probably presents better conditions for our experimental control than any of the others. The Elliot operation was performed on the better seeing eye. The filtering scar and resulting hypotension were perfectly satisfactory; and yet a latent iritis developed five months after the operation with subsequent blindness; while the old classic van Graefe iridectomy on the worse eye, still retains useful vision and satisfactory tension.

CONCLUSIONS: First—The corneo-

FIG. 1. CASE I. - NOTE POSITION OF TUMOR, AND LACK OF PIGMENT IN ANTERIOR THIRD.

FIG. 2. CASE I. - ANTERIOR SURFACE OF IRIS, POSTERIOR SURFACE OF CORNEA AND ANGLE OF ANTERIOR CHAMBER COVERED WITH TUMOR CELLS.

FIG. 3. CASE I. - ECTROPION OF PIGMENT LAYER OF IRIS.

FIG. 4. CASE I. - AREA FROM UNPIGMENTED EPIBULBAR PORTION OF TUMOR.

FIG. 5. CASE II. SECTION FROM METASTATIC NODULE IN LIVER.

FIG. 6 CASE III. - GROSS SPECIMEN NOTE TUMOR ON OPTIC STALK.

MELANOBLASTOMA OF THE EYE
FORMAN AND HUGGER.

FIG. 7, CASE III.—IRIS BOUND TO CILIARY BODY BY GRANULATION TISSUE.

FIG. 8, CASE III.—TUMOR IN OPTIC NERVE. INFILTRATING SPINDLE CELLS CONTAINING PIGMENT, SUPPORTED BY TISSUE OF THE PART.

FIG. 9.—TUMOR IN POSTERIOR PART OF CILIARY BODY.

FIG. 10, CASE IV.—VARIATIONS IN SIZE AND SHAPE OF CELLS.

FIG. 11.—TUMOR CELLS WITH NET-WORK OF TISSUE CELLS INTERSPERSED.

MELANOBLASTOMA OF THE EYE.
FORMAN AND HUGGER.

scleral trephine operation for glaucoma has late results which may develop even years after the operation.

Second—That rarely if ever should it be the operation of choice.

Third—That it should be tried where iridectomy has failed.

Fourth—Some reports in cases of buphthalmos would seem to warrant further trial in this condition.

MELANOBLASTOMAS OF THE EYE.

JONATHAN FORMAN, M. D., AND C. CLARON HUGGER, M. D.

COLUMBUS, OHIO.

Studies from the Laboratory of Pathology of the Ohio State University. Supplemented by clinical notes regarding the four cases reported; and reasons for preferring the term melanoblastoma to the more commonly used sarcoma of the uveal tract.

Tumors of the choroid may be either primary or secondary in origin. Primary growths are the more frequent and the large majority of these are composed of pigment producing cells.

The pigment bearing choroidal tumors are usually referred to as melanosarcomas. In this region pigment-free tumors are sometimes observed which differ in no wise either in structure or clinical course from those which contain pigment. Such tumors have been collected by Schieck¹ under the term leucosarcoma. This would appear to be an unfortunate term, because it is used by Sternberg and others to designate those cases in which a definite tumor-like mass is developed in some organ or tissue; and is composed of lymphoid cells which seem to escape into the blood stream, giving rise to leukemic alterations of the blood. Ribbert designated both the pigmented and the pigment-free tumors of this group as *chromatophoromas*. Mallory adopted the more euphonious and equally comprehensive term of *melanoblastoma*. The use of a group name for all tumors composed of melanoblasts, or chromatophores, has the advantage of not placing any emphasis upon benign or malignant forms. This compels a more careful study and accurate evaluation of the individual tumor.

Marshall² in his collection of 134 cases of melanosarcoma of the eye, found that the age ranged from 15 to

90 years with an average age incidence of 50 years. One eye, as a rule, is affected. The cases on record appear to be about equally divided between the left and the right eye. Injury is considered as an important etiologic factor by some authors. Marshall found a definite history of injury in about ten per cent of his series. Among the fifty-five eye specimens in the Museum of Pathology at the Ohio State University, there are four cases of melanoblastomas arising in the eye.

CASES.

Case I. *Clinical Notes:* A white woman, aged 30, consulted an ophthalmologist on October 8, 1914, because of pain and loss of vision in the right eye. Upon the appearance of a dark area on the upper surface of the globe of this eye about three years before, the vision of the patient began to decline. This area increased in size, bulged slightly, and was attended by an excruciating pain throughout the entire period. The patient recalled rather indefinitely having received an injury to the right frontal region. Ophthalmoscopic examination of the right eye revealed a normal fundus reflex. The details of the fundus were imperfect. There was a marked cupping of the disc. This was verified by the indirect method.

Removal was advised and the eye was enucleated October 9th, 1914, under ether anesthesia. The patient made an