CLINICAL REPORTS.

TWO INTERESTING CASES.

BY JOHN B. KEBER, M.D.

Professor of Diseases of the Skin and Syphilis, Beaumont Hospital Medical College,
St. Louis.

Triple Specific Infection.

FIGURE 1. Triple Specific Infection.

CASE I.—M. B., printer, age 21, applied for treatment June 23rd, 1896. He stated that about four weeks previously he had noticed the lesions on the upper lip, penis, and left cheek as small papules. They appeared about the same time, gradually increased in size, became eroded and indurated, but gave rise to no pain and little inconvenience. He, therefore, hardly gave them a thought. An eruption had made its appearance on his body within the past few days, and for this he had applied for treatment. This eruption was readily recog-

nized as a maculo-papular syphilide. Date of exposure could not be determined. The interesting feature in the case consists in the triple infection, giving rise to three initial lesions in regions widely separated. *Practically*, the case offered no difficulties, as the complex of symptoms was complete; *theoretically*, however, it presents many points of interest, but as their discussion would hardly come within the scope of this journal, it will be reserved for a future paper. The illustration shows the salient features of the lesions on the lip and cheek, the lesion on the penis having been omitted.*

Initial Lesion of the Tongue.

FIGURE 2. Initial Lesion of the Tongue.

CASE II. is interesting from a practical standpoint, inasmuch as the patient presented himself so early after infection that the importance of the lesion might easily have been overlooked. He stated that about ten days before, in using his knife as a tooth-pick, he had slightly wounded the tip of the tongue. The cut healed promptly, but three

* THE LARYNGOSCOPE will, on application, send a full length illustration of this case to any subscriber remitting a two-cent postage stamp.

or four days before consulting me he noticed a small spot as large as a pinhead covered with what he thought was a white membrane. Attempts at its removal proved a failure, and, as the spot increased in size from day to day, and inconvenienced him some by striking against the teeth, he called upon me for treatment. Inspection showed a lesion on the tongue, about one-quarter of an inch from the tip and a little to the left of the median line. This lesion was about the size of a lentil, slightly elevated, pearly white in color, and surrounded by a dark red halo about one line in width. It was perfectly round (later oval), sharply circumscribed, and on superficial examination seemed nothing more or less than a membranous deposit on a slightly congested base.

Closer inspection showed, however, that this was a mistake, and that the lesion really consisted of a small papule covered by a thickened and opaque epithelium. Palpation revealed a slight increase in the consistency of the part. The neighboring lymphatic glands were apparently normal. As the result of a process of exclusion, a probable diagnosis of "initial lesion" was made, and the patient instructed to report daily. The lesion increased slowly from day to day, its general character remaining the same, and in four or five days had attained the size of a split pea. Induration was now easily recognized, and the submaxillary glands on the right side showed signs of involvement. The centre of the lesion became depressed, and as the lesion grew in extent, gradually became eroded. The photograph was taken at the end of the second week, and shows the fully developed lesion, as well as the enlargement of the glands. At the proper time a macular syphilide made its appearance. Constitutional treatment was at once begun, and resulted in the rapid disappearance of all symptoms.

Ethmoiditis.

Dr. J. F. Barnhill (*Ind. Med. Jour.*) says that in simple cases the treatment should consist in removing all obstructions to the free drainage of the cells, and the application of soothing washes or sprays. In chronic cases he advises that a portion of the middle turbinated bone be removed, and that all the suppurative cells be converted into one cavity, if possible, by means of the curette or drill. The subsequent treatment should consist of hot antiseptic douches. Vigorous constitutional treatment will be needed.

By following this line of treatment all will be improved and many cured, but the patience of both surgeon and patient may be severely taxed.