3. The best means of accomplishing the last-named object is a few buried sutures in the apex of the axilla and a properly adjusted compressive dressing held in place by adhesive plaster.

DISCUSSION.

DR. WILLIAM L. RODMAN, Philadelphia, agreed with Dr. Porter as to the necessity of the free removal of the overlying skin, even in such cases as show that we can not get primary union. He inclined to the belief that possibly the greatest mistake that is made in practice generally is to subordinate the probabilities of a radical cure to securing primary union in every case. He said that we should keep constantly before us that Gross and Pancoast and the older surgeons reached 10 and 15 per cent. of cures, but only by very freely removing the skin. A complete axillary dissection was never done, and muscles were not destroyed. Whatever good was in these operations must have been in the free removal of skin. Therefore, he thinks that we should not accentuate the necessity for getting primary union, even in a majority of the cases. It is all very well if it can be done and without making too much tension, and this, he said, can be done, as Dr. Porter says, by undermining the skin, or by skin grafting, which can always be resorted to to cover in any defect that may exist.

Dr. Rodman has done the operation according to the method described by Dr. Porter a number of times, and he has also advocated strongly the operation of Warren, which is practically the same. He liked it very much and thought it the best operation he knew, inasmuch as it has generally made primary union possible, but he has found in a number of cases a bridle of skin which fetters and prevents the uplifting of the arm beyond that extent, thus interfering with the doing up of the back hair, etc. While Dr. Rodman still believes that that operation is one of distinct value, he has obtained recently a better result from the operation of Jackson, and prefers to place the incision in that way. He believes that it is a mistake to cut on to the arm, or, at least, to go low down on the arm.

Dr. Rodman emphasized the necessity of removing the muscles in every case of cancer of the breast, although he was not prepared to say that the muscles will be involved. Researches have demonstrated that the fascia of the muscles usually will stop limitations of the growth, but recent anatomic discoveries show that Halsted built better than he knew when he insisted on the removal of the muscles in every case. For, in the first place, a complete axillary dissection can not be made with the muscles *in situ*, and Gross and Pancoast demonstrated that occasionally there will be lymph glands which can not be felt with the muscles in position. Therefore, it is necessary to remove the muscles in every case, otherwise the operation is incomplete.

As to drainage, Dr. Rodman admitted that in the majority of instances the wound can be closed without drainage, and yet it certainly is the custom of American surgeons to drain after amputations of the breast. In preparing a paper on this subject some time ago, he wrote to fifty American surgeons, and of the twenty-one who replied, there were three who did not use drainage. Certainly drainage is not necessary in case of a small breast particularly, but Dr. Rodman believes that, as a rule, it is just as well to drain.

Speaking of the necessity of invading the neck, Dr. Rodman stated that he is satisfied that we do not invade the neck as often as we should. Recent anatomic investigations show that if the tumor is located superiorly, in the upper hemisphere of the breast, it is drained by the lymphatics which pass from the clavicle into the glands above the clavicle and not directly into the axilla. Therefore, in the case of tumors that are situated superiorly, it is unnecessary to enter the neck in every case.

DR. MILES F. PORTER, Ft. Wayne, Ind., was very glad indeed that Dr. Rodman emphasized the necessity of not sacrificing the chance of a perfect recovery for the sake of securing primary union. In the case he reported the breasts were removed as widely as any one removes them, and yet, in the majority of cases, union was achieved. Dr. Porter invades the neck, when it is the proper thing to do. As to drainage, he said that it is never too late to mend. The sooner those who make it a rule to drain, make it a rule not to drain, the better.

PLASTIC RADIOGRAPHY.

A PRELIMINARY COMMUNICATION.

WALTER G. STERN, M.D., AND E. ROSENBERG, M.D.

CLEVELAND, OHIO.

Radiographs, usually spoken of as skiagraphs, are not, as the latter name would imply, mere shadow-graphs or silhouettes; they are, rather, the result of the superimposition of semishadows or half-tones as they are called in photography, whose size, shape and intensity depend on the size, shape and molecular density of the object to be radiographed. But such pictures have never been interpreted as perspectives or "plastic" reproductions, and successful attempts have never been made to obtain "relief" effects except by means of stereoscopic reproductions.

Fig. 1.—A "plastic" radiogram, showing distinct shadow of soft parts, with high-lights and shadows of the bones, giving relief effects. Print on bromid paper direct from the combination of the negative and positive.

Béla Alexander, a country practitioner hitherto unknown, living in an obscure part of Hungary, first published¹ a method of obtaining "plastic x-ray" photographs. It is through the efforts of one of us (Rosenberg) that the substance of this and other contributions of Alexander and other Hungarian writers in the various Hungarian medical journals has been collected and abstracted. The method as they give it is long and tedious, full of archaisms and omissions of most important details such as the time and methods of exposure, copying, registering, printing, etc. We have tried

1. Orvosegyesület, March 10, 1906.

many methods, always rejecting anything which could smack of "dodging" (as the professional photographer would say) and are now ready to report the following provisional modification of Alexander's method:²

A careful exposure is made with two tubes on a specially prepared double coated x-ray plate. This plate is coated so that the slower emulsion is on top, and was made for this process, together with the plate for the positive (No. 2) by the Lumiere Company.

The first tube used for the double exposure is a very soft one, and without changing in the least the relations between the tube and the object and the object and the plate a second exposure is given with a hard tube. This change of tubes is easiest done by using a compression diaphragm apparatus of the Albers-Schoenberg or Frei-

ones the relative exposure will be about as 5 to 1. That is, if with a certain current one would ordinarily expose, say, a hand twelve seconds, then for "plastic" work the first exposure should be ten seconds and the second two. Of course no hard and fast rules can be drawn.

The plate is then developed in the usual manner and must be quite dense and contrasty; if the exposures have been correctly timed "plastic" effects will already be observed in this plate, which is, of course, the negative (No. 1). After the negative has been dried, a positive (No. 2) is to be made by direct contact in a printing frame, using a slow plate on the order of a lantern slide plate. The positive must be equal in density and contrast to the negative.

The plates are now placed back to back and accurately registered. This is best done by scratching on the negative before making the positive a short fine vertical line in the center of the upper edge of the plate and two short fine horizontal lines, one each in the lower right and left-hand corners. These fine scratches on the negative will, of course, be reproduced on the positive by corresponding fine black lines. If the plates thus

Fig. 2.—Ordinary radiogram of same hand as shown in Figure 1, made with a moderately soft tube; exposure 12 seconds. Note flatness and weak shadow of soft parts.

berg type, in which latter the tube holder is a separate piece and has two sights for centering the tube. My own apparatus has a shallow wooden cup accurately centered over the diaphragm for receiving the tube; thus in using tubes of the same size the target will always fall approximately in the same place.³ Neither the object nor the plate dare be disturbed in the least during this change. Although the European writers claim to have had great difficulty in determining the relative lengths of the exposure to the various tubes, we have found that for very soft tubes followed by medium hard

Fig. 3.—Plastic radiograph of a fractured elbow showing stereoscopic projection of the internal condyle. Picture was taken with the internal condyle and ulna nearest the plate. The picture shows best if held three feet away and two feet below the level of the eyes.

marked are placed back to back and held together by light spring clamps and examined in a strong light sending out parallel rays—an automobile gas lamp, for instance—they can easily be moved one over the other until these fine lines are found to overlap accurately. The plates are now properly centered and are permanently bound together with passe partout binding paper.

A print must now be made with the paper in contact with the positive plate, and for this purpose we advise the use of bromid paper, as it alone has the requisite speed to take the place of the Plates 3 and 4 of Alexander's method. The printing of No. 3, the "plastic," can only be done with parallel rays of light which must strike the plates at an oblique angle. Ordinary light is, of course, out of the question on account of non-parallelism of the rays, dispersion and reflection. For this purpose we use the aforementioned automobile gas lamp at a distance of from ten to twenty feet. The plate holder stands at an angle of about 45 degrees to the plane of the lamp, thus giving very oblique illumination. The distance from the light to the plate does away with any

2. Alexander makes use of four plates before his final print can be obtained. We have shortened the method and use only two.

3. By using a Mueller tube No. 16, for hard and soft rays, one can obviate the use of two different tubes, as this tube No. 16 is constructed in such a manner that by changing the connecting wires to different cathodes one can get soft or hard rays at will.

possible unequal illumination, for the nearer edge of the plate necessarily receives more light than the more distant one. The "plastic" illustrating this article (Fig. 1) was exposed fifteen minutes on bromid paper.

When developed the print (No. 3) will show beautiful "plastic" effects and the soft parts usually stand out in wonderful clearness. The more perfect the first negative—Plate No. 1—the better the "plastic."

How does this "plastic" result from the double exposure and the combination of the negatives? We do not know. According to the law and the prophets the combination of an equally dense and perfectly centered negative and positive should result in an absolute blank, as the opacities in the one and the transparent parts of the other should counterbalance.

How does the "plastic" differ from the ordinary skiagraph? As previously stated, the "plastic" does not consist merely of the overlapping of half-tones of different intensity, but contains "shadows and high lights"—the two light values which make an ordinary photograph or drawing stand out in relief.

Are the effects mere photograph dodges? So the skeptics have claimed, and at first sight we were inclined to the belief that they were mere artifacts. But one by one the skeptics have been silenced and have joined the ranks of the believers until even Schellenberg, the earliest and firmest opponent, has been converted and forced to admit the "plastic" radiography to the greatest advance since Roentgen.

As for the theories and rationale of "shadowgraphs and plastics" and the explanation of the various phenomena involved in their production, we are not yet able to speak from conviction. It has taken most of our time to perfect the details, and we shall give at another time and in a more suitable place an explanation of the various principles involved.

A SIMPLE AND SAFE OPERATION ON THE FRONTAL SINUS BY THE INTRA-NASAL ROUTE.

R. H. GOOD, M.S., M.D.
CHICAGO.

The instruments used in this operation are two frontal sinus rasps, two tapering curved chisels and three protectors.

The frontal sinus rasp (Fig. 1) is made on the same

Fig. 1.—Frontal sinus rasp, showing in outline exact length and width of rasp.

Fig. 2.—Chisel, showing in outline exact size.

principle as the old-fashioned coarse horseshoe rasp, from which I derived the idea. It is slightly curved at the rasp end to fit into the frontal sinus and the teeth are on the longitudinally concave surface, pointing toward the handle, so that when traction is made on the rasp, the bone anterior to it is rasped away, as illustrated in Figure 5. The posterior surface of the rasp is perfectly smooth and flat.

The chisels (Fig. 2) are short, curved and tapering. The handle at the chisel end is very slender so that the chisel can be manipulated under the direct vision of the eye. The edges are tapering as well as blunt, so that if the edge should strike a bony surface it would tend to glide off instead of penetrating.

The protectors (Fig. 3) are made of brass and in three different widths, as illustrated. They can be bent easily into shape on the flat, but not sideways.

I have performed these operations under local anesthesia. The cocain crystals are rubbed into the mucous membrane with a cotton swab on an applicator

Fig. 3.—Frontal sinus protectors.

which is first moistened with adrenalin chlorid 1 to 1,000, and then dipped into the cocain crystals. The entire olfactory region as well as the bulla ethmoidalis and unciform plate are thus anesthetized. The middle turbinate is then removed by the ordinary methods and the processus uncinatus of the ethmoid bone is removed with the larger chisel.

A frontal sinus catheter, with syringe attachment, is then inserted into the frontal sinus and a half dram of a 10 per cent. solution of cocain in adrenalin 1 to 1,000 is injected into the sinus. A pledget of cotton is placed over the infundibulum for a few minutes to take up the cocain as it returns from the sinus. The largest possible protector is now inserted into the frontal sinus, and if the smallest protector will not enter, which is rare, a good-sized probe is used, until later in the operation when the protectors can be inserted.

Having the protector or probe, as the case may be, *in situ*, the larger chisel is placed in position as illustrated in Figure 4, and a part of the processus frontalis of the superior maxillary bone, as well as a part of the spina frontalis of the frontal bone, and the anterior medial wall of the labyrinth ethmoidalis are chiseled through and removed with forceps or curettes.

The chisel must always be kept in direct line with the protector so that the tabula interna is not endangered. There is a considerable space between the spina frontalis and the tabula interna of the frontal bone, and the operator can almost always chisel without touching the protector, for the moment the chisel has chipped off a piece of bone it has to pass through a space before it encounters the protector, and the hand nearly always will stop the chisel before it touches the protector; and in case it does strike the protector, the hand has reduced the force to such an extent as to make hardly a mark on the protector.

As stated above, the edges of the chisel are dull and tapering, so that if the edge should touch the lamina papyracea of the ethmoid bone or the tabula interna of the frontal bone, it will slide off and by the time the chisel has traveled through the space of about half an inch, under proper care, would not have enough force to penetrate even if the edges were sharp.

The anterior ethmoidal cells are removed with forceps or curette, and by this time the frontal sinus rasp may