THE MATHEMATICS NEEDED IN FRESHMAN CHEMISTRY.

BY LEWIS W. WILLIAMS,

University of Illinois, Urbana.

This study was made to throw some light upon the criticism that high school students entering college with a year of algebra and a year of geometry do not have sufficient mathematics to handle Freshman chemistry. With this practical purpose in mind the book selected for examination was the text, "A Text-book in Chemistry," by Professor W. A. Noyes, required in the Freshman chemistry course in the University of Illinois.

Because the text consists of two distinct phases—exposition of the principles of chemistry and problems given for solution—the study is divided into two parts and these are treated independently. In the analysis of the expository portion a careful examination of the text was made to determine what words or expressions were used which were distinctly mathematical or which implied mathematics. All others were omitted. These were then classified and their frequency determined. In the second part of the study the problems were solved and where more than one solution was possible, the simplest was selected. The operations and quantities were classified as in the first part and the frequency determined.

It was early seen that some standards had to be set up for determining whether or not a word or expression should be included in the study or be omitted. The following principles were chosen for this purpose:

1. Expressions giving dates, either specific or general, imply mathematics. Example 1744, last century.
2. Denominate numbers are distinctly mathematics. Example 5° F., 76 cm.
3. Expressions having several meanings or uses are distinctly mathematical if, from the context, a mathematical meaning is evident; if not, they are omitted. Example, equivalent, constant.
4. Expressions having no use or meaning other than in a mathematical sense are distinctly mathematical. Example, volume, number.
5. Expressions not classed under Principle 4, but involving a use of mathematics, are considered as implying mathematics. Example, rate of expansion, equilibrium, divisions on a scale.
6. Cardinal, ordinal and numeral adjectives are distinctly mathematical. Example, three, first.
7. Equations, formulas, symbols, etc., are distinctly mathematical. Example, $AB = 180$, $F - 32 = C$, $C = R$

180	100	80
-----	-----	----

Words or expressions which could not be classed under the above principles were omitted.

Exposition.

Table I shows the mathematical concepts, a reading knowledge of which is necessary for an understanding of the expository portion of the text. It is obvious from an examination of the list that the concepts most commonly used are those most commonly found in arithmetic or in the first year high school algebra, or geometry. And those less frequently used are either found in the same sources or are not found in second year algebra. As a matter of fact, it would appear that if the less familiar terms, as rhombic hexahedron, are to be taught as a prerequisite to chemistry, it would be less serviceable to teach advanced mathematics than to give specific instruction in the meaning of this list.

Table II indicates the use of many common terms, as cents, miles, years, and two other groups of terms. These are the metric system, (32 terms), and certain terms in physics, as

TABLE I.
Mathematical Concepts Used in Exposition.

add.....	2	formulas.....	1
added.....	2	fraction.....	16
absolute zero.....	3	fractions.....	2
abscissa.....	3	geometrical.....	1
adding algebraically.....	1	hemispherical.....	1
algebraic sum.....	1	hemihedral.....	1
amounts.....	1	hexagonal.....	1
angle.....	4	hexagonal pyramid.....	1
angles.....	1	hyperbola.....	1
apex.....	1	invariable.....	1
average.....	8	inversely.....	4
axis.....	13	inverse proportion.....	1
British Thermal Units.....	3	inversely proportional.....	1
calculate.....	1	last.....	28
centimeter.....	1	length.....	5
constant proportion.....	6	lengths.....	4
constant volume.....	1	line.....	2
constant.....	1	lines.....	4
cube.....	4	monoclinic.....	2
cubes.....	2	multiple.....	2
curve.....	1	multiples.....	2
curves.....	4	multiplied.....	2
cross section.....	1	multiplying.....	1
denominator.....	2	multiple proportion.....	3
diameter.....	8	negative.....	4
difference.....	5	number.....	64
direct proportion.....	1	numbers.....	4
divided.....	5	numerator.....	5
division.....	1	octohedra.....	6
dividing.....	1	octohedron.....	2
empirical.....	1	origin.....	1
equation.....	43	ordinates.....	3
equations.....	14	parallel.....	6
face.....	2	per cent.....	310
faces.....	2	percentage.....	8

perpendicular.....	1	six-sided.....	1
plane.....	5	square.....	3
plotted.....	1	square prism.....	1
positive.....	5	square root.....	3
prism.....	1	straight line.....	1
product.....	4	subdivide.....	1
proportion.....	51	subdivisions.....	1
proportional.....	14	submultiple.....	1
proportionably.....	1	subtract.....	1
pyramid.....	1	subtracted.....	2
pyramids.....	2	subtracting.....	2
rate.....	29	sum.....	5
ratio.....	24	surface.....	1
ratios.....	3	tetragonal.....	1
reciprocal.....	3	tetragonal pyramid.....	1
rectangular.....	1	tetrahedron.....	2
rectangular prism.....	1	triclinic.....	1
rhombic.....	5	triple.....	1
rhombic dodecahedron.....	1	unit.....	3
rhombic hexahedron.....	1	unity.....	1
rhombic prism.....	2	varies.....	3
rhombohedra.....	1	vary directly.....	3
right angles.....	4	vary inversely.....	2
right angled prism.....	1	velocity.....	34
series.....	12	volume.....	208
single.....	1	volumes.....	41
simple ratio.....	6	zero.....	1

TABLE II.

Denominate Numbers Used in Exposition.

amperes.....	4	kilogram.....	21
calorie.....	29	kilograms.....	5
centimeter.....	74	kg.....	1
cento.....	3	kilometer.....	1
cents.....	3	kilometers.....	2
century.....	2	kilos.....	2
centuries.....	5	kilowatt.....	1
c. c.....	41	kilogram meters.....	5
cubic centimeter.....	22	liter.....	60
cubic decimeter.....	1	liters.....	45
cubic feet.....	6	meter.....	10
cubic millimeter.....	2	meters.....	12
day.....	2	metric ton.....	1
days.....	5	micron.....	1
decimeter.....	1	mile.....	1
degree.....	8	miles.....	1
°.....	865	milligram.....	22
\$.....	8	milligrams.....	29
erg.....	5	milliliter.....	1
ergs.....	1	mm.....	104
feet.....	4	millimeter.....	3
footpounds.....	3	minute.....	14
gallon.....	1	minutes.....	3
gram.....	169	mouths.....	5
grams.....	172	ohm.....	2
g.....	12	ohms.....	1
half dozen.....	1	ounces.....	4
hours.....	7	pound.....	12
horsepower.....	2	pounds.....	4
inch.....	2	lb.....	1
joule.....	5	second.....	1
joules.....	7	square centimeter.....	2

sqcm.....	2	watt.....	4
square meter.....	1	weeks.....	5
ton.....	13	year.....	3
tons.....	11	years.....	45
volt.....	4		

TABLE III.
Other Terms Used in Exposition.

first.....	115	twice.....	20
second.....	54	double.....	4
1st.....	1	doubled.....	2
2nd.....	1	triple.....	3
once.....	1		

TABLE IV.
Common Fractions—in Figures and Words.

<i>Common Fractions—in Figures</i>		<i>and Common Fractions.</i>	
1/2.....	12	1/15.....	1
one half.....	35	one fourteenth.....	1
one & one half.....	2	1/20.....	1
two & one half.....	2	one twentieth.....	2
3 1/2.....	1	71/29.....	1
fourteen & one half.....	1	hundredths.....	6
third.....	23	three one-hundredths.....	1
two thirds.....	6	1/120.....	1
fourth.....	17	one six-hundredth.....	1
three fourths.....	6	thousandth.....	3
3/4.....	1	1/1000.....	4
20/4.....	1	one seventeenth-hundredth.....	1
one & one fourth.....	1	one millionth.....	3
one fifth.....	32	ten millionth.....	1
two fifths.....	1	1/3000000.....	1
three fifths.....	1	1/7000.....	1
four fifths.....	6	1/500000.....	1
3/5.....	1	1/5000.....	1
sixth.....	8	1/272.....	3
five sixths.....	1	1/272.8.....	1
seventh.....	7	1/272.9.....	1
one eighth.....	1	1/272.6.....	3
seven eighths.....	1	1/273.....	2
ninth.....	2	1/3.8021.....	1
5/9.....	1	4/3.1416.....	1
tenth.....	23	18/24.....	1
tenths.....	2	22/24.....	1
1/10.....	2		

amperes, calories, and watts (13 terms). Here, again, it is apparent that advanced mathematics is less essential than an intensive drill upon the meanings of the common concepts and upon the metric system, together with specific explanation of the thirteen physical terms, in case physics has not been studied previously.

Table III presents a list of very simple terms quite understandable by even those who have not studied arithmetic.

The significance of Table IV is apparent. The student needs a reading knowledge of fractions of all-sized denominators, and, although fractions with denominators larger than ten constitute only eighteen per cent of the total, they are of sufficient importance to require attention. They involve, also, a knowledge of

decimals, though not so extensive a knowledge as is required in later tables. Here, again, it is obvious that an intensive review of fractions would be more efficient than an advanced course in mathematics where the more difficult of these terms, which presumably give more trouble, would be encountered by chance, if at all.

TABLE V.
Decimal Fractions Used in Exposition.

	Fraction	Frequency
.1	.9	33
.01	.99	140
.001	.998	50
.102	.996	75
.0002	.9996	49
.00002	.99999	66
.000008	.999982	14
.0000001	.0000026	6
.00000326		1

Table V clearly shows that a student needs to understand decimal notation to quite extended limits, although, as might be supposed, the lower places are most frequently used; four-place decimals, or less than four, constitute eighty-two per cent of the cases.

Table VI indicates that there were found 121 cases of mixed decimals with one place in the integral portion of the number and one in the fractional portion; as 4.3; and that 256 cases of the type 27.3, with two integral and one decimal place, were obtained, and so forth.

Table V contains the pure decimal fractions while Table VI contains the mixed decimals. The contrast in the use of common fractions and decimals is very evident when we note that fifty different common fractions were used and 920 different decimals. Were the frequency of the different fractions considered, this contrast would be much more emphatic. Pure decimals range from one place to eight places, the large majority being two, three, four and five places. Among the mixed decimals we find the following limits: From one unit place and one decimal place to four unit places and one decimal place, by far the greater number consisting of two or three unit places and one decimal place; from one unit place and two decimal places to three unit places to three unit places and two decimal places, most of them consisting of one or two unit places and two decimal places; from one unit place and three decimal places to three unit places and three decimal places, most of them consisting of two or three unit places and three decimal places. There were very few mixed decimals with four or five

decimal places. None were found with a higher number of decimal places than five.

TABLE VI.
Mixed Decimals Used in Exposition.

Place Integers 1	Place Decimals 1	Frequency 121
2	1	256
3	1	154
4	1	1
1	2	173
2	2	186
3	2	44
1	3	34
2	3	3
3	3	2
1	4	41
2	4	1
1	5	3
2	5	1

In Table VII, we find the integers used in the first part of the study. One place integers to the number of 570 were found in figures and 975 in words. They range from one unit place to twenty. There are very few beyond six unit places, however. Furthermore, most of the numbers consisting of more than four unit places end in one or more ciphers, which naturally simplifies their use.

Problems.

In Part II, all problems given for solution were solved. Each solution was then analyzed into its processes. In Tables VIII and IX, we have the processes and their frequency as found in the sixty-six problems. The striking thing about this table is the low frequency of equations, not only for two unknowns but also for one. This is in part explained by the fact that in a typical chemical problem, such as—How many grams of magnesium should be mixed with three grams of sand to form silicon?—the operation is not considered as algebraic but rather as the solution of an arithmetical proportion in which three terms are given to find the fourth.

2 Mg plus Si O₂ 2 MgO plus Si

48.6 60.3 96.6

60.3 : 3 :: 48.6 : ? $48.6 \cdot 3 = 2.42 \text{ Ans.}$

60.3

Certainly no algebra is needed for the solution of such problems. This brings us to the use of ratio and proportion.

TABLE VII.
Integers Used.

Place	Figures	Frequency	Words	Total
1	570		975	1545
2	1150		39	1189
3	893		13	902
4	382		9	391
5	40		0	40
6	25		6	31
7	2		4	6
8	9		0	9
9	1		1	2
10	2		1	3
14	0		1	1
20	1		0	1

TABLE VIII.
Processes Used.

Equation 1st Degree, 1 unknown.....	7
Equation 1st Degree, 2 unknowns.....	2
Equation 1st Degree, 3 unknowns.....	0
Equation 2nd Degree, 1 unknown.....	0
Equation 2nd Degree, 2 unknowns.....	0
Arithmetical addition (integers).....	72
Arithmetical subtraction (integers).....	11
Arithmetical multiplication (integers).....	149
Arithmetical division (integers).....	25
Arithmetical addition (with decimals).....	41
Arithmetical subtraction (with decimals).....	1
Arithmetical multiplication (with decimals).....	118
Arithmetical division (with decimals).....	88
Arithmetical addition (common fractions).....	0
Arithmetical subtraction (common fractions).....	0
Arithmetical multiplication (common fractions).....	9
Arithmetical division (common fractions).....	2
Fractional equations.....	6
Chemical equations.....	134
Percentage.....	29
Denominate Numbers.....	173
Substitution in Formula.....	9
Ratio and Proportion.....	77
Algebraic addition.....	2
Algebraic subtraction.....	0
Algebraic multiplication.....	0
Algebraic division.....	0
Highest Common Divisor.....	1
Clearing equations of fractions.....	6

TABLE IX.

Denominate Number Units Used in Problems.

calories.....	12	cubic feet.....	5
c. c.....	10	cubic meter.....	2
cm.....	2	degrees.....	17

gallon.....	1	metric ton.....	1
grams.....	53	milligrams.....	5
hour.....	2	minute.....	1
kilograms.....	9	pound.....	5
liters.....	32	square centimeter.....	2
mm.....	9	ton.....	3
meters.....	2		

There were fifty-nine cases of the type just indicated. There were eighteen cases involving simple ratios. It is very evident that the principles upon which ratio and proportion rest should be thoroughly understood and be thoroughly usable by the pupil.

Fractional equations were used very little. All of the six cases found might be considered as cases of compound proportion. However, the solution in each case involved substitution in a formula and use of the equality sign, so it was deemed best to list them as above. It is interesting to note, however, that in all of the cases, the denominators were whole numbers of not more than three unit places.

There were twenty-nine cases of percentage found among the problems. Without exception these were simple applications of the principles of percentage. Nine were cases of base and rate given to find percentage; eight of percentage and rate to find base; and twelve of percentage and base to find rate.

As was to be expected, there was found a wide use of denominate numbers. Fully 90 per cent of the problems involved denominate numbers in some form or other. Nineteen different units were used—with frequencies as indicated in Table IX. Usually, the problem was stated in units of some kind and their use was limited to manipulation merely and did not involve changes from higher to lower terms or from one system to another. This was true in sixty-three cases; in nine cases, a change from a higher to a lower denominator was required; in four, from a lower to a higher; in three, a determination of volume was required; in two, a change from the English system to the Metric and in two more, from the Metric system. Frequently, where changes were required from English to Metric or the reverse, the equivalent values were given directly.

There is very little use made of addition, subtraction, multiplication and division in algebra. These processes are used quite extensively, however, in an arithmetical way. This is particularly true of multiplication. Its frequency, both for its use with integers alone and with decimals, is large. Division and addition are used much less frequently.

The range of additions is indicated by the following facts: In sixty cases, additions were made of integers of 7 places or less to another integer of 7 places or less. In 30 cases, three integers of 6 places or less were added while in one case four numbers with 5 places of digits were added. Apart from these, additions of integers and decimals were found. In all, forty-two cases appear. Decimals of more than four places do not appear. The most complicated addition is that of three numbers consisting of one place digit and three decimals (as, 8.456), two place digits and three decimals, and three decimals. In only four cases was a decimal added to a decimal; in the remaining cases a decimal is added to a whole number.

The range of subtractions is very narrow. In the twenty-one cases reported, the most complicated is the subtraction of a six place integer from another of the same kind. The most common (eleven cases) consisted of the subtraction of a one place from a two place integer.

In the 235 cases of multiplication, 93 consisted of the multiplication of two place integers by a one place. In 156 cases, the operation involved nothing more complicated than a four place integer by a six place (of which there was one case). Three integers were multiplied only three times and no cases occurred of the multiplication of more than three. In the matter of decimals sixty-three cases were found in which a decimal or mixed decimal is multiplied by an integer. Of these, the decimal is never more extended than four places and the mixed decimal than five places in the integral portion and three places in the decimal portion. In sixteen cases a mixed decimal is multiplied by a mixed decimal, the most difficult being the multiplication of a number composed of two places in the integral portion and three places in the decimal by another composed of three places in the integral portion and two places in the decimal.

In the 139 cases of division, forty-six consisted of the division of one integer by another. These varied from one to twelve place integers. There was but one twelve place integer, one nine place integer and two eight place integers. Most of them were one, two or three place integers. There were twenty-five cases where a decimal was divided by an integer. In all but one case, the decimal was mixed. The most complicated of these were composed of eight places in the integral portion and one place in the decimal (1 case); 2 places in the integral portion and 4 places in the decimal (1 case); and 2 places in the integral portion

and 3 places in the decimal (1 case). There were twenty-six cases where an integer was divided by a decimal. In all but four of these the decimal was mixed, the most complex being composed of 3 unit places and 2 decimal places as 456.27. The numerator in most of these cases was not larger than a 5 place integer; in 2 cases, a 6 place integer. There were forty-two cases where a decimal was divided by a decimal, and in twenty-eight of these, both numerator and denominator were mixed decimals. A typical case is here given—34.598 divided by 27.4. The most complicated case is that of a number composed of 6 unit places and 2 decimal places divided by a number composed of 3 unit places and 1 decimal place.

Three important things remain for consideration—chemical symbols, chemical formulas and chemical equations. All have naturally a very extensive use and consequent emphasis upon the mathematics involved. With these is bound up the question of valence as well as that of atomic and molecular weight. A symbol stands for the name of an element, for one atom of that element and for one gram-atom of that element. More than one atom is indicated by a number placed at the right and a little below the symbol. A formula stands for the name of a compound, for one molecule of that compound, for one gram-molecule or for one gram-molecular volume, if a gas. Two or more molecules are represented by a number placed before the formula. An equation gives the substances which react with each other and the products formed, the number of grams of each substance or if the substances are gases, the number of gram-molecular volumes.

The mathematics used here is relatively very simple. Representing more than one atom or more than one molecule involves the simple principle of algebraic coefficients, the location of the figure in the case of the symbol being a mere matter of convenience. With atomic weight, the problem is very similar. Here the numbers are generally not simple but larger and often mixed decimals, as 35.46 for chlorine. The molecular weight is obtained by adding the atomic weights of the elements in the formula (or the multiples) of the atomic weights.

Chemical equations have some things in common with algebraic equations but they differ materially in several respects. We will concern ourselves with the mathematical phases only. The use of the equality sign, transfer of terms from one side of the equation to the other and general handling of the equation,

all conform to mathematical principles. Very few cases were found, however, where transfer of terms, simplification of the equation or general manipulation was used. Balancing a chemical equation is, however, a distinct process. It involves a simple numerical relation, keeping the number of atoms of each kind on one side of the equation the same as the number on the other. The success of this, of course, depends upon knowing the products of the reaction and the use of coefficients in expressing the formulas of products and their quantities. Analysis, therefore, shows that the mathematics used consists of ratios of simple numbers, multiplication of mixed decimals by simple numbers, addition of these products, algebraic coefficients and simple manipulation of equations. Nearly 1,000 different symbols and formulas were used and about 500 equations.

Valence is the property possessed by elements or radicals of combining with or replacing other elements or radicals in definite and constant proportion. The degree of this property, usually indicated by the number of monad elements, represented by hydrogen, with which an atom or radical can combine, is the meaning which concerns us in this study. Thus, valence is a simple, numerical relation and involves nothing beyond simple cases of ratio or proportion. A careful study of the formulas found in the text shows that the following are typical of the ones used almost entirely— HCl , Na_2SO_4 , CaCl_2 , H_3PO_4 , AlCl_3 , H_4SiO_4 , Al_2S_3 , CaSO_4 , $\text{Ca}_3(\text{PO}_4)_2$. The simple ratios involved here are 1:1, 1 and 1:2, 2:1 and 1, 1 and 1 and 1:3, 3:1 and 1 and 1, 1 and 1 and 1 and 1:4, 3 and 3:2 and 2 and 2, 2:2, 2 and 2 and 2:3 and 3. Only in the organic compounds do we find any noticeable departure from such simple ratios as are here indicated. Even then, whether it be a question of stationary valence or multiple bond, the basis is still simple, as, for example, in the structural formulas for C_2H_4 .

Carbon having a valence of 4, has 4 arms or bonds for each atom. These must be satisfied for each atom, by union with bonds of other elements or with bonds of other carbon atoms. Even in the second case above, that of C_2H_4 , the double bonds

required to satisfy conditions, do not require anything but simple ratios.

CONCLUSIONS.

1. Summarizing, we may say that the study shows a rather limited use of mathematics in a first-year course of chemistry based upon the text examined—limited in the sense of the number of processes used and in the relatively simple nature of most of those processes. Algebra and geometry have a very limited use, though it is possible to use algebra occasionally when arithmetic might not be quite as direct.

2. A good working knowledge of the important principles of the following processes is desirable: addition, subtraction, multiplication, division, fractions, both common and decimal, graphs of simple curves and proportions, percentage, surface measure, volume measure, metric system, simple progression (used but slightly), ratio and proportion, and square root.

3. For solving the problems given, a thorough knowledge of the fundamental processes of arithmetic is necessary. This should hold, no matter whether integers, decimal fractions or common fractions are used. Simple equations of one unknown and simple fractional equations should be mastered. The pupil should have thorough training in percentage, denominate numbers and ratio and proportion.

4. A knowledge of the fundamental processes in algebra seems desirable, though by no means as essential as might be inferred. Aside from their use with algebraic equations, themselves seldom found, they are scarcely ever used.

5. It appears entirely feasible to construct upon the basis of the facts here displayed a simple set of exercises or preferably practice tests for chemistry students who would be required to attain efficiency in them outside of the class. They might even be prepared and distributed to high school students expecting to take Freshman chemistry in institutions using this text. It appears very clear that such a device would give a much better hold upon preparatory mathematics than would an additional year of algebra, trigonometry or solid geometry.

CLASS ROOM SAYINGS.

Where does the food go, after leaving one's stomach? It passes into the testaments.

Rain is formed by the condensation of the humidity in the air.

People should be very careful not to get humane (ptomaine) poisoning.