accompanied by respirations that were shallow and irregular in both force and rhythm. As soon as the animal had adjusted itself to the new condition, the respirations were slowed and were followed by a period of rapid inspiration and prolonged expiration, probably due to the pressure of the stomach on the diaphragm. The blood-pressure remained high throughout the experiment. In another instance with the insufflation method, the reaction was by no means so marked so far as the blood-pressure was concerned, but there was a definite temporary embarrassment of the respiration from which the animal relieved itself by belching.

The reactions of the medullary centers to operative traumatism as shown in these experiments suggest that we should be cautious in the treatment of the parietal pleura, particularly in tearing it at the angles of the intercostal wound by injudicious application of the rib-spreader. The lungs may be handled freely, but manipulations that tend to transmit traction to the great vessels and bronchi at the root of the lung should be reduced to a minimum. In heart suture, the Sauerbruch method of temporary hemostasis leads to too serious a fall in blood-pressure to be safe except where other means of hemostasis fail. Any mechanical stimulation of the heart, either directly or through the pericardium, during suture should be avoided so far as possible. Furthermore, in packing off the lungs to obtain an exposure of other thoracic viscera, it would be wise to avoid any unnecessary trauma which might tend to reduce the blood-pressure excessively. These are the stimuli which, in the present set of experiments, have produced the most serious reactions

310 Temple Street.

ACUTE POLIOMYELITIS, TRANSVERSE MYELITIS TYPE *

A. L. SKOOG, M.D.

KANSAS CITY

We have given names to diseases according to the clinical phenomena exhibited, according to etiologic factors and according to anatomic localization or changes. The title of this paper might be altered, have additional descriptive terms appended, or even shortened.

The object of preparing this paper is to place on record a case studied clinically and anatomically, a case of which I can find no counterpart after having searched diligently through the European and American literature.

This report covers one of the most interesting of the many cases which I, in the capacity of special investigator for the Kansas State Board of Health during the past three seasons, have had the opportunity of studying. The patient was taken by the family physician, Dr. Beach, of Clyde, Kan., on the sixth day of the illness to the University of Kansas Hospital at Rosedale where she was under my observation continuously until death, a period of three months.

History.—Patient, F. G., schoolgirl, white, aged 13. Father and mother are in good health. One sister, aged 15, well. No brothers or sisters dead. Patient's birth, infancy and early childhood normal. Measles at 5. No serious illness at any

time. Somewhat emotional and considered an affectionate child.

Patient was in excellent health prior to date of onset. On Nov. 2, 1909, she complained of pain in left lumbar and pleural region. On the following day the pain was general in the back, especially along the spine, and in the abdomen. Both legs ached. Headache was present. These symptoms continued during the first four days. The attending physician recorded a temperature of 100 F. on first day of illness and 103.2 F. on the third day. The pulse ranged from 96 to 120. Respiratory rate was 24. There was no vomiting. Constipation preceded and continued during the illness. Retention of urine commenced on the third day of illness and required several catheterizations daily. Motor and sensory paralysis was first noticed on third day after onset when the patient was awakened by a sensation of numbness in both feet which crept rapidly upward to lower chest. The right and left side were equally involved. A mild delirium was observed on the third, fourth and fifth days.

Findings.—My first examination was made on the sixth day of the patient's illness, at the time of admittance to the hospital. Patient was well developed; pubescent. Respiratory organs were normal; circulatory organs normal except increased rate of heart action. Some abdominal tympany was present. There was mild rigidity of recti abdominis. Urinary bladder was filled. Liver and spleen were apparently not enlarged. Tongue was coated. Skin had several blebs on lower extremities and over pubis, accidents from local application of heat. The mental state showed a little depression, but accurate, coherent replies were made to all questions. Close examinations of all cranial nerves showed normal functions. Motor power in the neck group and upper extremities was normal.

There was an absolute flaccid paralysis of all muscles in the lower extremities, not a trace of movement being detected at hip, knee, ankle or toe joints. The abdominal muscles showed much weakness. There was complete retention of urine and feces. All reflexes of head and upper extremities were normal. No epigastric, abdominal, gluteal, patellar or ankle reflexes could be elicited. Plantar stimulation produced no response. There was a complete sensory paralysis of the area supplied by the eighth dorsal cord segment and below, there being absolutely no response to touch, cotton, pin-pricks, thermal, deep pressure or vibration stimuli in the lower extremities. There was no true hyperalgesia or hyperesthesia from the seventh dorsal area upward, but movements of the thorax involving all the upper dorsal vertebrae caused intense pain and patient preferred to be turned to right side. Sense of position was entirely absent at toe-, ankle- and knee-joints, but feebly present at hips, the right equal to the left.

Course.—The subsequent course of the case was one so frequently observed in severe paraplegic myelitis cases from any of the various causes, in which the patient clings to life for weeks or months and finally succumbs to exhaustion with slowly increasing bed-sores, cystitis and other complications. Emaciation and exhaustion reached an extreme degree and death occurred Feb. 22, 1910, almost four months after the onset of the paralysis.

Full mental faculties were retained until forty-eight hours before the exitus when they became dulled with a rapidly oncoming coma. The palsied state never showed any substantial improvement. The muscles of the lower extremities remained functionless, flaccid and became markedly atrophic. A feeble right and left toe-extension sign could be demonstrated in the third week but not two weeks later. No change appeared in the findings as to deep reflexes. A partial sense of position in toes, ankles and hips but not at knees could be demonstrated in the second week but never after the fourth week. The superficial sensations remained unchanged. Trophic disturbances in the lower extremities developed early, and finally numerous large bed-sores appeared, the sacral eroding into the neural canal. Edema of the lower extremities was present during the last month. There was continuous pyrexia, and temperature ranging between 100 and 104 F. During the last few weeks of the illness there was recorded an irregular

* Read in the Section on Pathology and Physiology of the American Medical Association, at the Sixty-Third Annual Session, held at Atlantic City, June, 1912.

subnormal temperature. It dropped to normal after each of the two venesections and returned on the following day. A consultant diagnosed a cholecystitis in the fourth week, but an exploratory operation revealed no pathologic changes in the gall-bladder, tracts or neighboring viscera. Although no anesthetic was administered, the patient was entirely unconscious of the operative manipulations excepting some visceral pain when traction was made on the liver.

Laboratory Analyses.—Rachicentesis at the third lumbar space was performed on the sixth day of the illness with no pain or consciousness of the operation. Only about 1 c.c. of cerebrospinal fluid could be obtained on account of the difficulty in obtaining a flow. It coagulated more rapidly than normal blood, the drops finally hanging at the external opening of the needle in the form of congealed globules. The fluid was perfectly clear. A microscopic examination showed a mild pleocytosis, about 90 per cent. lymphocytes. Much coagulation fibrin was present. No microorganisms were found. Rachicentesis at the second, third, fourth and fifth lumbar spaces on the ninth day gave no fluid each puncture. I felt sure that the needle entered the lumbosacral cistern each time.

Fig. 1.—Section of cord and meninges at the ninth dorsal segment, showing epidural fat, adherent and thickened dura, arachnoid and pia.

A venesection was performed on the seventh day of the illness and 60 c.c. of blood withdrawn. One guinea-pig, one rabbit and one rhesus monkey were inoculated with a few cubic centimeters of the blood by the intraperitoneal route. All yielded negative results. Seventy-five c.c. of blood were removed on the eleventh day of the illness. Bouillon cultures with blood gave negative results in each instance.

A blood-count on the seventh day gave a leukocytosis of 32,100 with 88.5 per cent. polymorphonuclears and 5.5 per cent. lymphocytes. Ten days later the leukocyte count was normal. A Widal test on the tenth day gave a negative result.

Urinalysis on the sixth day gave acid reaction, specific gravity 1.017, no albumin, sugar or indican, few hyaline and granular casts, red corpuscles, some leukocytes and long motile bacilli. Later analyses showed continued cystitis.

Autopsy.—The post-mortem was performed two hours after death. There was no rigidity. Great emaciation and enormous sacral and trochanteric bed-sores were present; also lesser sores over bony prominences of feet, knees and scapula. There was some edema in the lower extremities and lumbar muscles; no serious changes in organs of chest or abdomen except evidence of wasting. The gall-bladder oper-

ative wound had healed nicely. On opening the skull the dura appeared normal. There was a little excess of fluid in the subdural and arachnoidean spaces. Some retraction of brain was seen. The convolutions appeared normal. All the ventricles were normal. The spinal cord dura appeared normal above the seventh dorsal segment. The dura, arachnoid, pia and cord were closely adherent, obliterating all spaces from the seventh dorsal segment to the first lumbar. (Fig. 1.) No fluid was present. The meninges were three or four times thicker than in the normal. The cord, to the naked eye, showed nothing abnormal above the seventh dorsal segment. It was diminished in size from constriction at the seventh and eighth dorsal segments. The lower five dorsal and lumbar segments had several cylindrical microscopic cavities extending up and down in the white and gray tissue. Especially do they appear in the posterior columns. (Fig. 2.) In the last two sacral and coccygeal segments appeared a softening which

Fig. 2.—Section at ninth dorsal segment, showing a large cavity in region of posterior column, canal centralis and thickened meninges.

gave a *Staphylococcus aureus* culture. No bacteria were found in the cord above this area.

The cord was hardened and fixed in alcohol and sections from each segment stained by the Nissl, picrofuchsin and hematoxylin methods for microscopic studies. The meninges of the lower dorsal segments presented the same sclerosed or hypertrophic appearance under the microscope as seen macroscopically. There was no infiltration of the pia and arachnoid. Above the seventh dorsal segment no marked minute changes were observed outside of ascending secondary tract degenerations. Some of the anterior horn cells in the cervical enlargement showed mild degenerative changes. From the eighth dorsal segment to the third sacral were seen all kinds of degenerative changes in the anterior horn cells. In a few of the segments all cells were present, but only a few were normal. Studying cells from various segments revealed axonal degeneration, tigrolysis in various stages, loss of nucleus, various forms of nuclear displacement and vacuole formations. (Figs. 3 and 4.) Often a fragment or a mere trace of a motor cell could be seen. There were some degenerations of the smaller cells of the posterior horn. Clark's column group of cells was much degenerated. The anterior, posterior and lateral white columns presented diffuse degen-

erations in all the tracts from the seventh dorsal segment downward. Diffusely scattered through the columns were observed microscopic cavities. Surrounding these cavities as well as the macroscopic ones, there was no indication of inflammatory reaction to be seen. As in the gray substance there were seen no leukocytic infiltrations or connective-tissue overgrowth. All the blood-vessels present in both the gray and white substance appeared to have normal walls and had no infiltrations.

Diagnostic Remarks.—During the early period of the illness, with considerable caution, a diagnosis of acute epidemic poliomyelitis with a transverse myelitis was made. The subsequent course of the case and the autopsy only added support to this diagnosis. The findings from the two lumbar punctures indicated the adhesions of the meninges in the dorsal cord. The coagulation of the cerebrospinal fluid has been observed in a few exceptional cases of undoubted acute poliomyelitis. Otherwise the cytology of the spinal fluid was

unexplained. I am inclined to believe that the overgrowth followed a severe edema and inflammatory process of the lower dorsal meninges. The thickened meninges were probably responsible for much of the myelomalacia.

It is very evident that we have a transverse myelitis to deal with, but transverse myelitis type of acute epidemic poliomyelitis (with apologies to our rather inaccurate nomenclature) is exceptional, and particularly so severe and definite a case. Reviewing the history of acute poliomyelitis from the time of our earliest definite knowledge of the pathology by Cornil and Chareot, in 1864, and Vulpian and Prevost, in 1866,² we have had added from time to time different types of acute poliomyelitis until we now have quite a number. Most of these types are named by their localization as encephalic, bulbar, spinal, polyneuritic, meningitic, ascending, etc.

I have reviewed extensively the literature of acute poliomyelitis and transverse myelitis for cases that might be akin to the one which I am reporting. Harbitz and Scheel^{3,4} report a case of transverse myelitis with autopsy, but indicate some uncertainty about diagnosing acute epidemic poliomyelitis of the transverse myelitis type. Strauss⁵ reports an illustration of edema and marked softening of the central nervous tissue

Fig. 3.—Three anterior horn cells from lumbar cord with complete tigrolysis. The upper and lower cells contain vacuoles.

Fig. 4.—Four anterior horn cells from lumbar cord, showing marked chromatolysis, axonal degeneration and nuclear displacement.

quite characteristic. It was certainly not characteristic of any purulent infection. The laboratory work with the blood indicated no bacteriemia. The continued fever was probably a result of trophic disturbance and a cystitis. Urinary retention or incontinence is very infrequent in acute poliomyelitis, but would be expected in any case of complete transverse myelitis from any of the numerous causes. There are some undoubted cases of acute poliomyelitis reported with paralysis of the sphincters. Likewise severe sensory disturbances are rare in acute poliomyelitis. Hyperesthesias and paresthesias are not infrequent. I have reported¹ one case of spinal poliomyelitis in which there was diminished sensation in the paralyzed member and a return of normal sensation within a few weeks. In support of the often observed fact that patients rarely die of their original disease, this case is an illustration in which the patient probably succumbed to several sequelae which also were the cause of the most striking clinical phenomena. The mode of development of the thickened meninges cannot be stated. There are a number of causes given for hypertrophic meninges, but some remain

in a fatal case of acute poliomyelitis. The cases of transverse myelitis with flaccid paralysis are infrequent. Cases in which a transverse myelitis has appeared with a more or less sudden onset, with a flaccid paralysis, and in which acute poliomyelitis has been considered in a differential diagnosis, or in which the etiologic factors have been uncertain or hazy or in which no conclusions were made regarding the true cause are reported by Nageotte,⁶ Evans,⁷ Pilger,⁸ Thompson,⁹ Spear,¹⁰

2. Medlin: Nord. med. Ark., 1896, No. 1.

3. Harbitz and Scheel: Pathologisch-anatomische Untersuchungen über akute Poliomyelitis und verwandte Krankheiten, Christiania, 1907, p. 162.

4. Harbitz and Scheel: THE JOURNAL A. M. A., Oct. 26, 1907, p. 1420.

5. Strauss: Epidemic Poliomyelitis, Rep. of Col. Inv. Com. on N. Y. Epid. of 1907, N. Y., 1910, p. 78.

6. Nageotte: Deux cas de myélite transverse aigue, Nouv. Iconog. de la Salpêtrière, 1895, viii, 325.

7. Evans: Case of Acute Transverse Myelitis; Post Mortem Examination, Lancet, London, 1894, ii, 683.

8. Pilger: Ueber einen Fall von genuiner Myelitis transversalis, Inaug. Thesis, Berlin, July 26, 1898.

9. Thompson: Transverse Myelitis, Railway Surg., 1901-1902, viii, 130.

10. Spear: Report of a Case of Transverse Myelitis in Woman Five Months Pregnant Followed by Very Easy Labor, Hosp. Bull. Univ. Maryland, 1908, iv, 365.

1. Skoog: Three Cases of Acute Poliomyelitis in Same Family at Same Time, Med. Herald, St. Joseph, October, 1911.

Frohmann,¹¹ Nogues and Sirol,¹² Ferreri¹³ and Burr.¹⁴ Brissaud¹⁵ has reported a case of transverse myelitis with syphilis as a cause in which the paraplegia was spastic at first and became flaccid in three or four months. Primangeli¹⁶ reports a clinical case of acute transverse myelitis appearing on the seventh day of illness in a case of measles in which there was a severe flaccid paraplegia. Williams¹⁷ has reported a case of transverse myelitis at the first lumbar level with acute poliomyelitis as a stated cause. Evans¹⁸ ascribes the cause in his case as due to influenza. Having reviewed the clinical and autopsy report, I favor very much the diagnosis of acute poliomyelitis. Pilger¹⁹ considers poliomyelitis in the differential diagnosis of his reported case but finally concludes that the cause was "spontaneous." He mentions among the causes for myelitis: syphilis, tuberculosis, acute infections as small-pox, typhoid, etc., chilling, wetting, excesses, trauma and concussion.

CONCLUSIONS

It is not within the province of this article to enter into a lengthy discussion of the nomenclature, but I wish briefly to emphasize my choice, which is based on pathologic findings. Among the various terms applied to the disease during the past seventy years, none is more applicable than acute poliomyelitis, or possibly acute epidemic poliomyelitis might be used. The former can be employed at least until the organism has been discovered and probably even then continued. We have learned much about the nature of the virus during the past two years through the work of Landsteiner and Popper,¹⁸ Flexner and Lewis,¹⁹ and a number of other investigators. It remains a fact that in a high percentage of cases of acute poliomyelitis the greatest pathologic alterations occur in the anterior horns of the lumbosacral and cervical swellings of the spinal cord and from this the most striking clinical manifestations appear, namely, the sudden, flaccid motor paralyses. In cases in which we wish to designate some particular type, additional descriptive terms used in this paper and elsewhere may be employed in our titles.

Rialto Building.

THE SPINAL CORD IN PERNICIOUS ANEMIA

WITH THE REPORT OF AN INTERESTING CASE OF FAMILY INVOLVEMENT *

ROBERT N. WILLSON, M.D.

PHILADELPHIA

The clinical picture of pernicious anemia and the microscopic findings in the blood are so thoroughly understood at present that they may be passed over without comment, even in a discussion of the changes found in the spinal cord in this symptom-complex rather than disease. It is well, however, to remember that there do

exist spinal cord and perhaps brain changes in every case of true pernicious anemia, and that many cases that fail of prompt recognition, owing to nervous symptoms, would at once become clear if this fact were kept in mind. The likelihood that both the clinical picture and the cord degeneration are secondary to and the direct result of a definite toxemia has also occurred to and been hinted at by many investigators. Yet, up to to-day, the nature and source of this toxin and its method of action are still matters of speculation, and even its avenue of approach and of attack are still to be determined.

Liechtm,¹ in 1887, Putnam² and Dana,³ in 1891 and in 1899, and later Pickett,⁴ Burr,⁵ Spiller, Camac,⁶ and in the last months, Camp⁷ have all described the combined sclerosis and degeneration of the cord which have come to be well recognized features in not a few instances of pernicious anemia. Spiller has shown that, at least, occasionally, the degenerative process extends into the brain axis; Pickett,⁴ and much later, Marcus⁸ (1903), and in the current year, Camp⁷ have described cases of pernicious anemia in which there became prominent the mental picture and the signs of paresis. There has not yet been reported, however, a case which has suggested so definitely as does that which forms the basis of this brief study the likelihood of a direct causal relation between the intestinal tract and both the peculiar anemic state and the degenerative process in the spinal cord.

The spinal cord lesions of pernicious anemia, as already repeatedly demonstrated, consist of a combined sclerosis of the posterior and lateral columns. Occasionally the posterior columns are alone affected, and there attends a definite clinical picture of tabes dorsalis. The lateral columns never constitute the only field of involvement. There is little tendency to implication of the gray matter or of the nerve-roots, or to contraction and shrinking of the cord, such as are seen in true tabes. Hemorrhages are frequently in evidence and also processes that can perhaps best be explained on the basis of thrombosis of the spinal vessels. Usually the degeneration appears to start in the upper dorsal or in the lower cervical cord, the posterior columns sustaining the earliest and most severe attack. In my case the lumbar cord showed the most marked evidence of disease (Fig. 1). The lateral columns in most cases are also sooner or later involved, and later in the course of the intoxication the anterior horns show both clinical and microscopic signs of disease. There is more or less proliferation of the neuroglia with rarefaction of the nerve fibers and often a complete dropping out of the latter. The cells in the gray matter may show degenerative changes and may be pigmented. Camac and others have described cavity formation in the cord, probably due to edematous softening and consequent degeneration. In Camac's case the cavity corresponded accurately to the distribution of a branch of the anterior spinal artery, and apparently a thrombosis in one of the small veins was the original cause of the excavation. In a word, it may be said that the pathologic process consists of a degenerative vacuolization of the nerve-cells, accom-

11. Frohmann: Ueber einen Fall von Myelitis Transversa mit Muskelwogen und eigentümlichen Veränderungen der elektrischen Reaktion, *Deutsch. Arch. f. klin. Med.*, 1905-6, lxxxvi, 339.

12. Nogues and Sirol: Myélite transverse avec paraplégie flasque d'emblée, *Gaz. heb. de méd.*, 1899, iv, 601.

13. Ferreri: Di una mielite transversa acuta dorsale inferiore e del suoi postumi, *Riv. Veneta di sc. med.*, 1896, xxv, 221.

14. Burr: Acute Transverse Myelitis, *Med. Bull., Philadelphia*, 1890, xviii, 401.

15. Brissaud: Myélite transverse et paraplégie flasque, *Rev. neurol.*, 1898, vi, 582.

16. Primangeli: Un caso di mielite lombare transversa consecutiva a morillo, *Pollcinico*, Rome, 1900, xiii, 1161.

17. Williams: Transverse Myelitis at the First Lumbar Level, *Louisville Month. Jour. Med. and Surg.*, Oct. 18, 1911.

18. Landsteiner and Popper: *Ztschr. f. Immunitätsforsch.*, 1909, ii, 377.

19. Flexner and Lewis: *THE JOURNAL A. M. A.*, Nov. 13, 1909, p. 1639.

* Read before the American Association of Pathologists and Bacteriologists, April 5, 1912.

1. Liechtm: Zur Kenntniss der perniciosösen Anaemie, *Cong. f. inn. Med., München. med. Wchnschr.*, 1887, xxxiv, 301.

2. Putnam: *Jour. Nerv. and Ment. Dis.*, 1891, xiii, 60.

3. Dana: *New York Med. Jour.*, 1891, iii, 279; *Jour. Ment. and Nerv. Dis.*, 1899, xxvi, 1.

4. Pickett: *Am. Jour. Med. Sc.*, 1904, cxxviii, 1032.

5. Burr: *Univ. Penn. Med. Mag.*, April, 1895.

6. Camac and Milne: *Am. Jour. Med. Sc.*, 1910, cxxxix, 563.

7. Camp: *Med. Rec.*, New York, Jan. 27, 1912.

8. Marcus: *Neurol. Centralbl.*, May 10, 1903, p. 453.