preparations, moreover, is given an advantage and prestige enjoyed by scarcely any other science, in that it has legal standing in every country through their pharmacopeias.

The pharmacopeias, as is well known, are the legal authority for medicines in their respective countries, which gives to the nomenclature a significance and meaning scarcely realized. There are in the United States Pharmacopeia over thirty different classes of preparations and extra-officially nearly as many more modifications of these official preparations. Yet, the question may well be asked, Have these pharmaceutical terms a purpose? If so, why are they constantly ignored? Manufacturers either ignore them or vie with each other in coining new names to substitute for them. The pharmaceutical nomenclature keeps step with therapeutic science; as soon as a new name for a new form of preparation is required, the pharmaceutical literature quickly responds and the name appears. It is safely asserted that there is not a single medicinal preparation now offered to the medical profession which could not be classified under the present nomenclature. Is it because the terms, tincture, liquor, elixir, syrup, spirit, fluid extract, pill, powder or ointment, are too simple, too commonplace, or hackneyed, and have lost their charm, or is it that the manufacturers seek to evade them in order better to be able to palm off some of these classes of preparations under mysterious and captivating titles, possibly a cloak for some old-time mixture of doubtful composition and questionable therapeutic virtues?

It is recommended that every medicinal article have a scientific title or descriptive name, and that the exclusive control of an article through registration of the name as a trade-mark or through copyright, be regarded as a perversion of the purpose and function of the patent laws. The greatest manufacturers, those who have attained the greatest reputation, prestige and patronage of the medical profession, even to-day, are those who rely on their names as the criterion for the quality of their products. What trade-name product has ever enjoyed the reputation of Powers & Weightman's quinin, Rosengarten hypophosphites, Mallinckrodt's potassium iodid, Squibb's chloroform, Parke Davis & Co.'s fluid extracts, Wyeth's elixirs, etc.? Manufacturers are shortsighted to ape the methods of the anonymous and pseudochemical companies in employing trade names, thus losing the prestige their names would confer on any other products they may introduce from time to time.

ACUTE HEMORRHAGIC PANCREATITIS.*

C. F. NEW, M.D.

Pathologist Central Hospital for the Insane.
INDIANAPOLIS.

In the following case the woman was a patient at this institution under the care of Dr. Stockton, who makes the following clinical report:

Patient.—The woman, aged 55, was admitted to the institution ten years ago. She had delusions of persecution, hallucinations of sight and hearing and was restless, sleepless, dull and melancholy.

History.—Little is known regarding her life and condition previous to admission, but from what could be gathered from herself and from the commitment papers the indications are that she had led a dissolute, immoral life and that she was addicted to the use of morphia. From the time of her admission until the onset of her illness her physical health was

good except for a moderate constipation which required the use of a mild purgative from time to time. Mentally there were no marked alterations, but she manifested some retardation of thought and occasional periods of irritability and quarrelsomeness, with obscene and immoral talk. On arising on the morning of March 14, 1904, she complained of feeling ill and suffered from nausea and vomiting, but was up and about until 11 a. m., when she began to complain of great pain and distress in the region of the stomach, aggravated when lying on the back, and of more or less tenderness on examination. The severity of the pain was such that an opiate was necessary to relieve it. At 6 p. m. the temperature was 102.6 and pulse 130, with but little alteration in the local or general condition. Vomiting continued, the vomit matter being bile-stained. The pain was dull and aching, its acuteness probably lessened by the opiate. The following morning there was considerable tympanitis, rigidity and tenderness, most marked in the epigastric region, and general jaundice supervened. Vomiting had somewhat lessened. The pain was still dull and boring in character. While attempting to get out of bed about 11 a. m. she suddenly became semi-conscious and died in a very short time.

Autopsy.—The autopsy, performed three and one-half hours after death, revealed the following condition:

The body was that of an adult female, white, apparent age 55, weight 145 pounds, height 5 feet 5 inches. The body was well-developed, well-nourished and rather fleshy. There was considerable general jaundice. There was postmortem rigidity in the neck, hands and feet. Postmortem lividity was slight posteriorly. The abdomen was prominent, tense and tympanitic. Liver dullness was indefinite.

On opening the abdominal cavity, the omentum and the intestines protruded. The intestines were considerably distended with gas. The omentum and peritoneum were of a pink, light-red color, and a thin, dark-red, odorless fluid ran out of the incision. About 350 c.c. was diffused through the general peritoneal cavity. The appendix was 6 cm. long and lay to the inner side of the cecum, curled behind the lower end of the ileum, and was apparently normal. The pyloric half of the stomach, the duodenum and the transverse colon were pushed forward by a dark-red, rather soft mass, lying posterior to them, and apparently involving the whole of the pancreas. Scattered throughout the omentum, the mesentery and the walls of the stomach and intestines were numerous small, white nodules, varying in size from a pin head to a split pea, rather soft, white on section, and most abundant in the neighborhood of the pancreas. All the structures in the immediate surroundings of the pancreas were intensely swollen and edematous and infiltrated with a fluid similar to that lying free in the peritoneal cavity. The infiltration and edema extended into the retroperitoneal tissues, surrounding both kidneys, and down into the postcecal tissue on the right side, giving the whole a boggy, sodden appearance and a dark-red color. The mass involving the pancreas consisted of coagulated blood diffused through the whole of the pancreas, except a very small portion of the head and tail. The source of the bleeding could not be found. The mucous membrane of the stomach and intestines were pale in color, swollen and edematous.

The liver was of a pale, yellowish-red color, softened. The lobules were indistinct. The gall bladder contained about 20 c.c. of dark, reddish-brown bile and a moderate number of small, brown-colored calculi. The bile ducts were patent and opened by a separate orifice from that of the pancreas into the duodenum.

The spleen was slightly softened, and the splenic artery and vein were filled with coagulated blood.

The kidneys contained several small cysts and were markedly congested. The bladder contained a few centimeters of slightly albuminous urine, but no sugar.

Apart from the abdominal conditions, there was some fatty infiltration of the heart, marked atheromatous degeneration of the large arteries, with some calcareous deposit, bilateral chronic pleuritic adhesions, cicatricial contraction of the lower lobe of the right lung, excess of cerebrospinal fluid, consider-

* Read before the Marion County Medical Society, Indianapolis.

able subarachnoidal exudate and marked atheroma of the arteries at the base of the brain.

REMARKS.

Incomplete as the clinical report is, it yet indicates some of the most prominent symptoms that should lead one to suspect an acute involvement of the pancreas. The onset of this condition is usually sudden, but may be preceded by epigastric uneasiness for a day or two. It occurs more frequently in adult males, and particularly in fleshy people. It usually begins with abdominal pain, which is intense, more or less diffuse and persistent, with paroxysmal exacerbations. It is increased by movement, and is associated with well-marked tenderness above the umbilicus, distention of the superior abdominal region and by vomiting. Nausea and vomiting are early symptoms and increase in severity. No relief follows the act of vomiting. The vomited matter consists of food and mucus, later of bile and blood in various stages of degeneration.

A condition of collapse soon follows. The abdomen usually becomes distended and tender. The bowels, as a rule, are constipated, rarely there is diarrhea. Slight jaundice is usually present and increases the longer the patient lives. The mass may be palpable but usually the distention and tenderness are so great as to render palpation very difficult, if not impossible except under anesthesia. Severe attacks are usually fatal in 24 hours. When less acute, a gangrenous condition is often produced and proves fatal in from two to five days from absorption of the toxic material. When the condition is still less acute it may lead to abscess formation, or become subacute, or chronic; and in mild forms it may lead to resolution and recovery.

The hemorrhage may precede or take place during the inflammatory process. If it precedes the inflammation, the bleeding is usually within and without the glandular substance, and usually runs an extremely acute course. If it occurs during the inflammatory process, the bleeding is less profuse and runs a less acute course.

The diagnosis is difficult, as there are several conditions with a similar group of symptoms. The principal ones are: 1, Acute ptomain poisoning; 2, perforation of a gastric or duodenal ulcer; 3, gangrenous appendicitis; 4, phlegmonous cholecystitis; 5, acute intestinal obstruction.

Judging from the important part taken in the digestive process by the pancreatic secretion that is discharged into the duodenum, and in the process of metabolism by the internal secretion that is given off by the pancreas to the blood or lymph stream or both, one would expect that the clinical manifestations of a diseased condition would be so characteristic that a diagnosis would be comparatively easy, but there are several reasons why this is not the case:

1. The pancreas is rarely involved in a pathologic process without other organs being affected, particularly the biliary ducts, the stomach, and the duodenum.

2. Every function performed by the pancreas can be performed to some extent by some other organ in the body except the glycogenic function.

3. The greater portion of the gland may be destroyed and yet the remaining portion apparently may be able to carry on the normal function of the entire gland.

While these conditions may give rise to very conflicting and perplexing combinations of symptoms, yet there are some pronounced features manifested which should lead one to suspect a diseased condition of that special organ. Fitz has laid down a rule that when a previously

healthy person or sufferer from indigestion is suddenly seized with violent pains in the epigastrium, followed by vomiting and collapse, and within twenty-four hours by a circumscribed epigastric swelling, tympanitis, and resistance, with slight rise of temperature, one should suspect acute pancreatitis.

In a lecture delivered before the Royal College of Surgeons of England, Mr. P. J. Cammidge¹ stated that there were present in the urine certain products which were almost if not quite pathognomonic of acute pancreatitis. His method of examination to prove their presence is as follows: To 10 c.c. of filtered urine is added 1 c.c. of strong hydrochloric acid, boiled gently for 10 minutes on a sandbath. A mixture of 5 c.c. of filtered urine and 5 c.c. of distilled water is then added and the whole cooled in running water. The excess of acid is then neutralized by slowly adding 4 grams of lead carbonate. The whole is again filtered, the flask rinsed with 5 c.c. of distilled water. To this are now added 2 grams of sodium acetate and 0.75 gram of phenylhydrazin hydrochlorate and boiled for three to four minutes on the sandbath. It is then poured into a test tube and allowed to cool undisturbed, when a flocculent precipitate forms consisting of rosettes of golden-yellow crystals. The presence of sugar of albumin must be previously gotten rid of, as these substances interfere with the process and the results. As somewhat similar results are obtained in patients suffering from certain diseases in which active tissue changes are going on, a differentiating test must be applied, which consists in taking 20 c.c. of filtered urine and adding 10 c.c. of saturated aqueous solution of mercuric chlorid, filtering. To 10 c.c. of the filtrate add 1 c.c. of strong hydrochloric acid and boil as in the former process. It is then diluted with 5 c.c. of the former filtrate, and 10 c.c. of distilled water, cooled and treated as in the former instance.

The practical results of these examinations of the urine by these two methods Cammidge summarized as follows: If no crystals are obtained by either method the pancreas is not at fault, and the explanation of the symptoms must be sought for elsewhere. If crystals are obtained by the first method and not by the second, active inflammation of the pancreas is present and surgical interference is generally indicated. In acute inflammation of the pancreas the crystals are fine and dissolve in a 33 per cent. solution of sulphuric acid in about thirty seconds. In chronic inflammation the crystals are coarser and dissolve in one to two minutes. If crystals are obtained by both methods there may exist: 1, Malignant disease of the pancreas; 2, chronic disease of the pancreas; 3, non-pancreatic diseases.

In malignant disease the crystals are very coarse, and require from three to five minutes to dissolve, and operation is contraindicated. In chronic inflammation the crystals are smaller and dissolve in from one to two minutes. In non-pancreatic disease the crystals dissolve in one minute. Thus it seems that the size of the crystals and their solubility are directly dependent on the acuteness of the inflammatory process.

As regards the treatment, the pathologic condition indicates that practically only one line of treatment can be beneficial, and that is evacuation of the septic material, arrest of the hemorrhage, and free drainage by surgical methods. Even though the mortality following operation in the acute stage is very high, 36 out of 59

1. Cammidge: *Lancet*, July 1, 1905; abstracted in *THE JOURNAL, A. M. A.*, July 29, 1905, p. 359.

operative cases proving fatal, yet when one considers the fact that the disease almost invariably ends fatally when not treated surgically, an operation is strongly indicated.

POISONING DUE TO THE PAPA. (*ASIMINA TRILOBA*).

M. A. BARBER.

Associate Professor of Bacteriology and Pathology, University of Kansas.

LAWRENCE, KAN.

Some seven or eight years ago, in late summer, I found in this vicinity several fruits of the common papaw which had fallen to the ground and ripened prematurely. The fruit was ripe enough to be palatable, and I ate a small portion, perhaps equivalent to one fruit. About six hours later I experienced an urticaria, beginning with itching and burning in the hair at the back of the head and soon extending over a considerable part of the body. Accompanying symptoms were nausea, throbbing at the head and disturbance of the bowels. The urticaria would at times diminish suddenly; then the nausea would increase, to be relieved a few moments later with the reappearance of the "hives." The symptoms lasted several hours, but they had almost wholly disappeared by the next morning and the skin had resumed its normal condition. A friend ate some of the fruit at the same time that I did, but suffered no ill effects whatever.

I had never before experienced similar symptoms, and I suspected the papaw; but I was not sure of the cause until, a season or two later, I had an opportunity of repeating the experiment. This came when, visiting the same grove at about the same time of the year as on the former occasion, I found similar fruits and ate a small quantity to test the effect. The same train of symptoms followed after about the same interval of time, though this time the illness was considerably more severe, and I experienced a decided faintness along with the symptoms of the previous case.

In the autumn of 1904 I undertook some more systematic experiments on the nature of this poisoning. It seemed to me worth while to investigate the matter for the sake of any information I might be able to obtain on the general question of poisoning resulting from individual susceptibility to the action of certain fruits, especially since I had a subject so conveniently at hand. I desired particularly to find out if repeated poisoning confers any immunity on the susceptible person, and to ascertain if any particular condition of the fruit is necessary in order to cause the illness. I have eaten ripe papaws since childhood, but had never before experienced poisoning from them or from any other fruit. I was also interested in the papaw itself; for, while not altogether a popular fruit, it is much eaten in many parts of the Mississippi Valley, where it is abundant, and some attempts are being made to cultivate it and obtain improved varieties of the species.

The following is a brief summary of my experiments:

October 1. Ate small portions of several fruits. Eleven hours later I experienced urticaria and some of the other symptoms noted above, but the illness was very mild.

October 7. Ate 24 grams of fruit brought to me from a locality at some distance from the source of the fruits used in the other experiments. The fruit was probably picked up from the ground, because it was slightly blackened on one side. After about ten hours I experienced urticaria with slight nausea, but the symptoms were comparatively insignificant.

October 10. Ate 21 grams of the same fruit, which had been

kept in a refrigerator since October 8, and had ripened somewhat in the interval. No effect.

October 15 to October 18. Ate a whole fruit ripe from the tree and a whole fruit ripe from the ground. Also ate a whole fruit not fully ripened at the time of collection, but allowed to ripen three days after gathering, and ate 21 grams of a fruit taken green from the tree.

In none of these four experiments was there any effect. All of these experiments were made with fruit apparently in a normal condition, and differing only in the degree and manner of ripening.

On account of all these failures I began to suspect that I might have acquired some degree of immunity to this fruit, but the next experiment disproved this theory.

October 19. Ate 23 grams of a fruit which had fallen prematurely from the tree. The quantity eaten was equivalent to about one-third of a fruit of average size. It had been kept four days at room temperature and had ripened somewhat at the center, but the outer part was still hard. Its taste was bitter.

The effects began to be noticed three and one-half hours later. An urticaria began on the crown of the head, then appeared on the body where suspender or garter pressed the skin. Then it spread over the whole body, the face, hands and feet being the latest to be covered. Nausea with vomiting, faintness, chill and disturbance of the bowels followed, the symptoms being decidedly more severe than in any previous case. The following day little remained of the illness, except that the face and hands were slightly swollen, the throat a trifle sore and the skin ready to break out into a mild urticaria on irritation.

This experiment, the most decisive of all, indicated that the four previous poisonings, two of them within twenty days, had not conferred any great degree of immunity; and, further, that an abnormal condition of the fruit seems, in my case, at least, to be necessary in order that poisoning may take place.

During the autumn of 1905 I resumed the experiments, trying smaller doses of fruit similar to the one which had caused the poisoning of the previous year.

September 11, I ate 3 grams of such fruit, and September 12, 10 grams of the same specimen kept on ice over night. October 10, I tried 25 grams of a fruit taken from the tree soft but not fully ripe.

In none of these experiments was there any effect. It would seem that the dosage was too small, or more probably, the fruit not in the condition in which it is poisonous for me.

During the past year or so I have made many inquiries of physicians and others in regard to poisoning from this fruit, and I have learned of fifteen or twenty cases attributed to this cause. Dr. Cleveland of Burlington, Kan., a locality where the papaw is abundant, told me of two cases, in one of which he found the patient unconscious. On recovery to consciousness the patient exhibited urticaria. Dr. Manson of the same locality reports several cases of poisoning due sometimes to eating the fruit and sometimes to the mere contact of the surface of the fruit to the skin. Dr. J. E. Buckman of Modesto, Ill., wrote me of three cases which he attributes to the eating of papaws. In all three cases the patients reported that the fruit they had eaten was so ripe that the surfaces had begun to turn black. One case was characterized by fever, a burning sensation in the stomach and throat, with an eruption on the membrane of the throat. There was an eruption externally also, especially on the face where pustules formed which were large and slow to heal. The other two cases exhibited similar characteristics, except that there was no in-