This article was downloaded by: [University of Calgary]
On: 04 February 2015, At: 18:04
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer
Street, London W1T 3JH, UK

Annals and Magazine of Natural History: Series 5

Publication details, including instructions
for authors and subscription information:
<http://www.tandfonline.com/loi/tnah11>

Note on the carotids of *Rhea americana*

Franklen P. Evans Esq.^a

^a University College , Bristol
Published online: 07 Oct 2009.

To cite this article: Franklen P. Evans Esq. (1883) Note on the carotids of
Rhea americana, *Annals and Magazine of Natural History: Series 5*, 11:66,
458-458, DOI: [10.1080/00222938309459183](https://doi.org/10.1080/00222938309459183)

To link to this article: <http://dx.doi.org/10.1080/00222938309459183>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever

caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Note on the Carotids of Rhea americana. By FRANKLEN P. EVANS,
Esq., University College, Bristol.

In dissecting a young but almost full-grown specimen of *Rhea americana* I find that the right carotid is evidently present, though it is much smaller (about $\frac{1}{10}$ inch diam.) than the left. Its position for some distance past its origin corresponds to that of the left carotid; but subsequently, instead of converging to meet the latter, which runs along the hypapophysial canal, it continues onwards by the side of the right pneumogastric nerve and jugular vein. As the late Mr. A. H. Garrod, in his paper on the carotid arteries of birds (P. Z. S. 1873, p. 470), states that normally the left carotid alone is present in this species, it is possible that this difference is due to age; if so, it is interesting to find that a structure, the absence of which in the adult bird is regarded by Mr. Garrod as of sub-family importance, is originally present in the young bird, persisting even until close upon sexual maturity. Perhaps this may explain the conflicting statements of Prof. Owen with regard to the anatomy of *Apteryx* as quoted by Mr. Garrod.

On the Origin of Alternation of Generations in Hydro-Medusæ.

By W. K. BROOKS.

It is hardly possible that the form of development which we now find in most of the hydro-medusæ can bear any close resemblance to their primitive life-history; and there are many reasons for believing that alternation of generations has gradually arisen through the modification of "metamorphosis."

In *Cunina* we seem to have the ancestral form of development, a direct metamorphosis without alternation. The interesting and remarkable life-history of *Cunina* was first described by Prof. John McCrady, who found inside the bell of a hydro-medusa, *Turritopsis*, at Charleston, S. C., a number of hydra-like larvæ attached by short tentacles to the subumbrella, and furnished with a very long and flexible proboscis, with the oral opening at its tip. These larvæ are parasitic; and they obtain their food by inserting the proboscis into the mouth of the *Turritopsis*, and thus sucking from its stomach the food which it contains. In his first paper upon the subject Prof. McCrady stated his belief that these larvæ were the young of the *Turritopsis*, which carries its young inside its umbrella and nourishes them with partially digested food from its own stomach. Although McCrady soon corrected this error, and showed that they are not the young of *Turritopsis*, but the larvæ of *Cunina*, Romanes, in his recent work on 'Animal Intelligence,' has quoted McCrady's error without his correction, and refers to this instance of parasitism as a case of maternal care for the young among the Cœlenterata.

During the past summer both *Turritopsis* and *Cunina* were extremely abundant at Beaufort, N. C., throughout July and August, and I was fortunately able to trace the life history of each of them.