XANTHOMA.*

JOHN V. SHOEMAKER, M.D., LL.D.

Professor of Skin and Venereal Diseases in the Medico-Chirurgical Hospital.

PHILADELPHIA.

History.—A lad, 18 years of age, afflicted with peculiar, characteristic and unusual lesions on the hands, knees, posterior aspect of the thighs below the buttocks and on the eyelids, was brought by his father to the dermatologic clinic of the Medico-Chirurgical Hospital, was shown to the class on several occasions and made the subject of a clinical lecture.

Physical Examination.—Small in stature, general lack of development. Greatest weight, 79; present weight, 67. Thin, with almost no subcutaneous fat. General appearance that of a boy of twelve. Face marked with lines, giving the appearance of old age. Skin pale, showing a general anemic condition; complexion yellowish, subicteroid. Tongue flabby and pallid.

Chest narrow, scapulae prominent; marked depression above and below clavicle.

Circulation poor; pulse weak. Heart's action increased, apex beat displaced, diffuse; functional cardiac murmur.

Liver enlarged, extending from fifth interspace to an inch below umbilicus.

Spleen enlarged; could be easily palpated.

Urine: An analysis of the urine gave the following results: Sp. gr. 1014; acid reaction; no albumin; no sugar; no sediment; hyaline cylindroids; blood cells, uric acid and urates in abundance.

Blood.—An examination of the blood by Dr. L. Napoleon Boston revealed: Red corpuscles, 3,990,000; white corpuscles, 20,000; hemoglobin, 60 to 67 per cent. Dr. Boston also contributes the following notes concerning the blood: The red cells were pale. There was marked variation in the size of the red cells, many of them being greatly oversize, while others were scarcely one-half the normal diameter. The viscosity of the blood was diminished. The red cells stained feebly, as a rule; cells nearly colorless were often seen. The central pale spot in the red cells was much enlarged, and at times this pallor extended to near the margin of the cell. The hemoglobin had been extracted from areas in the protoplasm, and these pale areas overlying such degeneration resembled somewhat the effect produced by the malarial parasite. No pigment was seen in the red cells. The differential leucocyte count was as follows:

	Per cent.
Polymorphonuclear cells	66.0
Large lymphocytes	4.5
Small lymphocytes	15.4
Eosinophiles	6.0
Myelocytes	3.0
Large mononuclear	4.5
Transitional cells	0.6
	100.0

Description of Lesions.—The palmar and dorsal surfaces of both hands were covered with small growths, which had been in existence for about six months. They first came on the hands. The affected regions did not ache spontaneously, but were painful on pressure. The boy could not grasp objects firmly by reason of the tenderness of the palms. The earliest lesions occurred on the palms, especially in the flexures of the metacarpophalangeal and phalangeal joints. They were symmetrically arranged, and a description of one hand will do for both. An artist, with deliberate intent and skill, could not more accurately copy the picture from palm to palm. The palmar aspect of the thumbs and fingers were studded with discrete tubercles, most prominent in the flexures of the joints. At the base of the thumbs were numerous tubercles which presented a linear arrangement. Lesions were similarly distributed, corresponding to the principal lines of the palms. They were also perceptible between the fingers. The disease was likewise conspicuous over the knuckles, although the lesions in

these situations were less numerous and distinct than on the palms. An equal symmetry was observed in the location of the growths on the backs of the hands. Lesions were observed between the fingers, and in the flexures of the elbows they were confluent. The distribution was symmetrical on both arms. Large lesions were scattered, though more sparsely, over the back. Some of the largest tubercles, about the size of a pea, were situated below the gluteal fold. Lesions existed behind each ear and on the scalp. The upper and lower lids of both eyes were covered with numerous small lesions. They were present on the dorsal and plantar surfaces of both feet and between the toes. Walking was impeded and kneeling was painful.

In various situations the lesions were both nodular, or discrete, and confluent. Small nodules were seen on the penis. A patch existed on the mucous membrane of the mouth, corresponding to the attachment of the upper lip to the alveolar process. There was a spot of the same kind on the lower lip and others at the angles of the mouth, and one on the uvula. Several were situated within the nostril. Those located on the

Fig. 1—Palm of a typical case of xanthoma.

mucous membrane were rather more conspicuous than those on the skin, owing to the contrast between their yellow color and the red of the mucous membrane.

Many of the lesions were slightly elevated; others were plane with the surface. They were of a yellowish color and resembled fragments of chamois leather embedded in or attached to the skin. As the father expressed it, they are particularly numerous where the skin is "slack," or where it is habitually brought in contact with objects.

Diagnosis.—Although this case represented a rare affection, yet its features were so distinctive that there could be little difficulty in forming a diagnosis. The yellowish color might possibly be suggestive of pus, but the contents of the tubercles were neither fluid nor fluctuating; the lesions were rather soft, but nevertheless of firm consistency. There was no resemblance to acne or any other inflammatory disease of the skin. There was more analogy to milium, which is sometimes of a yellow color, appears principally on the upper eyelid and other parts of the face, and may remain stationary for years. Mil-

* Read at the Fifty-fifth Annual Session of the American Medical Association, in the Section on Cutaneous Medicine and Surgery, and approved for publication by the Executive Committee: Drs. H. W. Steilwagon, J. A. Fordyce and H. G. Anthony.

ium, however, contains sebaceous matter, which may be forced out by pressure after the surface has been pricked. This was not possible with the tubercles of the case now reported.

CLASSIFICATION.

The permanence of the lesions indicates that the disease should be classified among the new growths or tumors of the skin. The structure and the distinctive hue of the lesions are evidence that it could be no other than the affection which is named for its color—xanthoma, from *ξανθός*, yellow. Ordinarily, there is little scope for error, although multiple dermoid cysts have been mistaken for xanthoma, from which they could only be differentiated by microscopic examination. Urticaria pigmentosa, which occurs during infancy or early childhood, has sometimes been confused with xanthoma, a fact which is suggested by the very name which some writers have bestowed on that rare variety of nettlerash, viz., xanthelasma. The possibility of error is all the more

Fig. 2.—Lesions of xanthoma.

marked when we reflect that pigmented wheals may persist as permanent new formations of the skin and subcutaneous connective tissue, and that their color may be yellow. Wheals, however, are of firmer consistence than the tubercles of xanthoma. Itching is a much more prominent feature in urticaria than in xanthoma.

SYMPTOMS.

Xanthoma is divided into two varieties, according as the lesions are plane or raised—xanthoma planum and xanthoma tuberosum. The spots of xanthoma planum are smooth, sometimes slightly elevated and of a yellowish hue, which has been aptly compared to that of chamois leather. On account of the buff or yellowish color, the disease has also been known as xanthelasma. The abnormal pigmentation has suggested another title for the malady, vitiligoidea. A multiplicity of names or titles for the same disease is undesirable, as it is liable to lead to confusion.

The lesions of xanthoma may be round or elongated. In the case now under consideration, many exhibited a linear arrangement. They may be as small as a pin's head or as large as a pea, and, indeed, in some instances

have acquired much larger dimensions. Duhot has recorded a case in which some of the tumors were as large as a walnut. W. Moser encountered a case of a boy in whom multiple mushroom-like growths existed from the size of a pea to that of a hen's egg or larger; they were not tender or painful.

Xanthomatous skin does not seem to be thickened when pinched up between the fingers. After attaining a certain size the lesions generally cease to grow. The eyelid is a favorite location. Cases in which the disease is limited to the eyelid are much more common than those in which the lesions are more widely distributed, and which have been called xanthoma multiplex. In young persons most of these cases are congenital; in adults they are associated with jaundice or diabetes. The lesions have been seen on the cornea, in the mouth, esophagus, trachea, spleen, lining membrane of the bile ducts, liver, heart, aorta, and in the abdominal muscles. Tendons and serous membranes have also been implicated. Adjacent spots may coalesce. Xanthoma of the cornea is extremely rare. Gaucher and Herscher described a case¹ in which the cornea was involved, and stated that they had only been able to find one other case previously reported. The patient was three and one-half years of age, and was probably syphilitic. There was double interstitial keratitis. The keratitis was prior to the ocular

Fig. 3.—Typical lesions of xanthoma.

xanthoma, the nutrition of the cornea had been reduced, and it had become a spot of least resistance. On the right cornea was a yellowish deposit of irregular shape and plane edges. There were characteristic tubercles of xanthoma on the cheeks. The nodules of xanthoma tuberosum are, as a rule, about the size and color of those of the plane variety, and round in shape, though they have been seen as large as a small apple. The large growths are due to an aggregation of tubercles, and are more irregular in form. In both varieties of xanthoma the color may vary from a light to a dark shade of yellow.

Xanthoma is more frequent in women than in men. The most usual site of commencement is on the inner canthus of the eyelid, and more particularly of the left eyelid. It begins as a macule. The coalescence of several patches sometimes forms a semicircle around one or both eyes. Subsequently other parts may be attacked. The disease is generally tardy in its progress. Large growths overhanging the eye have been known to interfere with vision. Korach and Hertzka have reported

1. Journal des Maladies Cutanées et Syphilitiques, 1900, p. 48.

cases in which the evolution occurred within a few weeks, involving the greater part of the body, accompanied by icterus and pruritus. Dr. George T. Jackson of New York observed a case in a boy of 5 which began when the patient was 3 months of age, and attacked all parts of the body except the hands, feet and scalp. Prof. McCall Anderson witnessed an apparently congenital case in a child 3 years of age. Xanthoma, especially of the tuberous variety, is rare in children. Richter had as a patient a 9-month-old girl afflicted with generalized xanthoma, marked especially by tumors. The mother thought that the disease had followed the bites of insects when the babe was 3 months of age.² Most cases occurring in early life are hereditary or congenital.

Some cases of xanthoma have been described which seemed to be of hereditary origin. Török knew of a family in which it had affected three generations. Hilton Fagge mentions a mother and daughter who had xanthoma, and in whose family it had occurred for four generations. Hutchinson reports a case of xanthoma in two brothers and a paternal grandmother. Crocker had a case in a stout, country lad of 18, in which the lesions were symmetrically situated on the elbows and knees; an elder brother had the same disease on the elbows. Xanthoma is rather commonly associated with disease of the liver and jaundice. It has been known to develop during the course of diabetes or gout. Jonathan Hutchinson has described an interesting case. The patient had inherited gout, of which he had had several attacks, and of which physical evidence existed in enlarged bursæ and fusiform swellings in connection with numerous tendons. In consequence of a severe fright, jaundice supervened, after which xanthoma appeared on the eyelids. The patches were symmetrically arranged. Subsequently small streaks of the same disease appeared over the right olecranon process. The patient's father and maternal grandmother had had xanthoma of the eyelids. Under the title of xanthomatosis the case has been described of a woman, 35 years of age, who had slight mitral insufficiency, stationary for several years, rheumatism, hepatic colic with or without jaundice, but no gallstones. This patient had xanthoma planum of the eyelids and a small fibro-fatty tumor of the abdominal wall.

In the opinion of Besnier, xanthoma of diabetic origin manifests a tendency to spare the eyelids and locate itself in the mouth. This writer furthermore states that buccal xanthoma progresses rapidly, but may spontaneously disappear. Diabetic xanthoma runs a more rapid course than when the disease is due to other causes. A few cases are on record in which xanthoma preceded the glycosuria. Another case is mentioned in which sugar had entirely disappeared from the urine fifteen months before an eruption of xanthoma occurred. Sequiera had a case of diabetic type without glycosuria. The patient was a man, 45 years of age, in whom the disease had existed for six months without any impairment of the general health. He had never had jaundice, but there was slight enlargement of the liver. According to Malcolm Morris, jaundice never accompanies the xanthoma of diabetics. Up to the year 1900 but 30 cases of diabetic xanthoma had been reported. Several have been published since that date. An interesting case was studied by Sherwell and Johnston.³ The patient was a diabetic woman, 40 years of age, who had xanthoma on the arms, elbows, knees and buttocks. In fact, no part of the body

was totally exempt. It was present also on the palms, soles and scalp. The eyelids, an ordinary site, were spared. The lesions were painful when touched. It was difficult to handle objects, walk or lie down in bed. She had a sensation as if foreign bodies, such as peas or balls, were embedded in the skin. Antidiabetic diet had a remarkable influence on the eruption, which rapidly disappeared. Coincidentally she lost the feeling of foreign bodies in the skin. In the course of two or three weeks she was able to use her hands, walk and lie on her back. When the patient first came under their observation, the eruption, when seen from the other side of the room, looked very much like confluent smallpox in the suppurative stage. Dr. Abraham has had a case of xanthoma in a diabetic woman, 32 years of age. The eruption had been in existence for four months. She was subject to biliousness. The lesions were painful when pressed. Women are much less liable to diabetic xanthoma than men.

Xanthoma diabeticorum, or the diabetic type as it is also called, differs in some particulars from the other varieties of the disease. The onset is more rapid in diabetic cases and the eruption may speedily vanish, as in Sherwell and Johnston's case. The lesions are of somewhat different aspect. They present some signs of inflammation, and consist of yellowish points on hyperemic bases.

The lesions of xanthoma usually remain for a long period or during life. In a few instances they have spontaneously disappeared without leaving pigmentation or cicatrices.

In cases of xanthoma multiplex the lesions are generally both plane and nodular. The spots are more common on the eyelids and flexures; the tubercles most frequently occur on the knuckles, elbows, knees and buttocks. The disease is commoner on the limbs than on the trunk. Lesions on the knees may be the source of discomfort or pain.

PSEUDOXANTHOMA ELASTICUM.

Under this title Balzer described a form of eruption which bears some resemblance to that of xanthoma, being of a yellow color. Microscopically, however, it seems to be due to fatty degeneration of elastic tissue. A second case has been reported by Chauffard, and a third by E. Bodin. In these three cases the lesions were found on the front of the abdomen, base of the neck, axilla and groin; the knees, elbows and eyelids were free from disease. The evolution was slow; there was neither jaundice nor liver trouble, but there was tuberculosis in every case. The lesions were small masses of granular substance constituted by swollen and segmented elastic fibers, etc. Bodin regards this species as only provisionally attached to true xanthoma.⁴

PATHOLOGY.

Xanthoma is a connective-tissue new growth undergoing fatty transformation. The lesion consists of irregular, uncircumscribed aggregations of large cells which have a clearly defined outline and possibly a cell wall with small nuclei, sometimes several in a cell, and with cytoplasm, characterized by closely massed fat globules. The cells occupy the spaces between the bundles of connective-tissue fibers belonging to the reticular layer of the corium, sometimes extending to the papillary layer. The epidermis shows slight atrophic changes. According to Touton, the fat is deposited in proliferated connective-tissue elements. In sections made by Prof.

2. *Journal des Maladies Cutanées et Syphilitiques*, 1903, p. 531.

3. *Journal of Cutaneous and Genito-Urinary Diseases*, Sept., 1900.

4. *Journal des Maladies Cutanées et Syphilitiques*, 1901, p. 122.

Joseph McFarland, there were scarcely any evidences of inflammation exhibited. Some writers, as Crocker, Chambard and others, believe that the disease originates in a chronic inflammation with proliferation of round cells. As these are not invariably present, however, Touton and others refuse to accept the inflammatory theory. All authorities agree that it is the fat in the cells that gives them their yellow color.

Pollitzer, following Unna, teaches that xanthoma of the eyelids is due to degeneration of muscular fiber embryonically displaced, and that it differs from xanthoma tuberosum and xanthoma diabeticorum in this respect. Clinically, I should regard the different forms of the disease as the result of a process identical in its nature, though differing in some histologic details perhaps, for, as in the case which I now report, we see plane and nodular lesions on the same person. Not infrequently the disease begins on the eyelids and spreads to other parts of the body.

ETIOLOGY.

In reviewing the course and associations of the disease, I have anticipated some of the chief points of etiology. I have spoken of the close connection which often exists between xanthoma and disease of the liver, gout and diabetes, and that it may be of congenital or hereditary origin. Xanthoma palpebrarum is more common in women than in men, contrary to what occurs in xanthoma of diabetics. It generally begins in middle life and is most frequent in people of dark complexion.

TREATMENT.

When xanthoma is confined to the eyelids, or when the tubercles are not numerous, they may be removed by excision. Great care must be taken in order to avoid the occurrence of ectropion or epiphora. Cauterizing and curettage have sometimes succeeded. Kaposi used soft soap. Morrow, in a case of nodules on the soles and knees, effected an improvement by means of a 25 per cent. salicylic acid plaster. Dr. James C. Maguire made use of monochloroacetic acid, applying it to a small spot at a time and gradually going over the entire diseased area. A 5 to 10 per cent. solution of mercuric chlorid in colloidion has been recommended. Electrolysis has been used by Hardaway, Wende, Pansier and Leviscur. Dr. Willmott Evans treated a case by means of the x-ray. The disease affected both elbows, and was of seven years' duration. He subjected one elbow to ten exposures of fifteen minutes each with a very satisfactory result, the elbow becoming practically clear.

Wilson and Besnier advised constitutional treatment. The former gave nitrohydrochloric acid, with an occasional blue pill, followed by arsenic. Besnier administered phosphorus first, succeeded by arsenic, and reported a rapid removal of the growths by this method.

The diet and other measures which are of service in diabetes have exerted an excellent influence on xanthoma, due to that disease. Arsenic has also been given with asserted benefit.

TREATMENT AND PROGRESS OF THE PRESENT CASE.

The patient was first seen on Oct. 16, 1903. On account of his unsatisfactory general condition he was then ordered:

R. Ferri lactatisgr. 1/4|016
Strychninae sulphatisgr. 1/100|00065
Acidi arsenosigr. 1/100|00065
Salicinigr. ii|12

M. Ft. pil. No. i. Sig.: One such pill to be taken three times a day.

No local application was given, as the disease was so widely spread. On November 6 the boy's subjective condition was

somewhat improved. On November 13 he reported that he felt better, and it was thought that he was a shade less pale. He was kept under observation, and on Jan. 22, 1904, was again brought before the class. At that date it was evident that he was losing ground. The anemia had increased. On March 23 he was thinner—a mere skeleton—and was having night sweats. An examination of the chest gave a negative result.

On April 23 the patient's abdomen was distended and very hard on the right side. He was very weak, had lost appetite, had abdominal pain, and sometimes vomiting. He was placed on:

R. Acidi nitrohydrochlorici dil.....m. x|60
Tinct. gentianae co.3i|75
M. Sig.: One such dose to be taken after each meal.
He was also ordered:
R. Pil. strychninae sulphatis.....1/60|0011
M. Sig.: For one dose (before meals).

Xanthomatous deposit was probably taking place in the liver.

The boy had been watched most carefully by the clinical assistants. His father brought him to the dispensary at stated intervals. With the exception of some slight improvement at first, his condition gradually grew worse. At about the time of the last note one of the assistants visited the patient at his home—he lived in a distant part of the city—and found him in a feverish state, very much prostrated.

REMARKS.

This is a very interesting case, an excellent example of xanthoma multiplex. The exquisite symmetry with which the lesions were arranged, more particularly on the hands and arms, claimed attention, and is quite characteristic of the variety of the disease by which we were confronted. It was a case of xanthoma multiplex exhibiting patches, tubercles and nodules, both discrete and confluent. Although the eyelids are the commonest site of the eruption, and although they were typically involved in this case, yet it was not in this situation that the disease first made itself manifest.

The morbid process began at an early age in this patient. With the exception of the few congenital or hereditary cases, xanthoma develops, in most instances, after the age of 40. The lad was naturally of light complexion, which is also against the rule. The rather plentiful deposit of xanthomatous material on the mucous membrane is among the less usual manifestations. Although I have observed a number of cases of the disease, yet this was the first occasion in which I had seen it attack the mucous membrane.

The poor physical development of the lad and his unpromising systemic condition are to be specially noted. As a general rule, the prognosis of xanthoma is good as to life. The genesis of his case seems to depend on the renal involvement with an inactivity of the liver and anemia, as evidenced by the examination of the urine and blood. The stunted growth would seem to be only an interesting coincidence.

The evolution was rather rapid considering the large amount of surface covered. In this respect it somewhat approximates the cases observed by Korach and Hertzka. Its progress was accompanied by much itching. This symptom is more particularly distinctive of diabetic xanthoma, but no sugar was found in the patient's urine.

The pain elicited by contact with the lesions was a marked feature of the case.

Cases exhibiting multiple lesions are rare. The striae so well marked on the hands are not unusual in cases affecting the palm. The case recorded by Duhot also illustrated this arrangement. In Duhot's case the generalization was very complete, but, unlike that which I here report, the face, neck and genital organs were spared.

The lesions consisted of tubercles and tumors; they were yellow and firm. The palms and phalanges were striated. Some tumors had spontaneously and entirely receded. In Duhot's case there was no diabetes, but hepatic antecedents were noted.

The coalescence of adjacent lesions was particularly observable in the case of my patient.

DISCUSSION.

DR. A. W. BRAYTON, Indianapolis—Dr. Shoemaker has emphasized the extreme rarity of these generalized cases of xanthoma which justifies him in calling attention to them. I saw a case in Indianapolis several years ago, in a woman of 39. I think one-fourth of her body was covered to the same extent and with as great a variety of lesions as in this case. I reported that case briefly in the *Indiana Medical Journal*, April, 1900. It is the second case I have had knowledge of in which the lesions were so extensive. The urine was examined in that case with negative results. The patient has passed from my observation, but I think that if she is still living she can not recover. The lesions had been developing for three years, surrounding the eyes, involving most of the face, the knees and elbows, and particularly massive on the hands and arms and below the knees. On the shins the tumors were from the size of a wheat grain to a lima bean. On the palms the growths appeared as if strips of chamois skin were set in the "lines of life." The whites of the eyes were yellow; the urine was yellow, but gave no signs of bile pigments. The liver appeared normal. There was pruritus. Crocker states that four-fifths of the generalized cases have been associated with jaundice. The disease is rare. Ten years ago less than twenty cases of the diffuse form were known.

DR. WILLIAM ALLEN PUSEY, Chicago—A very exaggerated xanthoma was shown some time ago before the Chicago Dermatological Society and which has recently been under my observation. The interesting point was that the patient, a boy about 16 years of age, was suffering from polyuria without sugar. The lesions were of the type of the ordinary xanthoma and did not present the picture of diabetic xanthoma. So far as I know no other case of xanthoma associated with diabetes insipidus has been recorded.

DR. DAVID LIEBERTHAL, Chicago—Cases with such extensive distribution are not frequent. I wish to call attention to the fact that in this case, which is naturally one of infantile xanthoma, the eyelids are affected. In 1882 a report was made before the London Pathological Society by Startin and Mackenzie, of the published cases of infantile xanthoma, eight in all, and in none were the eyelids found to be the seat of lesions. About ten months ago, through the courtesy of Dr. L. Greensfelder, I had the opportunity to demonstrate a case of infantile xanthoma before the Chicago Dermatological Society. This was not the case mentioned by Dr. Pusey. It was a boy of 10 years, physically well developed, but morally defective, a kleptomaniac. The lesions were extensive, not only in the skin and subcutaneous layer, but tumors were also found on the tendons of Achilles. Dr. Greensfelder operated on the case to ascertain to what extent the tendons were affected, and it was found that their sheaths were free, but that the tendons themselves were diffusely infiltrated by the xanthomatous process. How shall we explain the development of fat in various localities normally free from it? There seems no doubt that there is an underlying, a severe, and probably an internal cause, especially when we consider Dr. Shoemaker's case, which shows a veritable cachexia. I would like to ask Dr. Shoemaker if lesions were found in the internal organs, inasmuch as there was not only extensive affection of the skin, but also lesions of the mucous membrane of the mouth and throat.

DR. EDMUND L. COCKS, New York City—In the Skin and Cancer Hospital of New York there was a case in a young woman similar to this boy's case, only the nodules of the lips and nose were larger. That case was gone over very carefully, but we could not find anything wrong except the condition of

the skin. Dr. Bulkley used the x-rays and the nodules diminished almost one-third, although the striæ have not diminished to any great extent. I feel sure from the reports I have had of her case that she will recover.

DR. JAY F. SCHAMBERG, Philadelphia—Dr. Shoemaker's patient appears to have on the cutaneous surface every gradation of the xanthomatous process. On the eyelids are typical lesions of common xanthoma planum; on the extremities are seen characteristic lesions of xanthoma tuberosum, and on the back is an inflammatory nodule closely resembling those seen in diabetic xanthoma. The extent of the eruption and the pronounced effect on the boy's general health would suggest that he is suffering from a general disease of which the cutaneous lesions are merely conspicuous external manifestations; indeed, the clinical picture is quite analogous to a malignant internal neoplasm with multiple cutaneous metastases. What has the physical examination of the abdominal viscera disclosed, particularly in reference to involvement of the liver and spleen? It is the careful study of severe cases of this character that will doubtless throw light on the lesions of xanthoma that are so commonly seen in persons enjoying apparent health.

DR. EDMUND L. COCKS, New York City—I have carefully examined the internal viscera and find the liver is very much enlarged; in fact, we have liver dullness extending below the umbilicus up to and even beyond the fifth interspace. The spleen is also enlarged. I think Dr. Shoemaker mentions the examination of the blood, which shows marked anemia.

DR. D. W. MONTGOMERY, San Francisco—The boy's protuberant abdomen would signify that some of the abdominal viscera are affected.

DR. M. L. HEIDINGSFELD, Cincinnati—I wish to record a case which I saw in November, 1902, in Evansville, Ind., in consultation with Dr. Sidney Eichel of that city. The individual was a cachectic white boy, aged 16 years, weight 68 pounds, height 4 feet 6 inches. Three years previously a ruptured compensation from a mitral insufficiency resulted in general edema, which was relieved by treatment, and jaundice and enlarged liver, which have continued to persist. The liver at the time of consultation extended below the umbilicus and almost completely filled the right half of the abdomen. The xanthomatous eruption began, shortly after compensation was ruptured, over the extensor aspects of the extremities, and was very generally distributed over the entire body in tuberculous form, except around the knees and elbows, where the lesions were very closely aggregated together and took on the form of xanthoma planum. Urine until lately contained albumin, but at the time of examination was normal, excepting that it was reduced in amount; its specific gravity was 1.015. His sleep until lately was disturbed by frequent micturition. Appetite remained good, bowels regular, general distress slight. All organs except liver and heart apparently normal.

DR. L. DUNCAN BULKLEY, New York City—When did the first signs of this disease appear?

DR. SHOEMAKER—I do not think he gives a clear history. Frank, how long have you had this?

THE PATIENT—About ten months.

DR. WILLIAM L. BAUM, Chicago—In hearing the result of the blood examination I think it would be well if some examination of the pancreas were made in this case, especially as to his digestion of fats. In view of the fact that in some cases of polyuria there is a relationship with xanthoma would point to investigation along the avenues of the pancreatic cells which might show some relationship with the central nervous system. This looks to me more like a secondary toxic effect and it would be interesting to know what Dr. Shoemaker's record of the case might be at some future time.

DR. HENRY G. ANTHONY, Chicago—This case, although presenting unusual features, is not difficult to diagnose. The cases which present the greatest difficulty in diagnosis are those in which there is no elevation, the lesions consisting simply of yellow spots in the skin. This variety is recognized as a separate form by the *Pratique Dermatologique*. Dr. Pusey showed a case of this character before the Chicago

Dermatological Society; the spots had been present on the forehead seven years. We had a discussion in regard to the diagnosis. I thought it was xanthoma; others thought it verruca planum.

UNCINARIASIS IN THE SOUTH, WITH SPECIAL REFERENCE TO MODE OF INFECTION.*

CLAUDE A. SMITH, M.D.

Pathologist to Grady Hospital; Demonstrator of Pathology and Bacteriology, Atlanta College of Physicians and Surgeons; Professor of Pathology and Bacteriology, Southern Dental College; Pathologist to Georgia Pasteur Institute; Secretary Anatomical Board of Georgia.

ATLANTA, GA.

In this paper I shall attempt to present some additional light on the mode of infection in uncinariasis or hookworm disease. The views as presented before this Section a year ago¹ have been confirmed, not only by my own observations, but also by those of physicians in different sections, who, after reading the article in THE JOURNAL, have written to me regarding the disease as it exists in their sections.

DISTRIBUTION AND PREVALENCE OF THE DISEASE.

It has been my fortune to see cases of the disease from all of the coast states in the south, from Virginia to Texas, inclusive, with the exception of Mississippi, but reports received from that state show that it is prevalent there as well. While my observations have necessarily been confined principally to my own state, yet the evidence indicates that the disease is equally as prevalent in all our southern states. I do not believe that there is any section of any extent in these states that is free from it. It seems as if the entire country was literally saturated with it. It is found on the highlands as well as the lowlands, and on the mountains as well as on the seaboard.

RELATION BETWEEN UNCINARIASIS AND GROUND ITCH.

I have previously called attention to the close relationship which I observed to exist between uncinariasis and ground itch, presenting clinical evidence that uncinariasis is always present in all cases which have had ground itch within eight years; that uncinariasis is rarely, if ever, present in those who have not had ground itch within that time; that of members of the same family, living under exactly the same conditions, those who have had ground itch within this time also have the uncinariasis, and vice versa; also that the severity of the disease and the number of parasites are almost invariably in direct proportion to the number of attacks of ground itch.

While the principal object of this paper is to report some experiments of placing the larvæ on the skin to artificially produce the disease, I wish first to make a few general remarks regarding the symptoms and diagnosis of uncinariasis.

SYMPTOMS.

No one set of symptoms can be put down as diagnostic of uncinariasis, as they vary with the amount of infection and with different individuals. For convenience, I have classified the disease under three types:

First. Mild cases, which have had only one or two attacks of ground itch, and therefore have only a small number of the parasites of the intestines, not sufficient to produce any diminution in the corpuscles and hemoglobin.

Second. The medium type of cases, which have usually had two or more attacks of ground itch, and have a much larger number of the parasites in the intestines, sufficient to produce a reduction of the corpuscles and of the hemoglobin, but which is not perceptible on inspection of the patient, but is readily shown by the blood examination.

Third. The severe type of the disease; those cases about which so much has been said in the past, and which have had numerous attacks of ground itch, and which usually have several hundreds of the parasites in the intestines and present all the symptoms of extreme anemia, which is readily recognized without the blood examination, and which is oftentimes so great as to interfere with the development of the individual.

We can not draw a sharp line separating these three types, but I make this classification for the sake of convenience in considering the prevalence of the disease, and also to draw attention to the fact that those cases which can be suspected from the symptoms of profound anemia are not so numerous as the other two types combined.

In cases of the first or mild type, there are practically no symptoms by which the disease can be suspected. There is no interference with the appetite and digestion which might not be attributed to other causes, no reduction in the hemoglobin, and the patient feels strong and vigorous, as the body is able to nourish a few of the parasites without taxing its reserve power.

In cases of the second type, there may be various symptoms of interference with digestion, and while the patient usually says that he feels fairly well and superficial inspection may not show any decided anemia, yet the blood examination will show the reduction in the constituents of the blood.

In the third or severe type, the patient presents all the appearances of profound anemia. The skin has a yellow parchment appearance, mucous membranes are blanched, the sclera shows clear white through the conjunctiva in which no blood vessels can be made out. The face is bloated and expressionless and the skin over the exposed parts often appears to be much thickened, and feels dry and rough. The patient may be greatly emaciated, but this may be hidden by edema. The patient is very weak, and if he attempts violent exercise quickly becomes exhausted and short of breath. There is a decided systolic murmur, and the blood examination shows a decrease of from 50 per cent. to 90 per cent. in hemoglobin, and the red corpuscles may be as low as 1,000,000, or even less. The appetite is capricious, perverted, and there may be nausea and headache. The perversion of the appetite may be shown in a fondness for meal, clay, sand, or other substances.

Many physicians have written to me regarding the habit of dirt eating as being the mode of infection of the disease. When I first discovered the disease in my section, this view naturally presented itself to me, as it had been mentioned in connection with the old-world parasite, but on careful investigation I found that children were sometimes falsely accused of eating dirt simply because they presented this appearance of profound anemia which is seen in those who acknowledge the habit of dirt eating. Also I found that those

* Read at the Fifty-fifth Annual Session of the American Medical Association, in the Section on Pathology and Physiology, and approved for publication by the Executive Committee: Drs. V. C. Vaughan, Frank B. Wynn and Joseph McFarland.

1. THE JOURNAL A. M. A., Sept. 19, 1903.