too much interference in the antrum, and too little interference with sepsis. I think that simple mastoids should be healed and well, fully recovered, within—and well within—two weeks. Many of them should require but two or three dressings before dispensing with drainage for simple external protection, and it is my hope and belief that Eustachian irrigation will still further reduce the time of recovery.

Tubal irrigation is indicated, I think, in every simple mastoid operation where the tympanum and tympanic membrane are intact. Naturally, for anatomical reasons, the procedure is more difficult of accomplishment in children than in adults. However, it is of less urgency in children, as the consideration of time of recovery is usually of less importance economically.

REPORT OF CASES

Case I.—C. S. C., male, aged 20 years, was a college student. X-ray diagnosis was made of an old focal infection in the right mastoid. At operation the knee of the lateral sinus was found within about $\frac{1}{2}$ inch of the posterior wall of the external canal, and the sinus was accidentally opened early in the operation. Mastoid cells were cleaned out; antrum not definitely involved and was left undisturbed. Eustachian irrigation. On account of the opened lateral sinus only one suture, at the top of the wound, was placed. On the fourth day the wound was dressed and the sinus packing removed. No pus was present. Dressed again on the sixth day. No pus. On the eighth day dispensed with drain and inserted one suture to help close the wound. The next dressing was on the tenth day. The wound closed nicely. On the fourteenth day the wound was entirely closed and the patient was discharged and sent home to another city. Note that this was an open wound case—well in fourteen days.

Case II.—J. T. S., male, aged 48 years, was a commercial salesman. Chronic mastoiditis, right side, of fifteen years' standing, with almost constant pain and occasional discharge from the ear. Tympanic membrane intact. Excessive tenderness of the posterior canal wall. At operation a large amount of carious bone and granulations with some pus were removed, including half to two-thirds of the posterior wall. (Pathologist reported pus cells numerous, but could not differentiate the micro-organisms.) Eustachian irrigation. Dressed on third day. No pus. Healing by first intention. Dressed again on fifth day, leaving out drain. Eighth day completely healed and patient sent home. A piece of plain gauze was kept over the wound area one more day merely as a protection from cold winds.

Case III.—M. B., female, aged 19 years, was a college student. Acute exacerbation of a chronic mastoiditis, right side. Posterior canal

wall markedly bulging and extremely sensitive. Tympanic membrane was intact. At operation the mastoid was found to be very large and filled with carious bone, necrotic tissue, granulations and pus, down to the tip, which was removed. Also resected one-half or more of the posterior wall. (Pathologist reported streptococcus.) Eustachian irrigation. Dressed the wound on the fourth day. No pus. Dressed it on the sixth day. Slight bloody serous discharge. Dressed on the seventh day and found moderate sero-purulent discharge. Better on the eighth day. No discharge on the ninth day. Left out the drain. Wound was closed on the eleventh day and the patient was well to all intents and purposes. One week later symptoms of pus appeared, the wound was opened under local anesthesia (ethyl chlorid) for one-half inch at the lower end. A small amount of pus was evacuated (perhaps eight or ten drops), and a drain inserted. Pus formation ceased in three or four days and at the end of a week closure of the wound was permitted. At this writing only a few days have elapsed, but the patient is apparently entirely well. As stated above, this was a streptococcus case.

AUTHORS' ABSTRACTS

Eye, Ear, Nose and Throat

Some Observations on Astigmatism. John Green, Jr., and William F. Hardy, M.D., St. Louis, Mo. American Journal of Ophthalmology, Vol. 2, No. 1, January, 1919, p. 43.

In order to eliminate uncertainty as to what is meant by astigmatism with the rule, against the rule, and oblique, Green and Hardy propose the following definitions:

"Astigmatism with the rule, or direct astigmatism, shall include all cases with meridian of greatest curvature at any point on the arc from 46° to 134° ."

"Astigmatism against the rule, or inverse astigmatism, shall include all cases with meridian of greatest curvature at any point on the arc from 0° to 44° and from 136° to 180° ."

"Oblique astigmatism shall include all cases of astigmatism with meridian of greatest curvature exactly at 135° and 45° ."

It is stated that inverse astigmatism is productive of unusually annoying symptoms. Reference is made to a recent study by the same authors, in which, in 1024 eyes with regular astigmatism, 283, or 27.7%, were inverse. This is a larger percentage than is generally surmised.

A photograph illustrates one reason for the greater degree of asthenopia in eyes with indirect astigmatism, e. g., the greater interference with the legibility of type.

Cycloplegic measurement is advocated in presbyopes who give conflicting answers when tested without "drops."

One case reported illustrates that high unilateral regular astigmatism may be due to a birth injury.

An appropriate cylinder will often greatly improve vision and enhance the comfort of a patient whose cornea is deformed by antecedent injury or disease.

An eye with irregular astigmatism due to incipient cataract does not often tolerate a cylinder, as the lenticular astigmatism is constantly changing with the progress of the opacification.

Eye Strain as a Factor in Gastric Neuroses.

Alonzo C. McFarling, Shawnee, Okla. The Journal of the Oklahoma State Medical Association, Vol. XII, No. 1, January, 1919, p. 13.

The far-reaching effect of eye strain in the production of gastric neuroses may be more easily be appreciated by a painstaking review of the physiological characteristics of the brain, the great "clearing house" of the nervous system.

From the various cells or cell units emanate the nervous impulses which activate and control the respective functions of the organs which they supply. The functional needs of such organs registering themselves as peripheral sensations upon their respective nerve centers call forth central impulses of sufficient force or intensity to meet the requirement.

In view of the fundamental physiological characteristics of nervous tissue and the anatomical arrangement of the fibers by which Nature has provided for those important processes designated as association, concentration, inhibition, etc., we may reasonably conclude that any cell or cell unit, when acting under a degree of excitement greatly exceeding the normal, would, through the processes of association and inhibition, modify the normal activity of those cells or cell units with which they were in physiological and anatomical relationship.

Reasoning from this, we may conclude that such a relationship exists between the nucleus of the vagus supplying the stomach and the nuclei of the third, fourth and sixth cranial nerves supplying the musculature of the eyes, all of which are situated beneath the aqueduct of Sylvius in the floor of the fourth ventricle.

Remote Symptoms (Especially Nervous) Due to Tonsillar Toxicemia. W. D. Jones and H. B. Decherd, Dallas, Tex. Texas State Journal of Medicine, Vol. 14, No. 9, January, 1919, p. 303.

Foci of infection, whether they be from tonsils or not, are in many cases responsible for nervous symptoms such as chorea, neuritis, etc.

The case reported in this article was one of toxic insanity and the patient rapidly recovered after removal of the tonsils. Wassermann was negative from cerebrospinal fluid at two different examinations and the culture was negative from the cerebrospinal fluid. Examination of the tonsils unfortunately was not complete as the report shows cocci and rare bacillus present, which is meaningless from a scientific standpoint.

The patient at times would be in a profound stupor, which would be followed by delirium and occasionally hallucinations. The thick white

flake patches on the tongue and similar patches on the tonsils, which were very red and friable, presented a picture not often seen.

This patient received no treatment except removal of tonsils and after care.

Congenital Tumor of the Cornea of Doubtful Classification. H. H. Stark, El Paso, Texas. American Journal of Ophthalmology, Vol. 2, No. 1, January, 1919, p. 5.

This report concerns a tumor on the cornea of a fifteen-year-old Mexican girl about 8 mm. across the base, bluntly conical in shape, and about 5 mm. high, protruding between the lids. There was a clear ring of corneal tissue surrounding it, showing that the origin had not been at the limbus. Older persons of the family confirmed the statement that the growth had been present since birth, and also that it had increased in size in the last few years.

The tumor was removed under a general anesthetic, leaving a clear cornea except where there were slight opacities from the operation.

On sectioning, it was found to be made up of fibrous connective tissue, which had undergone hyalin degeneration, with some fat in the base of the tumor.

Sections were submitted to four pathologists, two of whom reported it to be fibroid; one, dermoid; and one agreed that dermoid would be the most likely classification. However, to be positive, he would like to find some dermal tissue or hair.

The author takes up the literature on the subject of corneal tumors and speaks of the doubtful classification of some of the cases reported, contending that no tumor should be classified as corneal which has at any point a connection with the limbus. He considers the possibility of two origins of the dermoid: the lid and the amnion, contending that from either source it should be classified as a dermoid, basing the classification on the origin rather than the histological structure.

Bitemporal Hemianopsia Due to Acute Suppuration of the Posterior Nasal Sinuses. F. A. Conlon, M.D., Lawrence, Mass. American Journal of Ophthalmology, Vol. 2, No. 2, February, 1919, p. 92.

Bitemporal hemianopsia due to sinusitis is an extremely rare occurrence, there being but four such cases in all ophthalmic literature.

In the case reported the vision, a week after the onset of eye symptoms, was reduced to the ability to count fingers at two feet.

Following the exenteration of both posterior ethmoids and sphenoids, there was an immediate improvement in vision which in two weeks increased to 20/100 in each eye.

The teaching that the commisure or chiasm rests upon the optic groove of the sphenoid bone, is wrong. In the first place, the optic groove is not large enough to hold the chiasm which measures anteriorly 7 to 8 mm., while the average measurement of the groove is but 6 mm. anteroposteriorly.