tracheotomy tubes. In cases where the tube has become blocked with mucus it has invariably been coughed up with the obstruction. Bronchopneumonia does occasionally occur after intubation, but not nearly so frequently as in tracheotomy. Hemorrhages may result, but are usually of little consequence, coming as they do from the membrane of the fauces; this accident will always be avoided by the skilled operator.*

If the tube selected should be too small, the probable result will be its non-retention, as it may be at once coughed up. If this happens it will be necessary to replace it by a larger one. There is very little danger of the tube dropping into the trachea, because its collar is of sufficient size to prevent such an accident.

Extubation is always a critical period in the treatment of a case by the O'Dwyer method. This procedure is usually more difficult than the introduction of the tube. The child is placed in the same position as for its introduction, and with the index finger of the left hand the operator guides the extractor down to the larynx, where it is felt to strike against the end of the tube. It is then moved about gently, no force being used, until it drops into the opening of the tube. The blade should then be separated and firmly held while the instrument and tube are withdrawn, special care being observed not to relax the pressure just as the tube is being turned out of the larynx, for if this is done the instrument will slip and the tube may either fall back into the larynx or be swallowed. This accident can not happen with the new instrument that I have devised, because once the handles are locked they can not be separated unless done purposely by the operator. Special care should be taken that no pressure is made on the head of the tube in attempting to introduce the extractor, for the tube might possibly be pushed below the vocal cords, an accident that has happened in a few cases. There is, however, a very simple and sometimes quite an effective method of removing the tube, which requires the use of no instruments and which was discovered in Paris by one of the house surgeons at the Hospital des Enfants Malades. This is the method of expression. It is performed thus: The nurse takes the child on her knee, the operator faces the child and with the left hand over the vertex places the right thumb on the trachea, grasping the neck with the fingers. He then simultaneously compresses the trachea backward and upward, and jerks the child's head downward into the nurse's lap, and in the majority of cases the tube drops out without further trouble. This method of expressing the tube I believe to be an unscientific, dangerous and uncalled-for process, and it should only be resorted to in emergency where any procedure to save life would be justifiable.

34 Washington Street.

FUNCTIONALLY GOOD RESULTS IN THE TREATMENT OF FRACTURES AS VIEWED BY SKIAGRAPH AND PHOTOGRAPH.

BY B. N. TORREY, M.D.
CRESTON, IOWA.

By functionally good results, I mean those cases where the patient has good use of the part that has been injured; if it be a leg, he walks without a crutch or cane: if an arm, it can be flexed, extended or rotated sufficiently for practical purposes.

In order to more fully demonstrate the difference between the skiagraph and the photograph of functionally

good results, I have secured a series of pictures representing fractures of the limbs. These have not all occurred in my practice; a few have been treated by surgeons of reputation. They were not selected with a view of doing honor to the surgeon; had they been, I would have left out the very ones most needed to demonstrate the fact that we may have marked deformity and still have a useful limb.

In offering this collection of pictures, representing the different fractures and their conditions as shown by the skiagraph and photograph, I have no apologies to offer,

believing that they show just what the surgeon of the past has done, and what the majority are doing at the present time. In the examination of these, especially

248 and 249.—Fracture of leg. Four months after injury. Functional results perfect.

the skiagraphs, the first impression will be as mine was: that there must be something radically wrong in the procedure that would permit such departure from the proper anatomic appearance of the bones. Any surgeon who will look with patient honesty for an ideal re-

*Read before the Iowa State Medical Society, Des Moines, May 16-18, 1900.

sult, among his victims of the past, may look without finding the object of his quest, if such ideal involved the notion of exact anatomic alignment of the fragments; especially will he be disappointed if he expects such perfection in cases where the fracture was oblique or the fragments separated. He will be surprised to find that each and every one will present a deformity, more or less, of some kind, when viewed by the X-ray.

The results of a transverse green-stick fracture, or in those cases where the fragments have not been displaced, will show, from mechanical reasons only, angular or rotatory deformities; but in those fractures, buried as they are in excited muscular tissues, the obtaining and maintaining of exact anatomic relations would be an accident. It will be observed, in some of the pictures presented as illustrating this paper, that the results as viewed by photographs are all that could be desired; and others, while showing some deformities, are still within the limit of what would be considered good functional results. View the same cases by the X-ray, and it will be found to present angular rotatory and the overriding deformities, or a combination of any two or more—depending to a great extent on the bone fractured, the

It would not take the search-light of an X-ray to find a marked deformity in Case No. 258, a male of 200 pounds, who has quite a limp. He plows corn and does other work pertaining to the farm. The injury was caused by getting his foot caught in a hay-press, produc-

Photograph 254.

ing a compound comminuted fracture of both bones to the extent of leaving spiculæ of bone sticking to the hard wood of the press. Only his persistent demand that it should not be amputated is the reason that he has

Photograph 254.

254.—Compound fracture of leg 8 months after injury. Functional result perfect. There is no shortening. The apparent overlapping of fragments is due to the callus.

A.—End of fragments. B.—Callus which gives appearance of overlapping.

location of the fracture, whether a leg or an arm, a simple, compound, or compound comminuted, and the cause producing it. I am convinced that a careful, comparative study of this kind, made by any surgeon on his own cases, will smash some of his idols, will be conducive to humility, and make him cautious about critical remarks concerning the work of his fellow practitioner.

I would ask especial attention to No. 254, a male, aged 30, and weighing 190 pounds. He jumped from a spring-wagon, producing a compound fracture of the tibia, and simple fracture of the fibula. At this time, eight months after injury, he walks without a crutch or cane, with a very little limp, and he gets off and on trains in the capacity of a traveling man. Represented by the photograph and examined by manipulation, one would consider his a splendid result, all things considered, and I was especially satisfied with it until it was skiagraphed.

it in the place of a wooden one. Certainly a wooden one would be more symmetrical if not so handy. This case was treated in the early days of antiseptic surgery.

No. 248, a boy, 8 years old, was injured by a horse

stepping on his leg, producing a simple fracture of both bones. As shown by the photograph and by manual examination, one would think we had an ideal result, yet by the skiagraph we find not only an angular deformity, but a partial displacement of the individual fragments.

Nos. 265 and 266 represent a fracture of the femur and humerus, following a railroad accident in which the leg was lost below the knee. This case was treated in a Kansas City hospital, and by a well-known surgeon, but under some disadvantages, as all the injuries occurred at the same time.

258.—Compound comminuted fracture of leg. Twelve years after injury. Functional result good enough that patient has but a slight limp, and does general farm work.

I would call especial attention to Nos. 290 and 291, which present a photograph and an X-ray picture of a compound fracture of the leg, treated by a surgeon in Kansas, eighteen months ago. The X-ray findings are good, the alignment being practically perfect, while on the other hand the photograph shows marked rotatory deformity. The patient has a stiff ankle and a limb that is far from being functionally perfect.

The examination of fractures and the results obtained, by means of the X-ray, have been almost entirely confined to the profession, and it is a question of very grave importance to each individual member. To-day

you are criticising my ability as a surgeon through the representation of these pictures. To-morrow the general public will expect and demand the same privilege of looking over your work, and from the same standpoint. What will their criticism be? Have they not been led to believe, when the result from external appearance seems good and they have good use of the parts, that the union of the fragments was as smooth and complete as a cabinet-maker's splice? I do not refer to those severe crushing injuries—where they could see for themselves or where the astute surgeon has explained the probable results, if not allowed to amputate, but to those of more frequent occurrence, simple fractures of the leg or arm.

Through my investigation of fractures occurring in the limbs, with the fluoroscope and skiagraph, I have come to the following conclusions: 1. That both the surgeon and the public have believed they were attaining more perfect anatomic results than it is possible to get in the great majority of cases. 2. That the surgeon

Photograph 258.

should realize and the public be taught that the getting of a good functional result without deformity, as viewed by the unaided senses, constitutes the practically ideal, and that X-ray cosmetics is a refinement beyond the reasonable hopes of this day and age. 3. That we must not expect an ideal result in oblique or any other fracture where the fragments have been separated. 4. That it would be impossible to correctly coaptate a fractured bone buried in muscular tissues, and recognize it, without the use of the fluoroscope. 5. That very few, if any, oblique fractures of the larger bones of the leg or arm are treated without shortening and other deformities. 6. That the public has the impression that all fractures are transverse and the surgeon should always procure an end-to-end approximation. 7. That angular deformities alone can be corrected or prevented by lateral splints. 8. That the only way we can be sure of a correct reduction would be under the fluoroscope, with an anesthetic and superextension, or by incision, with

mechanical fixation of the fragments. The use of the fluoroscope would be impossible in many cases, and the indiscriminate practice of incision would do more harm than good. 9. That oblique fracture of the tibia, when both bones are broken, is one of the hardest of fractures to treat without marked deformity—especially if it be

muscles are located on one side of the bones; there is an element of torsion in the force that produced the fracture. 10. That extension would give us less shorten-

265.—Fracture of humerus (Left) 14 months after injury. Functional result good.

266.—Fracture of femur 14 months after injury. Result of railroad accident. Other injuries necessitated amputation of leg. Doubtless making proper extension impossible.

Photograph 265.

Photograph 266.

compound or produced by indirect violence—from the fact that there is not room for traction; the stronger

ing, less deformity and better results in fractures of the long bones; but too much extension and too long contin-

ued, would interfere with the physiologic condition of the joints, while stricture of the tissues between joints, sufficient for traction, would interfere with the circulation, and might produce blister, ulceration and non-

except those having end-to-end approximation, and that without perceptible inconvenience or pain to the patient. 12. That a skiagraph picture may exaggerate a deformity, but it might be hard to make an interested party

290 and 291.—Compound comminuted fracture of leg, 18 months after injury. Operation to encourage union was done 5 months after injury; another 11 months after injury. Result: good *apparent* alignment as seen by X-ray, but there is marked rotary deformity, ankylosis of ankle joint and fixation of toes in position of extreme flexion.

believe it. 13. That an X-ray picture, if interpreted according to our former notion, or the ideas now held by the general public, would be unjustly exacting for legal purposes. 14. That we may expect a mechanical device

Photograph 290.

Photograph 291.

union. 11. That weight of the limb, such as would result from the too early use after fracture of the leg, would increase or might produce deformity, in all cases

for fixation independent of muscular tissues; whether it is foreshadowed by Dr. Parkhill's clamps, the peg, screw or some other contrivance, no one can tell. 15. That

every up-to-date surgeon should have access to or own an X-ray machine—a portable one would be very handy. 16. That the fractures viewed and presented in illustrating this paper, fairly present the average results obtained by the past and present methods of treating fractures. 17. That the application of the X-ray to surgery, while it presents many advantages, may also bring annoyance; but since it is here and here to stay, it behooves us to make a careful study of its findings in order that while we enjoy the advantages, we may avoid the injury that might arise from its misapplication, and prevent its being made to bear false witness through misinterpretation of its findings.

I wish to acknowledge my obligation to Dr. F. E. Sampson, of Creston, who has prepared the skiagraphs and photographs used in illustrating this paper.

ON REMOVAL OF THE CERVICAL SYMPATHETIC IN GLAUCOMA AND OPTIC NERVE ATROPHY.*

BY JAMES MOORES BALL, M.D.

Professor of Ophthalmology in the St. Louis College of Physicians and Surgeons; Oculist to St. Joseph's Ophthalmic and Surgical Sanatorium.

ST. LOUIS, MO.

I propose to consider the surgery of the cervical portion of the great sympathetic nerve in certain ocular diseases. European oculists and surgeons have performed sympathectomy for glaucoma and exophthalmic goiter. I have gone further, and in one instance removed the superior cervical ganglion for simple atrophy of the optic nerve. I have performed sympathectomy four times up to July 20, 1899. First the cases will be reported; then the conclusions will be drawn.

CASE 1.—EXCISION OF SYMPATHETIC FOR GLAUCOMA ABSOLUTUM.

Mrs. B. S., aged 36, has had pain in and around the right eye for two months, and examination showed vision in this eye reduced to light perception; tension +3, and the pupil widely dilated. The anterior chamber was shallow, the cornea cloudy and slightly anesthetic, the media slightly cloudy, still allowing the fundus to be seen. The episcleral vessels were enlarged. Circumcorneal injection was present and the optic nerve cupped. A diagnosis of chronic irritative glaucoma was made. The left eye presents immature cataract, and vision in this eye is 20/70.

Knowing of the flattering results obtained by Jonnesco and others, by excision of the superior cervical ganglion in absolute glaucoma, I explained the operation to the patient, and obtained permission to operate. On May 15, 1899, the patient was anesthetized, chloroform being employed. An incision four inches in length was made on the right side downward from the mastoid process, extending along the posterior border of the sternocleido-mastoid muscle. The external jugular vein was cut and tied. The sterno-cleido-mastoid was then separated from the trapezius muscle, and the spinal accessory nerve was cut. A deep dissection was then made, exposing the carotid sheath. This was opened to enable us to locate the pneumogastric nerve beyond question. The carotid, internal jugular vein and pneumogastric nerve were then pulled forward, enabling us to see the rectus capitis anticus major muscle, on which the superior cervical ganglion rests. Tearing through the fascia the ganglion was found and stripped. The ganglion was

then cut high up with curved scissors and all its branches severed. About one inch of the trunk of the sympathetic below the ganglion was removed. The wound was closed with interrupted sutures and the neck placed in a plaster cast. The time required for operation was fifteen minutes, and immediately after it was noticed that the right eye was suffused with tears, the right conjunctiva much injected and the right nostril moist. The intra-ocular tension was +2. The patient slept well all night, without medicine, being free from pain for the first time in over two months. Tension had steadily decreased to +1.

On May 16, slight ptosis was noticed on the right side. This symptom is yet present. On May 19 the circumcorneal injection was much less; the conjunctival hyperemia and lacrimation were still present, while the ptosis was slightly increased and tension was +1.

At the present date—July 23, 1899—this patient has no pain. The retinal arteries are increased in size. Tension is +1. Vision has increased from light perception to ability to count fingers at three feet. The conjunctival injection which followed the operation has disappeared; the optic nerve has a color more approaching the normal. The ptosis is less.

This was the first sympathectomy made in America for glaucoma.

CASE 2.—DOUBLE SYMPATHECTOMY FOR GLAUCOMA SIMPLEX.

Miss M. E., a German, aged 43, was sent to me on June 14, 1899. For two years sight had been failing, until at this time vision was as follows: R. E. = 0; L. E. = light perception. Tension was +3. Both optic nerves showed marked cupping of the disc; the vessels were pushed to the nasal side. She stated that she had never had pain in the eyes and had not consulted an ophthalmic surgeon.

I advised her to submit to an excision of the left superior cervical ganglion; she consented, and on June 15 the operation was performed by myself, assisted by Dr. E. C. Renaud, at St. Joseph's Sanatorium, in the presence of Drs. J. C. Murphy, A. R. Kieffer and S. A. Grantham. The operation was difficult owing to the abnormal position of the vagus nerve. This was outside of and external to the carotid sheath, and was much smaller than normal; it was not larger in diameter than the head of a pin. It was identified by irritating it and watching the effect on the heart. The superior cervical ganglion was removed and one-half inch of the trunk of the sympathetic below. Shortly after the operation there was lacrimation, ocular congestion and contraction of the pupil on the corresponding side. On the second day she counted fingers at 2½, and on the third at 3½ feet. Slight ptosis was present.

She left the hospital on the eighth day. At this time she counted fingers at four feet. There was only slight if any reduction of tension during the eight days she was in the hospital. In counting fingers she saw the nasal side of the retina—temporal field. I did not see her again until June 30, and she was then counting fingers at five feet. Tension on that day was normal. She had light perception in the right eye.

On July 16, I excised the right superior cervical ganglion, without difficulty, and on July 7 she counted fingers at seven feet with the left eye, and could see the hand at four inches with the right. I examined her on July 20, when vision remained the same, the tension of the right eye was +1, and of the left +2. She was well pleased to have the small amount of vision she possessed.

*Read before the Ninth International Ophthalmological Congress, at Utrecht, Aug. 17, 1899.