before he is moved into it. In all our treatment, to bring about reaction, we should stop short of producing overaction and its sequence, which is secondary hemorrhage. Whenever it becomes necessary to operate on one who is very old it is important to reinforce their vital powers by a tonic course of strychnia a week or more before the operation, as it does much more good in this way than when given on the operating table. Confinement in bed is very damaging to these old people; they ought to be propped up in bed, or removed to a chair just as soon as it is practicable. It is important that operations on old persons be as bloodless as possible.

In these conditions of post-operative shock, do we not, under some circumstances, overtreat the case? Many times, with the heart scarcely beating for the lack of blood stimulation, with respiration sighing and shallow, with patient bathed in cold perspiration, we have seen one hypodermic injection after another containing stimulating poisons given in such quick succession that we may well pause to ask ourselves the question, are we not confusing and overpowering a circulation and respiration almost extinct by the rapid successive administration of such strong drugs as nitroglycerin, strychnia, digitalin, atropin and morphin into a subject already overpowered by a general anesthetic, loss of blood, shock and collapse? Under the excitement of such conditions, proper time very often does not elapse from one dose to another to see what the effect will be. Again, drugs having diametrically opposite effects are often used and stored up in the system. We are not claiming this to be a common practice in well-regulated operating rooms, but I am sure that many of us have witnessed cases in which the postoperative treatment was overdone.

It has always been my idea that, if surgeons would spend more time in getting the patient in proper condition to stand the operation, in having everything in readiness, in having assistants trained in what they are expected to do, and also if the least possible time were consumed in performing the operation, and if all lectures and explanations were suspended until the operation is finished, much less of the first necessary poison, viz., the anesthetic, would be required, and we would have fewer cases in *extremis* to deal with after operations.

THE MECHANISM OF STREPTOCOCCUS INFECTION.*

GUSTAV F. RUEDIGER, M.D.

[From the Memorial Institute for Infectious Diseases.]

CHICAGO.

[Work aided by a grant from the American Medical Association.]

INTRODUCTION.

Although a great deal of work has been done on the mechanism of streptococcus infection, both with normal and with immunized animals, many points are still left entirely in the dark. In the experiments described in this paper, an attempt is made to throw additional light on some of these points, but on account of the many inherent difficulties, the results which I have achieved by no means exhaust the problem.

Shortly after the publication of his classical paper on phagocytosis in daphnia, Metchnikoff¹ made a study

of phagocytosis of streptococci in the human body, in cases of erysipelas. He concluded from his observations that streptococci gain entrance through abrasions of the skin, multiply and set up an inflammation. At the same time there is a gathering of microcytes about the streptococci, and these soon take up the latter and destroy them. Macrocytes also are found in considerable numbers, but they do not take up the streptococci, but have, nevertheless, a phagocytic action, in that they take up and remove dead and disabled microcytes.

In 1895 Denys and Leclef² studied the mechanism of immunity in rabbits which had been repeatedly injected with small but gradually increased doses of virulent streptococci. Their conclusions may be briefly stated as follows: Immune rabbit serum is not as good a culture medium for streptococci as normal serum, but it does not possess any marked streptococcidal powers. The cell-free fluid from a leucocytic exudate retards multiplication of virulent streptococci and sometimes kills them. Normal leucocytes in normal serum, and leucocytes from an immune rabbit suspended in normal serum, do not greatly retard multiplication of virulent streptococci. Leucocytes from a normal rabbit, or those from an immune rabbit, suspended in immune rabbit serum rapidly take up and destroy virulent streptococci. The serum has acquired something in the process of immunization which neutralized something in the cocci, by virtue of which they were protected against phagocytosis.

Denys and Marchand³ showed that there is better phagocytosis of virulent streptococci when inoculated into a suspension of rabbit's leucocytes in normal rabbit serum, to which had been added 1 per cent. of immune horse serum, than when they were inoculated into the same mixture of leucocytes and rabbit serum, with the addition of 1 per cent. of normal horse serum.

Bordet⁴ was able to find no protection against streptococci in normal and in immunized rabbits and guinea-pigs, except that due to phagocytosis. If a rabbit was treated with antistreptococcal serum, and later injected with several times the minimum fatal dose of streptococci, the organisms were all taken up by phagocytes and destroyed. In untreated rabbits he also found phagocytosis, but the organisms soon got the upper hand and the rabbit died of streptococcus infection.

Marchand⁵ studied phagocytosis of virulent and non-virulent streptococci, and came to the conclusion that the failure on the part of the leucocytes to take up virulent streptococci depends on a physical property of the organisms and not on a secretion. Tchistovitch⁶ injected rabbits intravenously with fatal doses of a highly virulent streptococcus, killed the animals in one-fourth to six hours, and examined the organs for evidence of phagocytosis. He was always able to find some phagocytosis in the lungs and in the liver of these animals, and concluded that this phagocytosis of virulent streptococci may be due to the fact that every culture contains some cocci which are less resistant than others, and that those are the only ones which are taken up by the phagocytes.

Simon⁷ found very little evidence of phagocytosis, *in vitro*, when rabbit's leucocytic exudate was inoculated with a non-virulent streptococcus. The washed

2. La Cellule, 1895, vol. xi, p. 177.

3. Quoted from Denys, Centralbl. f. Bakt., 1898, vol. xxiv, p. 685.

4. Annales de l'Institut Pasteur, 1896, vol. x, p. 104, and 1897, vol. xi, p. 177.

5. Archiv de Méd. Exper., 1898, vol. x, p. 253.

6. Annales de l'Institut Pasteur, 1900, vol. xiv, p. 802.

7. Centralbl. f. Bakt., 1901, vol. xxix, pp. 81 and 113.

* Read in the Section on Pathology and Physiology of the American Medical Association, at the Fifty-fifth Annual Session, June, 1904.

1. Virchow's Archiv, 1887, vol. cvii, p. 209.

leucocytes suspended in salt solution, and also the cell-free exudate, killed non-virulent streptococci, but not the virulent organisms. Leucocytic exudate mixed with rabbit serum had no effect on non-virulent streptococci. He was unable to confirm the view advanced by Bordet and others that a rabbit which is injected with streptococci into a pleural cavity containing leucocytic exudate can stand a larger dose than can a rabbit of the same size which is injected in the normal pleura. The pleural exudate was produced by the injection of aleuronat suspension forty-eight hours previously. These results are quite contradictory to those of other investigators, and it was, therefore, thought desirable to repeat these experiments with leucocytes.

EXPERIMENTS WITH LEUCOCYTIC EXUDATE IN VITRO.

A large Belgian hare was injected in each pleural cavity with 10 c.c. of aleuronat suspension, and bled to death in from eight to ten hours later. About 30 c.c. of thick exudate, containing chiefly polymorphonuclear leucocytes, was obtained and used in the experiment which is tabulated in Table 1. No sodium oxalate solution was added to this exudate, as this salt is strongly streptococidal, but the coagulation of the exudate was prevented by occasional agitation of the tubes during the first hour. The contents of each tube were 2 c.c., which were inoculated with three loopfuls of a twenty-four-hour broth streptococcus culture, and three loopfuls were plated at intervals. Tube 8 differs from Tube 7, in that it contains the washed leucocytes which had been centrifugated out of 2 c.c. of the exudate, suspended in rabbit serum, whereas Tube 7 contained a mixture of 1 c.c. of the entire exudate +1 c.c. of serum. The fluid of the exudate in Tube 8 was removed to show that the effect of these mixtures is not due entirely to the liquid portion of the exudate, but in a large measure to the leucocytes. The table shows that non-virulent streptococci are destroyed by the leucocytic exudate, by mixtures of leucocytes and serum, by leucocytes and defibrinated blood, and by the cell-free exudate. Defibrinated blood and serum alone, the heated exudate and a suspension of leucocytes in salt solution have little or no effect on these organisms. The virulent streptococcus is not destroyed by the leucocytic exudate. Smears made from these tubes in from one to two hours showed evidence of phagocytosis, but this is most easily demonstrated in tubes containing suspensions of leucocytes in serum. Table 2 shows the same experiment as that shown in Table 1, except that guinea-pig's leucocytes and blood were used. With these leucocytes the eighteen-hour plates are often sterile, or nearly so, while it is only occasionally that they are sterile when rabbit's leucocytes are used.

TABLE 1.—Showing the effect of rabbit's leucocytic exudate and mixtures of exudate and body fluids on streptococci.

Streptococci.	Immed-ately.	3 hrs.	18 hrs.
300 Leucocytic exudate	4600	0	10000
300 Exudate 58° for half hour.....	3250	8000	Many
300 Cell-free exudate	4800	1400	3000
300 Defibrinated blood	4800	4700	Many
300 Blood 1 c.c. + exudate 1 c.c.....	4000	750	1500
300 Rabbit serum	4200	4200	Many
300 Serum 1 c.c. + exudate 1 c.c.....	5200	1100	Many
300 Leucocytes in serum	5000	900	Many
300 Leucocytes in NaCl sol.	4900	5000	Many
324 Leucocytic exudate	420	11	Many
324 Defibrinated blood	430	7000	Many
324 Blood 1 c.c. + exudate 1 c.c.....	750	540	Many
324 Rabbit serum	552	6500	14000
324 Serum 1 c.c. + exudate 1 c.c.....	320	1	Many
B104 Leucocytic exudate	3500	3600	Many
270 Leucocytic exudate	1700	2	Many

Sources of streptococci: Nos. 300 and 324 from the heart's blood of scarlet fever patients, postmortem. No. B104 from an abscess in a guinea-pig which had been injected with a fungus. This organism has been passed through 78 rabbits and is very virulent for both rabbits and guinea-pigs.

TABLE 2.—Showing the effect of guinea-pig's leucocytic exudate and mixtures of exudate and body fluids on streptococci.

Streptococci.	Immed-ately.	3 hrs.	18 hrs.
300 Leucocytic exudate	6500	70	170
A Leucocytic exudate	650	600	Many
B104 Leucocytic exudate	8000	10000	Many
300 Cell-free exudate	5500	800	6000
B104 Cell-free exudate	8000	10000	Many
300 Defibrinated blood	6000	5000	Many
300 Defibrinated blood + leucocytes...	6800	240	72
300 Serum	6000	6000	Many
300 Serum + leucocytes	6000	2000	Many

Streptococcus No. 300 is not virulent for guinea-pigs. Nos. B104 and A are virulent. No. A was isolated from a peritonsillar abscess following an attack of follicular tonsillitis and was used in this experiment in the first generation on artificial media.

EXPERIMENTS WITH LEUCOCYTIC EXUDATE IN VIVO.

In order to determine whether a rabbit is more resistant when injected in a pleura containing leucocytic exudate than when injected in the normal pleura, the following experiments were carried out:

EXPERIMENT A.—A gray rabbit weighing 1,450 grams was injected into the right pleura with 7 c.c. of aleuronat suspension. Fifteen hours later 2 c.c. of a 40-hour culture of a virulent streptococcus were injected into this pleura and also into the right pleura of a normal rabbit weighing 1,500 grams. The normal rabbit was found dead in 18 hours, and the one which had been previously injected with aleuronat died 45 hours later.

EXPERIMENT B.—A gray rabbit weighing 2,100 grams was injected in the right pleura with 6 c.c. of aleuronat suspension. Eighteen hours later 1 c.c. of a 24-hour culture of a virulent streptococcus was injected into this pleura and also into the right pleura of a normal gray rabbit weighing 2,150 grams. The normal rabbit died in about 18 hours, while the aleuronat rabbit died only after 45 hours—21 hours after the death of the control.

The experiment was now varied so that the conditions were the same as in Simon's experiments.

EXPERIMENT C.—A rabbit weighing 1,700 grams was injected in the right pleura with 8 c.c. of aleuronat suspension, and 48 hours later this rabbit and a normal rabbit weighing 1,700 grams were each injected in the right pleura with 1 c.c. of a 30-hour streptococcus culture. In this experiment the aleuronat rabbit died in 24 hours, while the control lived for five days.

EXPERIMENT D.—A rabbit weighing 1,800 grams was injected in the right pleura with 9 c.c. of aleuronat suspension. Forty-eight hours later this rabbit and a normal rabbit weighing 1,800 grams were each injected in the right pleura with 1.5 c.c. of a 24-hour streptococcus culture. The normal rabbit died in 37 hours, while the aleuronat rabbit died in 45 hours—eight hours after the death of the control.

The results of the last two experiments agree fairly well with those obtained by Simon, and we may now ask why they are different from the results of Experiments A and B. Apparently, this difference is due to the fact that in Experiments C and D the inoculation was made some thirty hours later than in Experiments A and B.

That this is the correct explanation is indicated by the following experiment:

EXPERIMENT E.—A rabbit weighing 2,300 grams was injected in the right pleura with 9 c.c. of aleuronat suspension. Forty-eight hours after this injection the rabbit was bled to death, and on opening the pleural cavity it was found to contain no leucocytic exudate.

Had this rabbit been inoculated in the right pleura forty-eight hours after the injection of the aleuronat suspension, the inoculation would not have been made into leucocytic exudate. That being the case we would not expect that this rabbit could stand a larger dose of streptococcus than a normal rabbit. In Simon's experiments, however, the inoculations into the pleural cavity were made forty-eight hours after the injection of aleuronat, and this is probably the reason why he did

not find these rabbits more resistant than normal rabbits. Had he made his inoculations within twenty-four hours after the injection of aleuronat his results would very probably have been different.

EXPERIMENTS WITH ORGAN CELLS.

It seems quite reasonable to suppose that the organ cells might play an important part in the protection of the body against invasion by streptococci, and with this point in mind the following experiment was carried out:

EXPERIMENT.—Two guinea-pigs, A and B, were bled to death and the blood collected under aseptic precautions. The organs were now carefully removed from the body, placed in sterile glass dishes and rubbed up; A with guinea-pig's serum, and B with defibrinated guinea-pig's blood. The organs and blood were not allowed to become cold. The suspensions of organ cells were now introduced into small test tubes, inoculated with two loopfuls of a 24-hour broth culture of a non-virulent streptococcus, and two loopfuls were plated at intervals. The colonies that developed in each plate were carefully estimated and the results are shown in Table 3. In no instance did the suspension of organ cells have an inhibitive effect on the multiplication of the streptococci, but on the contrary the organisms multiplied more rapidly in these suspensions than in either the serum or the defibrinated blood alone. If, however, bone marrow was added to the defibrinated blood or serum, the number of streptococci in these tubes was greatly diminished during the first three to five hours.

TABLE 3.—Showing the effect of suspensions of organ cells of guinea-pigs on streptococci.

Streptococcus.	Immediately.	3 hrs.	18 hrs.
B104† Defibrinated blood	1800	1600	1500
B104 Blood + muscle	2000	8000	Many
B104 Blood + spleen	1900	Many	Many
B104 Blood + liver	1800	Many	Many
B104 Blood + kidney	1900	Many	Many
B104 Blood + adrenal	1500	Many	Many
B104 Blood + lymph gland	1500	6000	Many
300 Blood + bone marrow	2500	1000	Many
300 Serum	3800	3750	6000
300 Serum + leucocytes	2700	1800	1500
300 Serum + spleen	2800	5000	Many
300 Serum + liver	2000	12000	Many
300 Serum + kidney	2400	12000	Many
Control liver + serum	0	3	9
Control spleen + serum	0	0	0

†The strain of streptococcus B104 used in this experiment has not been passed through rabbits, and has lost its virulence.

These results agree with those found by Bail and Peterson⁸ in their study of the mechanism of anthrax immunity in chickens, and show that suspensions of organ cells from the guinea-pig possess no streptococidal power except the suspensions of bone marrow. Suspensions of marrow cells in defibrinated blood or serum kill large numbers of avirulent streptococci, but that is not the case with suspensions in salt solution. These results have also been obtained with suspensions of rabbit's bone marrow.

PHAGOCYTOSIS IN HUMAN INFECTIONS.

During the course of this work I had an opportunity to study a case of acute purulent rhinitis, in the discharge of which streptococci were found in pure culture.

Patient.—A man 27 years of age. During the first two days the discharge was clear and watery and contained very many streptococci in short chains and in diplococci. Figure 1 shows a photomicrograph of a smear made on the second day. There were only a few leucocytes, and those did not contain any streptococci. On the fourth day the discharge had become thick and purulent, and smears made at that time presented an entirely different picture. There were no free streptococci, but very many leucocytes, some of which were loaded with streptococci. Figure 2 shows a photomicrograph of such a prepara-

tion made on the fourth day. The purulent discharge continued for about a week, and during this time it was easy to find leucocytes in the smears which contained large numbers of streptococci. The streptococci finally disappeared, and the patient made a complete recovery.

In this case of streptococcus infection, phagocytosis of the organisms by the leucocytes seemed to be an important factor in combating the infection. In the smears made on the fourth day no free streptococci could be found, although diligent search was made for them. Plates made with a small quantity of discharge, which was obtained high up in the nares by means of a sterile cotton swab, contained, however, many colonies of streptococci after twenty-four hours' incubation. The fact that many colonies developed in the plates, although no free streptococci could be found in the smears, seems to indicate that the organisms were not dead when they were taken up by the phagocytes, but were taken up alive. It is possible, however, that there were free streptococci in this purulent discharge, even though they were not found in the smears.

PHAGOCYTOSIS OF LIVING STREPTOCOCCI IN VIVO.

The following experiment was, therefore, made to show more conclusively that there is phagocytosis of living streptococci in the animal body:

EXPERIMENT.—A guinea-pig was injected in the right pleura with 7 c.c. of aleuronat suspension. Twelve hours later he was injected in this pleura with 2 c.c. of an 18-hour glucose broth culture of a non-virulent streptococcus, and two hours after this injection he was bled to death. Several cubic centimeters of leucocytic exudate were now taken from the right pleura, under aseptic precautions. Smears made with this exudate contained no free streptococci, but many of the leucocytes contained cocci in large numbers.

Figure 3 is a photomicrograph of one of these smears showing streptococci in the body of a leucocyte.

Preparation of Slides.—By means of a sterile, soft cotton swab a small quantity of the exudate was now gently smeared on each of eight agar-covered slides; the slides were incubated in a moist chamber for from four to six hours, stained and examined under the microscope. The preparation of these slides requires the following special technic: Three ordinary clean glass slides and a heavy watch glass are placed in a large Petri dish and sterilized in dry heat. When the Petri dish has cooled sufficiently to be handled the slides are taken out with sterile forceps and hot glucose agar is poured on each to form a uniform thin film. Each slide is immediately replaced in the Petri dish and the agar allowed to solidify. The exudate which contains many streptococci in the bodies of the leucocytes is now gently smeared on these agar films, the watch glass is half filled with sterile water, and the large dish is placed in the incubator for from four to six hours. The water is used to prevent drying of the agar.

Staining.—The slides are now taken out of the Petri dish and dried in a thermostat at 55 C. for from fifteen to twenty minutes, or longer if necessary; fixed in 40 per cent. formalin for four or five minutes, washed in water, stained for only a few seconds with Loeffler's methylene blue solution, again washed in water and dried in the thermostat.

The best stains were obtained when the methylene blue solution was poured over the wet slide, the slide being held so that the stain ran off readily.

The slide is immediately placed in a dish of cold water and gently rinsed in several changes of water. Violent agitation in the water pulls off the film of agar. When the agar is perfectly dry (which requires only about half an hour), a cover-glass is placed on it, either in glycerin or Canada balsam. The slides are now ready to be examined with the oil immersion lens.

Examination.—Examination in various stages shows positively that the streptococci in the leucocytes multiply, burst open the cell and form a small colony surrounding the cell

which contained them. Figures 4 and 5 are photomicrographs of such preparations at different stages. In Figure 4 the leucocyte has just been burst open and the streptococci are beginning to grow out of it; in Figure 5 a considerable colony has already formed around the ruptured leucocyte. These colonies may be found in all stages, by varying the time of incubation.

The leucocytes do, however, kill streptococci if the conditions are right, as is shown by the following two facts: 1. Organism No. 300 is not virulent for guinea-pigs, and we have seen that the leucocytes take up the cocci alive. The guinea-pig must destroy these organisms somewhere, and we can find no substance in its body that is capable of doing this except the leucocytes and the bone marrow. We have seen in Table 2 that the leucocytes destroy streptococci *in vitro*, and the inference is that they also destroy them *in vivo*. 2. Plates made with three loopfuls of the exudate when taken from the guinea-pig contained 2,540 colonies of streptococci after twenty-four hours' incubation, while plates made with three loopfuls, after the exudate has stood in a test tube in the incubator for six hours, contained only 480 colonies. Evidently some of the streptococci

to have obtained powerful toxins. Marmier¹³ got a powerful toxin by growing streptococci in peptone solution. He precipitated the toxin from the filtered cultures, and found that the precipitate killed rabbits in doses of 40 mg. Schenk¹⁴ extracted the livers of rabbits which had died of streptococcus infection, with salt solution and filtered the extract. These filtrates and also those of a broth culture of his organism were toxic for mice. Homén and Laitinen¹⁵ prepared a toxin from broth cultures by precipitation with ammonium sulphate. These cultures varied in age from three to thirty days, and apparently were not filtered, hence it is doubtful whether the precipitates were sterile. Lingelsheim¹⁶ found that some strains of streptococci possess a high toxicogenic power, but that they usually lose that property after being grown on artificial media, even if they are passed through rabbits from time to time. He obtained his most powerful toxin from an organism of medium virulence.

Besredka¹⁷ succeeded in preparing filtrates that are strongly hemolytic for red corpuscles of various animals, by growing a virulent streptococcus in heated rab-

Fig. 1.—Photomicrograph $\times 1500$. Streptococci in nasal discharge on second day of sickness.

Fig. 2.—Photomicrograph $\times 1500$. Streptococci in a leucocyte from nasal discharge on fourth day of sickness.

were killed in this tube, but the smears show that all of the organisms are in the leucocytes, hence they must have been killed in the leucocytes.

THE SO-CALLED STREPTOCOCCUS TOXINS.

The so-called toxin-producing power of streptococci has been studied by a number of investigators.† In 1888 Manfredi and Traversa⁹ succeeded in preparing a toxic filtrate from a streptococcus which had been isolated from a case of erysipelas. Four years later Roger¹⁰ prepared a filtrate which killed rabbits in doses of 13 to 20 c.c. per kilogram weight of rabbit. Marmorek¹¹ got a more powerful toxin by growing a virulent streptococcus in a mixture of bouillon and human serum for three months and filtering. One cubic centimeter of this filtrate killed a large rabbit in three to four days. More recently he¹² recommends a special culture medium containing glyocol and leucin, in which he claims

bit serum for eighteen to twenty-four hours and filtering. His filtrates were not toxic for rabbits, but I have shown¹⁸ that filtrates thus prepared may also be toxic. Simon¹⁹ has pointed out recently that the hemolysin and the toxin are probably two different substances, and this may explain why my results do not agree with those of Besredka.

Arohnson²⁰ found that the filtered broth cultures, varying in age from two to fourteen days, possessed only weak toxic powers, even if the organism was highly virulent. Macfadyen and Rowland²¹ prepared a highly toxic extract of streptococci by grinding up the organisms when frozen and reduced to a very low temperature by means of liquid air, and exhausting this mass with salt solution. One-tenth c.c. of this extract killed a guinea-pig in six hours when injected intraperitoneally.

†It should be remembered that it has not yet been shown that any of the so-called streptococcus toxins correspond fully with the dogmatic definition now given of toxin.

9. Quoted from Lingelsheim, *Beit. z. Exp. Ther.*, No. 1, p. 16.

10. *Rev. de Méd.*, 1892, vol. xii, p. 944.

11. *Annales de l'Institut Pasteur*, 1895, vol. ix, p. 611.

12. *Berliner klin. Woch.*, 1902, vol. xxxix, p. 253.

13. *Annales de l'Institut Pasteur*, 1895, vol. ix, p. 568.

14. *Wien. klin. Woch.*, 1897, vol. x, p. 937.

15. *Ziegler's Beiträge*, 1899, vol. xxv, p. 10.

16. *Beiträge zur Experimentellen Therapie*, No. 1, 1899.

17. *Annales de l'Institut Pasteur*, 1901, vol. xv, p. 880.

18. *THE JOURNAL A. M. A.*, 1903, vol. xli, p. 962.

19. *Centralbl. f. Bakt.*, 1903-4, vol. xxv, pp. 308-440.

20. *Berl. klin. Woch.*, 1902, vol. xxix, p. 981.

21. *Centralbl. f. Bakt.*, 1904, vol. xxxv, p. 415.

Simon has shown that streptococci secrete more toxin when grown in a medium which has some resistance to their development than when grown in a medium which is favorable for their rapid growth. He therefore did not use broth or serum as a culture medium, but used the cell-free fluid from rabbit's leucocytic exudate. The exudate was obtained by injecting the rabbit in the pleura with 6 to 10 c.c. of aleuronat suspension and bleeding it to death the following day. He grew an organism of medium virulence in this fluid for from three to nine days under anaërobic conditions, and passed the culture through a filter. He also prepared a toxin by injecting into the pleural exudate a virulent streptococcus and filtering the fluid found in the pleura after the rabbit had died from the infection. These filtrates killed rabbits in from three to twelve days, in doses of from 4 c.c. to 7 c.c.

At the time of publication of Simon's paper I was working along the same line, and have since been able to confirm his results, as is shown by the following experiments:

EXPERIMENT A.—A mixture of two parts of cell-free exudate and one part of broth was inoculated with a virulent streptococcus (B. 104), incubated for three days and passed

Fig. 3.—Photomicrograph $\times 1500$. Streptococci in guinea-pig's leucocytes, taken from the pleura two hours after injection of streptococci.

through a porcelain filter. One c.c. of the filtrate added to a tube of ascites broth gave no growth in 24 hours. A rabbit weighing 1,335 grams was injected with 10 c.c. of the filtrate. During the first two days the rabbit lost 50 grams and then held his weight, but died on the fourteenth day.

EXPERIMENT B.—A rabbit was injected with 4 c.c. of aleuronat suspension in each pleura, and the following morning he was injected in each pleura with 3 c.c. of a 24-hour culture of a virulent streptococcus. The rabbit was found dead the following morning and 25 c.c. of exudate was obtained from the pleural cavities. This exudate was centrifugated and passed through a filter. One c.c. of the filtrate added to a tube of ascites broth gave no growth. A rabbit weighing 1,690 grams was now injected with 10 c.c. of the filtrate. The rabbit appeared to be ill the next day, but did not die until the end of the fifth day. Cultures made from the heart's blood, liver and spleen remained sterile.

Several other experiments similar to these were carried out, but not all of the animals died from the injection of the filtrate. It is clear, however, that filtrates thus prepared possess a considerable degree of toxicity.

At the suggestion of Professor Hektoen, I tested some of the filtrates to determine whether they are toxic for leucocytes, as has been shown by Van de Velde,²² and by Neisser and Wechsberg,²³ to be the case with staphylococcus filtrates. The organism used in these experiments is B. 104,¹⁸ which has been passed through 78 rabbits, and is very virulent for rabbits and for guinea-pigs, and produces an active hemolysin when grown in heated rabbit serum or in a mixture of equal parts of rabbit serum and ascites fluid. The filtrates used were from twenty-four to thirty-six-hour cultures in a mixture of rabbit serum and ascites fluid, which had been heated to 56 C. for one-half hour before inoculation.

Neisser and Wechsberg²³ have shown that living leucocytes reduce methylene blue solution when the supply of oxygen is shut off, while dead leucocytes no longer possess this property. In my experiments this fact was made use of to determine whether or not the leucocytes had been killed by the filtrate; 0.3 c.c. to 0.4 c.c. of leucocytic exudate from a rabbit was added to 1.5 c.c. of the filtrate, incubated for two hours, then two drops of a methylene blue solution added and the con-

Fig. 4.—Photomicrograph $\times 1500$. Multiplication of streptococci after their ingestion (*in vivo*) by a guinea-pig's leucocyte.

tents of the tube shaken up and covered with a layer of sterile olive oil. If there was reduction in two hours the result was considered negative, but if there was no reduction in that time this fact was taken as evidence that the leucocytes were dead. Control tubes with a mixture of equal parts of heated rabbit serum and ascites fluid were always made, and the results recorded only when there was marked reduction in the control in two hours. A large number of these experiments was carried out, but it is not necessary to tabulate more than one of them here.

Experiment.—3 c.c. exudate + 1.5 c.c. 24-hour filtrate; no reduction in 2 hours, but a trace of reduction in 3 hours.
3 c.c. exudate + 1.5 c.c. 36-hour filtrate; no reduction in 3 hours.
3 c.c. exudate + 1.5 c.c. ascites-serum;²⁴ nearly complete reduction in 2 hours.

The leucocytic exudate was obtained from the pleura of a rabbit which had been previously injected with aleuronat suspension. The methylene blue solution was

22. Annales de l'Institut Pasteur, 1896, vol. x, p. 580.

23. Zeitschr. f. Hyg., 1901, vol. xxxvi, p. 299.

24. The term "ascites-serum" as used here refers to a mixture of equal parts of ascites fluid and rabbit serum, heated to 56 C. for one-half hour.

made according to the formula given by Neisser and Wechsberg, which is given below:

Solution A.—Methylene blue, 1; absolute alcohol, 20; distilled water, 29.

Solution B.—1 c.c. of solution A; 49 c.c. of physiologic salt solution.

Two drops of Solution B, which must be sterile, are added to each tube in the experiments.

The fact that the filtrates have a decided effect on leucocytes has also been shown in a second experiment.

EXPERIMENT.—Two cubic centimeters of a rich leucocytic exudate were placed in each of five small test tubes and centrifugated. The clear, supernatant fluid was now decanted from each tube and replaced by 2 c.c. of bouillon, heated ascites-serum or filtrate from a 24-hour ascites-serum streptococcus culture. The leucocytes in each tube were evenly suspended in the fluid, the tubes placed in a water bath at 37 C. and kept there for from 1 to 1½ hours. Now .3 c.c. of normal rabbit serum (guinea-pig serum when guinea-pig's leucocytes were used) was added to each tube, the tubes inoculated with a loopful of a non-virulent streptococcus broth culture and plates made at intervals, the tubes being kept in the incubator.

The following table shows the results of an experiment:

Fig. 5.—Photomicrograph $\times 1500$. Multiplication of streptococci after their ingestion (*in vivo*) by a guinea-pig's leucocyte, in slightly later stage than Figure 4.

TABLE 4.

	Im.	4-5 hrs.	20 hrs.
Leucocytes + 2 c.c. filtrate B 104 + .3 c.c. normal serum	2000	3200	Many
Leucocytes + 2 c.c. filtrate 10 A + .3 c.c. normal serum	1200	2000	Many
Leucocytes + 2 c.c. filtrate 300 + .3 c.c. normal serum	1100	24	Many
Leucocytes + 2 c.c. ascites serum + .3 c.c. normal serum	1000	140	Many
Leucocytes + c.c. bouillon + .3 c.c. normal serum	1800	60	Many

†The numbers B 104, 10 A and 300 refer to different strains of streptococci. B 104 has a high virulence and 10 A a medium virulence; both these strains produce hemolysin in heated serum. Although 10 A has never been passed through animals. It was recently isolated from suppurating axillary glands. 300 is non-virulent, and does not produce hemolysin.

The suspensions of leucocytes in filtrates from serum cultures of virulent streptococci have no effect on streptococci, while the suspensions in filtrates from non-virulent organisms, in heated ascites serum, and in bouillon destroy large numbers of cocci.

These experiments do not always give positive results, but there is a small percentage of tubes in which the filtrate has apparently no effect on the leucocytes. This is not surprising, however, when we consider the fact that in about 100 filtrates which I have tested for

hemolysis, there were nearly 10 per cent. that had no hemolytic properties, whereas some of them were so active that .005 c.c. completely laked 1 c.c. of a 2½ per cent. suspension of rabbit's washed corpuscles in two hours.

In all these experiments with leucocytes it is necessary that no time should be lost after the exudate has been removed from the animal. It has been my custom to have all tubes ready and at incubator temperature before the exudate is taken from the animal, so that it can be immediately introduced into the warm tubes containing the serum or filtrates.

The addition of normal serum to the tubes in the last experiment was carried out because it was found that there is less destruction of streptococci by leucocytes when suspended in heated serum than when they are suspended in fresh normal serum. This is shown by the following experiment:

	Im.	3 hrs.	20 hrs.
Rabbit's leucocytes + 1 c.c. normal rabbit serum	750	17	Many
Rabbit's leucocytes + 1 c.c. rabbit serum 60 C. for one-half hour	700	320	Many

The leucocytes contained in each tube had been centrifugated out of 1 c.c. of exudate and were washed three times in .85 per cent. NaCl solution which had been warmed to 38 C.

This result agrees with those of Wright and Douglas,²⁵ who have shown that human washed leucocytes when suspended in heated serum take up less streptococci, colon bacilli and other organisms than when they are suspended in normal human serum.

Attempts have been made to determine whether the destruction of streptococci by the suspension of leucocytes in serum is a complement action, but these results are as yet unsatisfactory. It is evident that the rôle of the plasma in phagocytosis of streptococci needs to be further investigated.

SUMMARY.

It will be seen that my work corroborates and extends the work of most previous workers in this field.

Leucocytic exudate, the cell-free exudate and suspensions of leucocytes or of bone marrow in serum or in defibrinated blood kill non-virulent, but not virulent, streptococci. Suspensions of organ cells from guinea-pigs, in serum or in defibrinated blood, do not kill streptococci. Leucocytes in the animal body take up living streptococci.

Putting all facts together, it seems that we are warranted in concluding that the leucocytes and the bone marrow are the most important (if not the only) factors concerned in combating streptococcus infections in rabbits and in guinea-pigs. Dr. Weaver and I²⁶ have shown that normal human serum is a good culture medium for streptococci, and I have shown above that there was active phagocytosis of streptococci by the leucocytes in a case of streptococcus infection in man. It would seem, therefore, that in man also the leucocytes are an important factor in protecting the body against invasion by streptococci.

It was shown in the foregoing that the leucocytes take up living streptococci *in vivo*, and in all probability destroy them, but the fact that the cell-free exudate also destroys many organisms indicates that there may be destruction of cocci extracellularly as well as intracellularly.

Virulent streptococci when grown in heated serum

25. Proceedings of the Royal Society, 1903, vol. lxxii, p. 357, and 1904, vol. lxxiii, p. 128.

26. Trans. Chicago Path. Soc., 1903, vol. v, p. 285; Medicine, 1903, vol. ix, p. 515.

secrete a substance which is toxic for leucocytes. This may throw some light on the question why virulent streptococci are not taken up by the phagocytes. The practical value of this demonstration is self-evident. If we accept the view that the leucocytes are the most important agent in combating streptococcus infections, we see at once that the destruction of these cells by the toxin, as they collect about the focus of infection, robs the body of its strongest defense against these organisms.

In conclusion, I wish to thank Professor Hektoen and Dr. Weaver for many suggestions.

ADDENDUM: Since this article was put in type there has appeared an important paper on experimental streptococcus infections by Neufeld and Rimpau.²⁷ These experimenters have shown that the leucocytes play an important rôle in combating streptococcus infections in rabbits. They have shown also that the addition of antistreptococcus serum to a suspension of leucocytes and streptococci aids phagocytosis, and that this is not due to a stimulation of the phagocytes but to an effect of the serum on the cocci.

CYSTS OF BARTHOLIN'S GLANDS,

WITH BRIEF REMARKS ON THE ANATOMY OF THE NORMAL GLAND STRUCTURE.*

THOMAS S. CULLEN, M.D.

BALTIMORE, MD.

Bartholin's gland was named after Bartholinus, who described it in detail in the seventeenth century. In 1761 Duverney¹ gave in detail the results of further studies of the gland, and pointed out the resemblance between its tissue and that of the prostate gland; he also mentioned the duct of the gland and demonstrated its opening at the vulva. Huguier² (1850) published a most extensive and thorough article on diseases of this gland and referred to it as the vulvovaginal gland.

In 1892 Müller gave a résumé of the literature of the development of Bartholin's glands. He says that they are first recognized in the fetus when it has attained a length of 4.5 cm. Here they are seen as solid cords passing off from the *sinus urogenitalis*.

Pohlman, who has been giving this subject much study in Professor Mall's laboratory, tells me that the beginning of Bartholin's gland becomes apparent just as soon as the first differentiation of sex appears externally. Formerly it was not thought to be demonstrable before the fourth month, but Pohlman has found it in embryos 2 cm. in length. The glands at this time consist of simple pouchings out at the sides of the urogenital sinus. Bartholin's gland in the female corresponds to Cowper's gland in the male. It does not, however, develop so rapidly as the male gland. There is much difference of opinion as to whether Bartholin's gland is ecto- or endo-dermic. Pohlman thinks that it is probably ecto-dermic.

ANATOMY.

These glands are two in number and are situated in the para-vaginal tissue. They are 2 to 3 mm. in diameter and 3 to 5 mm. long. The gland duct passes down on the inner or vaginal aspect and opens into the vagina just below the outer side of the hymen. Huguier, who

Fig. 1.—Relation of Bartholin's glands to the vagina. (Redrawn after Huguier.) In this case both glands show inflammatory enlargement and from the duct on each side a drop of pus is escaping. The glands are seen in the lateral vaginal walls, a short distance above the fourchette.

Fig. 2.—Deep relations of Bartholin's glands to the vagina. (Redrawn after Huguier.) In this case the duct to the right of the vagina has become occluded and is cystic, forcing the gland, which is considerably enlarged, further out than usual. On the left of the vagina the duct is also markedly cystic. Under ordinary circumstances the glands lie in the adipose tissue just external to the vagina and are partially covered by the band of muscle shown in the drawing.

27. Deutsche med. Woch., 1904, xxx, p. 1458.

* Read in the Section on Pathology and Physiology of the American Medical Association, at the Fifty-fifth Annual Session, June, 1904.

1. Oeuvres anat., Paris, 1761, vol. II, p. 319.

2. Mémoire sur les maladies des appareils sécréteurs des organes génitaux externes de la femme. Mémoires de l'Académie de Médecine, Paris, 1850, vol. xv, p. 531.