Original Articles.

THE CLINICAL VALUE OF THE WASSERMANN REACTION.

BY ABNER POST, M.D., BOSTON.

WHEN a practitioner sends a specimen of blood to a laboratory with a request for a Wassermann test as to the presence of syphilis and receives an answer "positive" or "negative," what ought that answer to mean to him?

This short paper is intended to answer such a question. In other words, it is an attempt to express the diagnostic value of the Wassermann reaction without any reference whatever to its theory or to the technical procedure in language which may be understood both by physician and layman.

In the first place—the Wassermann reaction is not directly a test for syphilis at all. It simply shows that syphilis has or has not produced a certain change in the blood. Syphilis does not produce that change at the moment of inoculation nor is it usually present until after the chancre or primary sore has existed for three or four weeks. Occasionally, it shows very early but oftener it is late, that is, three, four or five weeks after the appearance of the primary sore.

When we are told that the reaction is positive in primary syphilis in such and such a proportion of cases, it leaves the interest centered about the cases in which it has failed. The story is not complete unless we know the age of the lesion at which the blood was taken for examination.

The spirochetæ may be discovered some time before the change is recognizable in the blood by the Wassermann test. There may come, also, a time later in the disease when the blood ceases to respond to the Wassermann test, although the disease itself is still present. The disease is then latent or attenuated or powerless, so far as the production of the Wassermann is concerned.

The Wassermann reaction is most delicate and complicated. It can be performed only by skilled scientists specially trained for the purpose. It depends upon a certain substance in the serum of the blood which is known only by its influence on other substances. The amount of blood required is about a drachm and is usually drawn from a vein by the insertion of a hollow needle. The amount required is too large to be readily taken from the ear or finger tip. The reagents required in the original process were:

The patient's blood serum, heated for half an hour at 56° C. just before the experiment
Watery extract of congenital syphilitic liver
Guinea pig's blood serum, freshly drawn
Rabbit's blood
Washed sheep's blood

These are manipulated in five test tubes. The presence or absence of syphilis in the patient's serum is shown by its action on the sheep's corpuscles in the presence of the other agents. Two other test tubes are used for control experiments.

The reaction differs in intensity and this intensity is indicated by the number of plus signs used, though the same symbolism is not used universally. One series of symbols used in this vicinity is as follows:—

+	+	+	=	markedly positive
+	+		=	positive
+			=	slightly positive
-			=	negative.

When the Wassermann is positive, it means that the patient has syphilis. That is practically reliable in this country. There are, however, certain diseases other than syphilis which may give a positive reaction. These are yaws, leprosy, trypanosomiasis. The blood in certain temporary conditions has also given positive reaction, anesthesia, narcosis. A positive reaction has also been found in a small proportion of other diseases—malaria, scarlet fever, pneumonia, typhoid,—but not constantly. These may almost be said to be isolated observations of single observers, not confirmed by others.

Blood in certain other conditions has occasionally been reported positive. The amount of latent syphilis in the community makes it extremely probable that unexpected positive reports should occur not infrequently. For instance, a woman presented symptoms of disease of the internal organs. She bore the scar of an extensive operation for removal of the breast. There was every reason to believe that she was suffering from malignant disease of the mediastinum, secondary to the disease of the breast. For some reason, a Wassermann was done with positive result. She was given repeated doses of 606 without any change whatever in her symptoms.

A further inquiry into her history gave room to believe that she had had both syphilis and malignant disease. The positive report was evidently misleading. Because she had syphilis and a positive Wassermann, her attending physician drew the inference that everything was syphilitic.

The exceptions to the truth of positive reports are very few. Unfortunately they occur, when they do occur, in cases which are troublesome, as for instance, in the diagnosis between syphilis and leprosy.

From what has been said, it will be evident that a negative report does not necessarily mean no syphilis. It means either no syphilis or that syphilis, if present, has not produced that change in the blood which creates a positive reaction. A negative report is, then, simply negative evidence. It simply shows that there is no evi-

dence of syphilis in the particular specimen of blood submitted for examination, and as we have seen that syphilis may be present in the body and not show in the blood, it is evident that a negative Wassermann is not conclusive evidence of the absence of syphilis. In fact, it sometimes occurs that we get a negative Wassermann while characteristic syphilitic lesions are present on the body. Some of the serious cases of nervous syphilis occur after a negative Wassermann. This does not mean that a negative report is valueless.

The Wassermann reaction is of use for two different purposes. First, for diagnosis, and second, as a measure of the effect of treatment. So far, it has been considered in this article only as a help to diagnosis. As a measure of treatment, it is taken before treatment is begun, at intervals during treatment, and after treatment has been omitted for a length of time. When found positive before treatment and negative after treatment, it shows that treatment has had an effect on the disease. Some times there is a gradual reduction in the strength of the reaction in successive tests. It may be reported +++ before treatment, ++ after treatment has been begun, + a little later and negative a little later still. When the reaction is thus extinguished, there is always room for hope that the disease is absolutely cured. Unfortunately, the reaction is known to return to positive in some cases. This has led physicians to take repeated tests at long intervals after it becomes negative. How long such reactions should be followed before one can pronounce his patient positively cured is unknown. It has become evident that the disease does occasionally show its old characteristic of returning after an apparent cure.

The reaction may be rendered negative by treatment when the disease is not cured, exactly as other symptoms may be made to disappear. This change in serum reaction may sometimes be brought about by mercury before the symptoms for which it was given have disappeared. The same is true of salvarsan administration though the external symptoms are more apt to disappear before the serum reaction changes. A negative reaction is also said to be brought about by acute alcoholism.

A negative report is also of value as confirmatory evidence. This is best illustrated by an example. There was a difference of opinion between two doctors as to the character of certain lesions on a patient. One said that they were syphilis and the second denied it. They agreed to submit the case to a third doctor and to abide by his opinion. The third doctor found no evidence of syphilis. A few years ago, he would have said so—at present, he must have a serum examination before he can deliver a positive and satisfactory opinion. In this particular case, a negative Wassermann was necessary to complete the case. The report would have been incomplete without it.

A third report on blood submitted for the Wassermann test is 'occasionally returned—"anticomplementary." That means that complement, the unisolated and little understood substance, known only by its influence on other substances, and necessary to the Wassermann reaction, is not present in the specimen submitted. This substance, present in all fresh serum, disappears from blood too long preserved. That change in the blood renders it necessary that the specimen sent for examination should be freshly taken.

No sooner had Wassermann made public his discovery than scientists set to work to improve it, either by simplifying the procedure or by the introduction of more delicate reagents which should detect a larger number of cases than the original. At present, certain pathologists claim that their reagents recognize a larger number of cases of syphilis than those of other pathologists. This difference in procedure gives different results and different reports when specimens from the same individual are sent to different pathologists. In these cases, the positive report is returned usually as mildly positive, or is indicated by a single plus sign, showing that the two pathologists do not differ widely and that the case is on the border line, but they are very annoying to the clinician.

After the publication of the Wassermann reaction, a few enthusiastic pathologists felt that the diagnosis of syphilis could always be made in the laboratory and statements to that effect were widely circulated, but the reaction is a symptom rather than an unfailing test. When it proves positive, it is a very great help but a negative report by no means settles the matter. It means no more than the absence of any other important symptom. The case must be further studied clinically.

As a measure of the effect of treatment, it gives us something absolutely new and of very great value, but its greatest usefulness can never be reached until pathologists themselves reach unanimity in their methods.

But its value is not confined to individual cases. Its assistance in the study of the disease has been very great. It has helped to show how very large the proportion of syphilis is in the country; the dissemination of the disease; that certain diseases called para-syphilitic are still syphilitic; it has been of wonderful use in the study of hereditary disease.

If one will regard the Wassermann reaction as a symptom to be considered like other symptoms, and in relation to other symptoms, it will prove a wonderful help. If one insists in calling it a test, he may be confused rather than aided.

Our knowledge of the Wassermann reaction is not yet perfect. In fact, no definite conclusion has been reached as to what the Wassermann reaction really is. Our present knowledge is only a step towards a wider knowledge of yet hidden truth.