mid-leg, through an incision along the inner border of the gastrocnemius, retracting it and the soleus outward; the nerve lies beneath the soleus on the interosseus membrane; in the lower fourth, through an incision one finger's breadth behind the outside of the internal malleolous progressing at an angle of 20 degrees outward. The nerve lies deep to the tendons about one-third way between the internal malleolous and the tendo-achilles.

Several muscles have been omitted because of the impossibility of demonstrating their individual function or loss of function when paralyzed.

CONCERNING URETERAL STONES: THE REMOVAL BY NON- CUTTING OPERATION*

By E. P. MERRITT, M.D.,
Instructor in Genito-Urinary Surgery,
Atlanta Medical College (Emory University); Urologist, Georgia Baptist and Grady (City) Hospitals and the Anti-Tuberculosis Association,
Atlanta, Ga.

The question of ureteral calculi in the past few years has not been neglected in current literature, and great advancement has been made toward the absolute and the differential diagnosis. It is very easy in taking a history of some patients suffering with ureteral calculi to be misled from the point of pathology being sought; especially is this true of the right side, where the appendix is usually the law-breaker of the easy life. The gall-bladder in many instances furnishes symptoms clear enough to lead astray the most careful clinician.

ETIOLOGY

1. Hunner tabulates the etiology very clearly into three periods. The first period dates from early medical history to the later part of the nineteenth century, namely, the "Chemical" epoch. Most

writers confined their theory to that etiology.

The next, or second period, includes the last two decades of the last century and first part of the present, and may be called the "Bacteriological."

The third may be confined to the "Mechanical" epoch, which embraces injuries to the ureter from diseases and other abnormalities.

Of course, any argument on the etiology would almost necessarily fall under

Misleading Shadows Shown to be Phlebotisms by Employment of Iron Oxid Catheter.

one of the above theories, although some of the arguments are so varied and fascinating that the writer will briefly mention a few.

Van Helmont attributed stone-formation to the coagulation of uric acid which comes from the existence of potential alcohol, and the breaking up of urinary ferments.

2. Chetwood divides calculi into (A) Primary, which forms in the kidney where there is no morbid process; and (B) Secondary, where the formation is due to some pathologic condition in the kidney, making a nucleus for the stone-formation.

*Prepared for Section on Surgery, Southern Medical Association, Twelfth Annual Meeting, Asheville, N. C., Nov. 11-14, 1918, postponed one year on account of influenza epidemic.

3. Cabot does not believe that climate or varieties of diet play an important part in stone-formation: As for heredity,

Ureteral Stone, Afterwards Removed by Operating Cystoscope.

he says, "It is difficult of estimation." His theory of stone-formation is very plausible. He says supersaturated urine is secreted by the kidneys under certain conditions which are at present unknown. In this supersaturated urine, crystalization occurs. These crystals, under certain conditions of retention, whether renal or vesical, are held in the urinary tract a sufficient time to enable them to form a true concretion of calculus.

Infection is only a factor in so far as it gives rise to conditions favoring retention, thereby affecting the chemical composition.

4. Hunner is inclined to believe that infection plus stricture of the ureter is responsible for a large percentage of ureteral stones, the infection being the primary cause, possibly liberating stone nuclei.

It is quite plausible to think of a stone's forming after an infection, which gives up or throws off desquamated epithelia, clumps of bacteria and the like. Especially is this true with certain individuals

who are susceptible to stone formation due to hereditary inclination. Most of these patients were closely questioned. Heredity seems to be the answer.

Most authorities are of one accord that the ureteral stone-formation takes place in the kidney pelvis, which, to my mind, in a large majority of cases, is correct.

Personally, I think a ureteral stone will cause a stricture in more instances than a stricture will cause a stone. Nearly all stones are found at one of the normally constricted points of the ureter.

SYMPTOMS AND DIAGNOSIS

To recognize ureteral stone by symptoms other than the typic attacks of colic, requires considerable skill, as the symptoms vary so much. This part of the discussion will only be superficially covered. The ordinary individual suffering from this abnormality comes complaining of "pain" usually in the back, possibly either side, reflected to lower limbs and genitals. This is worse at some times than others, and he may be passing a small "gravel" or a little blood. On one or two occasions

Ureteral Stone Exhibiting Typical Shape

a "hypo" was necessary for relief. These attacks are generally intermittent, and thought to be colic or appendicitis, sometimes forgotten until a repetition occurs.

The diagnosis, after centralizing our minds on ureteral stone, would be practi-

cally easy if all the stones would show under the exposure of the x-ray. Thanks to the rays, a very large percentage does show; but now and then a pure phosphatic or uratic stone will not show. Then, if

be taken into consideration also, but will not be gone into in detail here.

NON-SUGRICAL REMOVAL

There has been much written of late years concerning the removal of ureteral stones by non-operative methods, and this is one of the sensible procedures to my mind. There are several different methods used for the removal of ureteral stones.

6. Walther reports three cases of ureteral calculi removed by injections of papaverin below the stone, causing dilation of muscle fiber.

7. Crowell and Thompson, in selected cases, use a 2% cocaine solution, injected below the stone, afterward using 10 c. c. sterile olive oil above.

Walther does not go quite far enough, only using the papaverin. Crowell, in my judgment, goes a little too far by the injection of too much oil into the kidney pelvis. Cocaine, in my estimation, is too toxic for general use.

The method which I have used successfully for some time is as follows:

Shadow Demonstrated as Ureteral Stone by Employment of Iron Oxid Catheter.

the clinical symptoms point to stone, shall we say "negative?" Not until the cystoscopic methods have proven so, namely, obstruction to catheters and negative findings per wax-tipped catheters.

5. Braasch thinks that the failure to show kidney or ureteral stone per x-ray is partially due to the roentgenographic technique. He is right, as the technique is the main problem in the x-ray work.

Shadows resembling ureteral calculi in the ureteral region are not always to be taken seriously until further investigation. For instance, pathologic mesenteric glands and phleboliths give shadows that are misleading. The differential diagnosis is made easy by inserting an iron oxid catheter into the suspected ureter, having a roentgenogram made and noting the obstruction met and the distance from catheter to shadow.

The history and urinary findings are to

Ureteral Stone, Later Removed by Operating Cystoscope.

Dilate the ureter below the stone with ureteral dilators. After the introduction of papaverin or novocain, 2 to 4 c. c. of 2% solution, the catheter is withdrawn so that the ureteral walls may be infiltrated thoroughly. If a small ureteral orifice

obstructs, then use ureteral scissors to cut the meatus to desired size. After this, pass a stiff catheter to or beyond the stone, if possible. Inject then from 2 to 4 c. c. sterile olive oil. Of late it has been my method to inject a few drops of 10% argyrol or silvol afterward for the anti-septic value it possesses, possibly reducing soreness from manipulation.

The patient, if not in pain, may sit up. If in pain, large doses of morphine, hot applications and an abundant intake of fluids are indicated.

Misleading Shadows Previously Diagnosed as Ureteral Stone. At Autopsy Found to be Tubercular Mesenteric Glands.

This procedure is employed at intervals of three days if the condition of the patient will permit. The average number of times for the removal has been three, with me.

8. As mentioned by the writer, the manipulation through the urethra with the cystoscope aids materially by setting up reflex peristaltic waves.

If in watching for a "big stone" we do not see it pass, we should not be discouraged, as it will oftentimes break up (especially is this true of the phosphatic type) and come out in small particles; or there may be more blood clot sometimes than there is stone. It is a good idea to be convinced of the absolute removal by another ureteral examination.

This method is convincingly reasonable and safe, even if we do not succeed every

Case Showing All Symptoms of Ureteral Stone with Misleading Shadow. Note Inward Displacement of Ureter.

time. There is no harm done, but in the majority of cases, especially, selected cases,

Probable Pathologic Mesenteric Gland. Shown to be Outside of the Ureteral Channel.

this method saves a cutting operation. Twenty-three cases have come under my

care. Of that number, twenty-one passed the stone after the above-mentioned manipulation.

BIBLIOGRAPHY

- (1) Hunner, Guy L.: "Etiology of Ureteral Calculus," Surgery, Gynecology and Obstetrics, Sept., 1918.
- (2) Chetwood: "The Practice of Urology," p. 562.
- (3) Cabot, Hugh: "Modern Urology," p. 133.
- (4) Hunner, Guy L.: "Etiology of Ureteral Calculus," Surgery, Gynecology and Obstetrics, Sept., 1918.
- (5) Braasch, W. F.: "Clinical Data of Nephrolithiasis," Surgery, Gynecology and Obstetrics, Jan., 1917.
- (6) Wallner, H. W. E.: "Papaverin in the Treatment of Ureteral Calculus," The Urological and Cutaneous Review, Sept., 1916.
- (7) Crowell, A. J., and Thompson, S. R.: "Diagnosis of Ureteral Calculi and the Technique of Removal with Operation," Journal American Medical Association, Aug. 10, 1918.
- (8) Merritt, E. P.: "A Consideration of Ureteral Stones, Their Removal by Aid of the Operating Cystoscope: Report of Cases," Surgery, Gynecology and Obstetrics, Nov., 1918.

A RARE SURGICAL CASE

By V. B. PHILPOT, M.D.,
Houston, Miss.

The writer does not feel certain that the diagnosis made in this case was correct, but is of the opinion that the trouble with this little patient was a lymphangioma or a cyst of the lymphatics.

The history of this case is as follows:

The patient is now four years old. She was born with two small tumefactions—one below the axilla and one just above the clavicle to the left of the neck. The tumors grew but little until six months previously, at which time a very rapid growth took place and constitutional symptoms became manifested, the patient becoming weak and emaciated and at times having abnormally high temperature which was supposed to be due to some inflammation in the tumor.

I saw her four months ago and advised surgical interference and at the same time informed the family that I could promise nothing as to the outcome. Her people hesitated for a couple of weeks, and as the condition continued to grow, decided to take a chance in having it removed.

We made an incision in the tumor below the axilla, hoping to remove it, sac and all; but we found after draining the fluid out that the sac extended deep between the ribs and formed pockets all through the upper left chest. One of the deepest pockets extending to the scapula behind and another, between the clavicle and scapula, forming the large cyst to the left of the neck above. The sacs were of course adhered to the bones everywhere and the idea of removing them was eliminated. All the sacs communicated, however, and drainage of all was accomplished. About a quart of blue fluid was drained off. We next curetted the cavities in every direction as far as possible and swabbed with iodine, hoping to destroy the secreting cells.

Our efforts were only partially accomplished as in a few days a large tumor began to appear

to the left of the neck. Another anesthetic was given and an incision was made above the clavi-

Before operation

cle and with dressing forceps we swabbed the cavities from the incision here to the previous incision below the axilla. Another lot of fluid

After operation

was drained. After this the tumefaction above and below were apparently cured, but in five or six days a new growth formed above the left nipple. So the third anesthetic was given and an incision made here and the pockets which we found to be three or four in number were curetted and swabbed as in the previous operations.