The Origin of Cancer.

By

Dr. Arthur Turnbull (Glasgow).

The chief stumbling block towards the elucidation of the problem of cancer has been the nature of the internal change at work within the tissues, or, in other words, the nature of the variability by which what we call a „normal“ becomes an „abnormal“ tissue. Let us illustrate this by the case of the tooth rubbing on the tongue and occasioning a cancer of the tongue. Here we have within the tongue:

A moment's reflection shows that this intermediate process is the heart of the cancer problem.

What then are our views of this internal variability of tissue? They are two, views so utterly contradictory that they cannot both be right. They may be briefly summarised in this way:

1. Cancer has a cause.
2. Cancer has no cause.

This distinction may be expanded as follows:

- (1) In this case we look upon the body as a test-tube varying only in response to the external environment, and naturally, if we have this idea, we shall try to explain the development of cancer in terms of the external environment. We shall regard the cancer-change as the effect of the action of parasites. We shall note that nearly all the known diseases are produced by bacteria, we shall lay stress on evidences of tissue-socialism (symbiosis), we shall discover ultra-microscopic bacteria, we shall describe cases in which one person

passed on cancer to another, we shall prove the existence of cancer houses in which family after family became affected. For instance, one great authority, Sir William Macewen, points out that cancer affects chiefly the flexures of the gastro-intestinal tract, that is, the points where the supposed external parasite will impinge, e. g. at the lip, tongue, lower ends of gullet and stomach, at the commencement of and hepatic, splenic, sigmoid and rectal twists of the great bowel, at all of which points cancer is alleged to occur with especial frequency. On this idea, therefore, a cancer is something foreign and pathological to the body.

- (2) In this case we shall look upon the body as varying not in response to external causes, but from within. We shall therefore regard cancer as an internal eruption of the body from one phase (the normal) into another phase, which we call cancerous or malignant. We shall hold cancer to be a physiological and not a pathological process. We shall regard it as compensatory in every case. We shall discredit the idea of parasites and shall lay stress rather on its usually slow and insidious course. We shall find a complete series of linking growths from the benign galls, warts and polyps to the most rapidly fatal sarcoma. We shall believe that not merely are there men and women diverging into cancer to-day, but that there are many who have diverged in the direction of cancer only to recover themselves and many who are to-day curing themselves from within. We shall regard cancer as a tissue-divergence, which may sometimes cure itself as a tissue-retention.

Now could this difficulty be dissolved? The following is an attempt at solution. It started from two stray remarks, each from a famous expert, the one proceeding from Prof. Dr. D. v. Hansemann, of the Virchow-Krankenhaus, Berlin, the other from Dr. W. H. Mayo, of Rochester, Minnesota, U. S. A.

Prof. v. Hansemann said: You will remember one thing, cancer has already been experimentally produced in man, namely, by the X-rays.

Dr. Mayo said: There is only one thing known to cause cancer, and that is chronic irritation.

The deduction drawn from the first of these remarks was that there did not seem to be much sense in wasting time on experiments, when the X-rays had already carried out the experiment for us; clearly, the one thing now remaining was to understand the internal principle by which some men became cancerous on X-rays and others did not.

The deduction drawn from the second remark was the equation:

Chronic Irritation i. e. Mechanical-strain = Cancer.

Now, nearly all cancer workers have attacked this equation from the cancer side; it was decided to proceed from the other side of the equation and investigate the result of strain upon the normal tissues.

How could this be done?

Firstly, by securing a means of measuring the variability of tissue: secondly, by securing an easily measurable tissue exposed to an excessive strain.

- (1) It is needless to discuss the variability measures in common use. A simple illustration of a close or low as against an extended or high variability will be understood by a glance at the two following series:

10 . 11 . 12	13 . 14 . 15
8 . 9 . 11	14 . . 16 . 17
Mean-Value.	

- (2) Bones were the only tissues measurable easily and in large numbers. The neck of the thigh-bone (femur) appeared to be a part exposed to an excess of strain.

Accordingly, 1100 femora were measured,

- A. at the narrowest diameter of the neck,
- B. at the narrowest diameter of mid-shaft,
- C. at the narrowest antero-posterior diameter at the lower end.

The variability-figures were as follows:

- A. 10.7
- B. 8.7
- C. 8.7

It appears therefore that when a tissue is exposed to especial strain, and a series of units of that tissue be examined, the units of the series are found striking away from a mean-value and towards an extreme-value.

Returning now to the tooth rubbing on the tongue (or the pipe on the lip, the blow on the bone, etc. etc.) and substituting this new principle of tissue change into the sequence of events, we find:

1. In the tongue:

2. In the thigh-bone:

Therefore 3: In the tongue we get

This solves the problem of cancer in a sense opposite to the parasitic idea, and determines that cancer has no cause, being nothing more nor less than a tissue divergence.

These represent the measurements and corresponding variability figures for the narrowest neck and narrowest mid-shaft diameters of 1600 ancient Egyptian femora, lying in the New Museums, Cambridge¹). Access to these was secured through the courtesy of Professor Alexander Macalister.

Femora.

1st 100.				2nd 100.			
Neck		Mid-Shaft		Neck		Mid-Shaft	
		cm				cm	
$v = 10.2$	1	1.8		$v = 11.3$	1	1.7	
	7	1.9	1		3	1.8	1
	18	2.0	13		8	1.9	
	15	2.1	13		16	2.0	2
	18	2.2	18		10	2.1	19
	9	2.3	19		12	2.2	26
	12	2.4	15		11	2.3	18
	9	2.5	9		13	2.4	15
	8	2.6	10		13	2.5	10
	2	2.7	1		8	2.6	7
	1	2.8			4	2.7	2
		2.9	1		1	2.9	
$v = 8.7$				$v = 7.5$			

1) The 10th hundred contains some english bones.

Femora.

3rd 100.			
Neck		Mid-Shaft	
cm			
$v = 9.3$	2	1.8	1
	15	1.9	1
	28	2.0	13
	15	2.1	22
	17	2.2	21
	5	2.3	22
	10	2.4	11
	6	2.5	7
	1	2.6	1
	1	2.7	1
		2.8	
$v = 7.3$			

6th 100.			
Neck		Mid-Shaft	
cm			
$v = 10.2$	2	1.8	1
	9	1.9	8
	12	2.0	17
	15	2.1	13
	11	2.2	16
	19	2.3	15
	15	2.4	9
	8	2.5	15
	4	2.6	4
	3	2.7	1
	2	2.8	1
$v = 9.7$			

4th 100.			
Neck		Mid-Shaft	
cm			
$v = 9.6$	2	1.8	1
	5	1.9	1
	17	2.0	8
	22	2.1	22
	15	2.2	10
	7	2.3	23
	11	2.4	16
	14	2.5	12
	7	2.6	5
		2.7	2
		$v = 8.2$	

		7th 100.	
Neck			Mid-Shaft
		cm	
$v = 13.3$		1.4	
	1	1.7	1
	9	1.8	5
	9	1.9	10
	16	2.0	18
	18	2.1	23
	15	2.2	12
	4	2.3	5
	8	2.4	9
	6	2.5	3
	5	2.6	10
	6	2.7	3
	1	2.8	1
	1	2.9	
	1	3.3	
			$v = 11.8$

5th 100.				
Neck		Mid-Shaft		
cm				
$v = 11.0$	5	1.8		
	10	1.9		
	15	2.0	10	
	13	2.1	26	
	20	2.2	18	
	17	2.3	17	
	5	2.4	15	$v = 7.7$
	3	2.5	9	
	5	2.6	3	
	6	2.7	2	
	1	2.9		

8th 100.			
Neck		Mid-Shaft	
cm			
$v = 10.8$	1	1.7	
	4	1.8	1
	9	1.9	3
	14	2.0	12
	9	2.1	14
	14	2.2	15
	13	2.3	13
	16	2.4	20
	13	2.5	16
	4	2.6	3
	2	2.7	1
	1	2.8	1
	1	2.9	1
		$v = 9.5$	

Femora.

9th 100.		
Neck	Mid-Shaft	
cm		
$v = 10.2$	1 1.7	
	3 1.8	1
	5 1.9	2
	8 2.0	5
	19 2.1	13
	12 2.2	18
	16 2.3	24
	12 2.4	17
	12 2.5	12
	8 2.6	6
	4 2.7	2
		$v = 7.8$

12th 100.		
Neck	Mid-Shaft	
cm		
$v = 9.3$	1 1.7	
	2 1.8	
	5 1.9	
	14 2.0	17
	18 2.1	15
	18 2.2	17
	17 2.3	22
	10 2.4	19
	9 2.5	12
	4 2.6	1
	2 2.7	3
		$v = 7.5$

10th 100.		
Neck	Mid-Shaft	
cm		
$v = 12.9$	1 1.6	
	2 1.8	
	5 1.9	4
	8 2.0	8
	13 2.1	17
	13 2.2	12
	9 2.3	17
	16 2.4	18
	10 2.5	9
	11 2.6	6
	4 2.7	5
	1 2.8	4
	2 2.9	
	3 3.0	
	1 3.1	
	1 3.2	
		$v = 9.7$

13th 100.		
Neck	Mid-Shaft	
cm		
$v = 10.2$	1 1.7	
	6 1.8	
	6 1.9	2
	15 2.0	7
	13 2.1	22
	25 2.2	16
	12 2.3	18
	6 2.4	22
	10 2.5	6
	4 2.6	6
	1 2.7	1
	1 2.8	
		$v = 7.7$

11th 100 (85).		
Neck	Mid-Shaft	
cm		
$v = 9.0$	1 1.9	1
	1 2.0	1
	4 2.1	5
	9 2.2	8
	7 2.3	15
	17 2.4	20
	17 2.5	19
	13 2.6	9
	5 2.7	6
	5 2.8	1
	6 2.9	
		$v = 7.2$

14th 100.		
Neck	Mid-Shaft	
cm		
$v = 9.6$	3 1.8	
	10 1.9	2
	19 2.0	10
	18 2.1	14
	12 2.2	27
	17 2.3	14
	9 2.4	26
	9 2.5	4
	1 2.6	2
	1 2.7	1
	1 2.8	
		$v = 7.1$

Femora.

15th 100.			
Neck	cm	Mid-Shaft	
$v = 10.4$	1 1.5		
	1 1.7		
	4 1.8	1	
	5 1.9	2	
	10 2.0	9	
	18 2.1	16	
	14 2.2	18	
	13 2.3	15	
	16 2.4	21	
	11 2.5	12	
	6 2.6	5	
	2.7	1	
	1 2.8		

 $v = 8.1$

16th 100 (80).			
Neck	cm	Mid-Shaft	
$v = 9.8$	4 1.9		
	9 2.0	2	
	12 2.1	9	
	15 2.2	25	
	8 2.3	12	
	11 2.4	12	
	10 2.5	9	
	6 2.6	8	
	4 2.7	1	
	1 2.8	2	

 $v = 7.8$