PART III.

MEDICAL MISCELLANY.

Reports, Transactions, and Scientific Intelligence.

Resection of the Stomach, a portion of the Duodenum, and the greater part of the Pancreas. By M. RICARD. Translated by GEORGE M. FOY, M.D., F.R.C.S.; Surgeon, Whitworth Hospital, Drumecondra.

At a meeting of the Academy of Medicine of Paris (*Gazette des Hôpitaux*), M. Ricard asked permission to relate the history of an illness, and to present the patient herself. She is a thrifty woman of fifty-two years. She entered the Hôpital St. Louis on the 20th April, 1899, for gastric pains which commenced two years before. After six months her sufferings increased, and were intensified after food. Her appetite completely went; she had vomiting daily. She wasted to a perfect skeleton, not weighing more than 102 lbs. Her skin was pale and bloodless. A tumour irregular in shape and hard to the feel was felt in the epigastric region. Carcinoma of the stomach was diagnosticated. A certain degree of mobility, the form of the lump, the change of position of the lump when the stomach is empty and it is distended, caused the surgeon to think that the tumour was situated on the greater curvature of the viscus. On the 26th of April the patient was anæsthetised, and a sub-umbilical incision made, by which it was discovered that the tumour not only occupied the greater curvature of the stomach, but also extended over the two surfaces, and principally over the posterior wall. The mass was with difficulty moved, and at first seemed incapable of removal; nevertheless, a careful examination showed that the two ends of the mass were accessible—easily at the duodenal end; with difficulty at the cardiac end. This revealed the difficulty that beset the operation.

After making an opening in the gastro-hepatic epiploon, sufficient for the insertion of a pair of clamps to cut off the blood supply of the section, the duodenum was cut off at the beginning of its vertical portion. Two long forceps were placed at the left of the neoplasm, and the stomach was cut at the cardiac end of the œsophagus. An oblique fold of tissue, from 3 centimetres to 4 centimetres long was

left at the great extremity. The stomach was now detached from its peritoneal bands, and the divided peritoneal fold sewed together with catgut. Withal, though the extremities of the stomach and its greater and less curvatures were free, it was found that the diseased mass had a deep attachment that prevented the removal of the viscus and of the neoplasm. Traction from left to right and from right to left was now made on the mass, and it was found that the growth was attached to the pancreas. The body of the pancreas was now freed, most of the vessels of the spleen liberated, and the to-and-fro movements recommenced. The liberated portion of the pancreas was now dissected from the vena cava and removed. A mass of infiltrated ganglia was detached from the aorta. Some difficulty was experienced in arresting the bleeding on section of the pancreas, hæmorrhagic points having to be freely touched with the thermo-cautery, and finally compression by a pad of gauze had to be resorted to.

The first step of the operation—that of excision—was now finished; the second and more difficult step, that of reparation, was now to be commenced. The open extremity of the duodenum was now closed by suturing the different tissues together, like-to-like, and finally the peritoneal tissue closed over by Lembert sutures. By careful placing of the preserved tissue around the cardiac orifice of the stomach, a small cavity, about the size of a hen-egg, was formed. At this period of the operation the alimentary tube was divided into two disconnected parts, so far removed and each so fixed that they could not be drawn together. Catching the jejunum, it was drawn up to the cardiac orifice, and the ampulla formed of portion of the stomach was fixed in it as in an ordinary case of gastro-enterectomy. The ampulla looked like a reversed V, the apex being at the œsophagus, and the other extremity being fixed in the jejunum. The operation was now practically finished.

It was found necessary to retain the compress on the cut pancreatic stump, and from it a lamp-wick-like fold of gauze was brought out of the cutaneous wound for drainage. The superficial incision was sewn together by ordinary sutures and a few interrupted silver wire sutures: One hour and a quarter was occupied in the operation. During the last fifteen minutes the patient got a hypodermic of caffèïn and an intravenous injection of five hundred grammes of serum.

Considering the gravity of the operation the sequelæ were good. The highest temperature for the first three days was 100° F. during the sojourn of the patient in hospital; afterwards the temperature

did not exceed 99° F. On the first day after operation the patient got seventeen hundred and fifty grammes of serum, five hundred being given by intravenous injection. On the evening of the second day she vomited blood. Rectal feeding with peptonised food was commenced on the second day. On the third day the tampon of gauze was removed, and a pancreatic fistula followed, which remained for twenty-five days. On the fourth day she took some milk and some weak grog. A little pottage and some milk were taken on the fifth day. The patient remained in hospital for one month, during which time she put on flesh and gained strength. A histological examination confirmed the diagnosis of cancer of the cylindrical epithelioma variety, which invaded the pancreas and the neighbouring ganglia. To-day the patient, nine months after operation, has regained her strength and increased fifty-two pounds in weight. She has a fresh, ruddy colour, and, in a word, is in good health.

This case is interesting from the extent and complexity of the operation, from the good convalescence of the patient, from the freedom of any symptom of the return of the disease, and from the good health she enjoys.

M. Ricard considers that success will follow all similar operations. He affirms that extirpation of a portion of the stomach is practically safe, and refers to his eight cases of pylorectomies with seven recoveries published by M. Guillot in the *Gazette des Hôpitaux* in confirmation. He has the conviction that the excision of cancers of the stomach, in good time—with average skill, with pure asepsis—are less fatal than extirpation of cancers of the uterus or rectum, which all the world practises freely.

It is for the physician to diagnosticate such cases for the surgeon; it is his to recognise the contra-indications for operation which experience has taught him. He will learn to be bold even to rashness. This case is reported from others equally successful that it may encourage physicians and surgeons who hesitate.

A MAN WITH A QUIVERFUL.

A FRENCH Canadian living in a Rhode Island town was recently presented by his third wife with his forty-first child. His first wife gave birth to several pairs of twins, and his second presented him with three sets of triplets. Thirty-six of the children are living and many of them have families of their own. Eight of the grandchildren also are parents.

CLINICAL RECORDS.

The Treatment of Dysentery by Intestinal Asepsis. By JAMES R. WALLACE, M.D., F.R.C.S.I.; Surgeon to the Home Hospital, Calcutta.

THE following cases illustrate the successful treatment of ordinary tropical dysentery, or what should really be styled muco-enteritis, by a simple method of converting a septic alimentary canal into an aseptic one.

The routine was the same in all the cases—namely, the administration of a dose of an emulsion of castor-oil, terebene and opium, morning and evening, 20 drops of liquor hydrargyri perchloridi B.P. every 6 hours, and a copious enema of boro-glyceride in a warm water solution every 12 hours. The diet consisted of light, solid food, not milk slops and soups, with fruit and vegetables given freely.

CASE I. (No. 7 in Hospital Case Book).—Mrs.——, a young English woman, aged twenty-six, married, one child two and a half years old, menstruation regular, appetite fairly good, very emaciated and anæmic in appearance. Has been ailing with small bloody mucus discharges from the bowels for three months, and has had from six to a dozen motions every day for the past three weeks. Before that they were not so frequent, though three or four stools of a similar character were voided daily throughout this period. There had been, and was still, considerable pain and tenderness about the bowels and a great deal of tenesmus during defæcation. The patient had lost weight very remarkably during her illness. She was admitted into the Home Hospital on the 6th December, when it was found that the stools consisted of about a couple of ounces of dark green, blood-tinted mucus of a jelly-like consistency. They were of an extremely offensive putrid odour. There was painful tenesmus with each evacuation. There was a distinct tenderness over both iliac fossæ. The liver was tender, but not enlarged; the spleen was normal; the kidneys and urine were healthy; the lungs were healthy; the heart's action was weak; there was slight fever, temperature 100·8°; appetite poor, but there was a slight inclination for solid food and a dislike for slops. She was placed on the above treatment and on solid food, with the effect that within 48 hours the stools had lost their foetor, and their frequency had been diminished by half, while their character was also altered, there

being a deal of grumous faecal matter with much less blood and mucus. Within three days more the stools became normal, the appetite was restored, all pain had subsided, the temperature was 98·6° and the patient was practically cured. She was detained in hospital till the 19th December to allow her to pick up, but all medicine was stopped on the 14th.

CASE II. (No. 8 in Hospital Case Book).—Mr.——, a Scotch engineer, aged forty-seven, married, who gave evidence of a once robust and well-built frame, but who was now emaciated and cachectic looking, was admitted into the Home Hospital on the 6th December for typical tropical dysentery, from which he had suffered at intervals of a month or so for the past three years. The present attack began about a month ago. He was having 10 or 12 motions in the 24 hours accompanied by great pain and straining while at stool. The motions were very offensive and consisted of glutinous mucus and blood, with occasionally a little liquid faecal matter. He had slight fever, and his temperature in the afternoon was 102°. This rise of temperature was often preceded by either a sense of chilliness or sometimes a distinct ague fit. He also was subject to a good deal of nausea and often rejected his food. There was no enlargement of his liver or spleen, the lungs and heart were normal, his abdomen was distended, and the whole intestinal area was tender to the touch. He was placed on the antiseptic treatment above described. In addition, 5 grains of quinine were given at 9 o'clock in the morning and 3 o'clock in the afternoon, to combat the evident malarial infection. Within four days the motions assumed a healthy character, consisting of grumous faecal matter, with only a slight trace of mucus without any blood-tinging or foetor. There was still occasional nausea after food and some tenderness over the bowel on palpation. There was also a rather marked voraciousness of appetite. Suspecting the lodgment of worms in the gut, he was given 5 grains of santonin with 2 grains of calomel, followed by a double dose of castor-oil emulsion. Within 12 hours he voided a regular nest of lumbrici, made up of no less than 21 small round worms, matted together in the shape of a ball. Next day he passed 5 more worms. All medicine was stopped for 48 hours, and he was then given a tonic of dilute nitromuriatic acid and chiretta. He rapidly gained flesh, and was discharged on the 30th December quite well. I saw him two months later when he visited Calcutta, and learnt that he had had no further bowel trouble, and that he had gained nearly a stone in weight.

CASE III. (No. 19 in Hospital Case Book).—Mr.——, a German, aged thirty, four years in India, had an attack of dysentery a year ago, and was treated by me at his own home. He was cured in ten days. He suffered from a similar attack six months later, and was all right again after a fortnight's treatment. On both occasions the antiseptic method was adopted. He was admitted into the Home Hospital on the 27th January, 1899, with copious bloody motions mixed with mucus, and attended with painful tenesmus and tenderness over the bowels. Within 24 hours after the aseptic treatment the motions were completely altered, and he was discharged cured on the 29th January, 1899, three days after admission. He remained in good health for three months in Calcutta, and then proceeded on a voyage to Germany. While at Colombo, he got another attack of hæmorrhagic dysentery, which killed him in three days.

CASE IV. (No. 42 in Hospital Case Book).—Mrs.——, an English lady, aged twenty-seven, married, two children, had suffered from dysentery on several occasions in Darjeeling during the past four or five years, each attack lasting about six weeks and confining her to her bed. She had been ill for a week before her admission into the Home Hospital on the 10th April, 1899. She was very pale and thin, and was in great pain. Her bowels were moved almost every 20 minutes, the evacuations being scanty, bloody-mucus, and very offensive. There was also intense retching and vomiting. The whole intestinal area was acutely tender, more especially about the cæcum. There was some tenderness over the liver, but no enlargement. There was slight fever, temperature 101° . A single boro-glyceride enema gave marked relief, while the action of the bichloride and terebene worked an almost immediate change in rendering the motions non-offensive in odour. The fever subsided in 48 hours, and on the third day she had only three movements of the bowels, and these were only slightly tinged with bloody mucus, and consisted chiefly of grumous fecal matter, largely tinted with bile. On the fifth day the stools were normal, and on the 16th April, seven days after admission, she was allowed to go home, perfectly cured.

CANCER OF THE RECTUM.

HOCHENEGB says—Do not operate radically if the patient is very weak; if there are metastases; if the mass is firmly anchored in the pelvis; if glandular infection passes beyond the reach of the finger; and if acute obstruction arises.

ROYAL ACADEMY OF MEDICINE IN IRELAND.

President—EDWARD H. BENNETT, M.D., F.R.C.S.I.

General Secretary—JOHN B. STORY, M.B., F.R.C.S.I.

SECTION OF STATE MEDICINE.

President—SIR C. A. CAMERON, C.B., M.D.

Sectional Secretary—NINIAN FALKNER, M.D.

Friday, February 16, 1900.

The PRESIDENT in the Chair.

Presidential Address.

SIR CHARLES A. CAMERON, C.B., on taking the Presidential chair, thanked the Council of the Section for having nominated him as President, and the Academy for having elected him. He would give a short historical account of the origin of the Section. They all were aware that the Academy was formed in 1882 by the amalgamation of the Medical, Surgical, Pathological, and Obstetrical Societies. At that time there was in existence a Society of Metropolitan Officers of Health, of which he had the honour of being President. The Society met monthly, and many valuable papers were read before them, and they had an annual dinner. He suggested that, as the various Medical Societies were being amalgamated, it would be desirable that their Society should enter into the Union under the title of Section of Hygiene. There was considerable opposition to this proposal, on the grounds that it would unduly increase the number of sections, but ultimately the proposal was adopted in a modified form, and a Sub-section of State Medicine was constituted. The four Sections were each to have a President, but the Sub-section was to have only a Chairman. Some years subsequently the Sub-section was elevated to the dignity of a Section, and the Chairman was converted into President. He was the first Chairman, and now, after an interval of 18 years, he again filled the highest office, under the title of President. Sir Charles then referred to several topics, including the notification of infectious diseases. He regretted that the Notification of Infectious Diseases Act was not a compulsory one, and he also regretted that so many medical men were opposed to notification. Measles and whooping-

cough were not amongst the diseases enumerated in the schedule of the Act, but sanitary authorities had the power of adding them to the schedule with the assent of the Local Government Board. The Corporation of Dublin had included both diseases in the schedule, but whooping-cough had since been eliminated, as no good had resulted from notification of that disease. Measles was hardly notified in any town save Dublin, but he believed it was desirable that it should be notified. If the sanitary authorities in Ireland had the power of closing schools at the beginning of an epidemic (a power possessed by sanitary authorities in England), epidemics of measles might be checked earlier than they can now be. Notification of diarrhoea was required when these countries were threatened with cholera, but were only scheduled until all danger had passed away. Sir Charles referred to the inquiry now going on in London as regards the use of preservatives in food, and believed that some of them, such as boric acid, if used in moderation, would be useful. It would be better to consume food with a very moderate amount of preservative, which kept the food fresh, rather than food in a state of incipient decomposition, and containing deadly ptomaines.

Infantile Insanity.

DR. WALTER BERNARD read a paper on this subject.

DR. CONOLLY NORMAN said he had listened in vain for any practical suggestion from Dr. Bernard. He had spoken with too much generality. A great deal was heard about "insane predisposition," and it should interest us all, for we all have it. It was a very short and obscure family which did not contain many lunatics. Dr. Bernard had employed another vague phrase in constant use, and to which it was difficult to attach any definite meaning—"judicious training." We were told lately that a British youth leaving a university had about as much education as an eight year old German child. Which system was right?

DR. SYMES said feeble-minded children were very different from idiotic children, but he thought we did not distinguish between the two classes in this country. He thought many of those charged with such offences as causing fires, &c., suffered from some mental defect. In the homes for children to which he was physician he divided all the children into the following six classes:—(1) Ordinary normal children, in health; (2) children physically but not mentally defective; (3) defective and feeble-minded children incapable of attending school; (4) epileptic children; (5) blind and deaf children; (6) idiots and imbeciles.

DR. DAWSON agreed with Dr. Conolly Norman's remarks as to the excess to which the theory of heredity is carried now-a-days. No doubt, in certain cases, there would be no difficulty in making out a neurotic heredity, and in these cases prophylaxis should be employed if possible. He (Dr. Dawson) thought Dr. Symes had missed the point of Dr. Bernard's paper. Dr. Bernard had referred to extreme early infancy.

DR. DOYLE thought self-control was the point needing greatest emphasis. He had seen very marked improvement effected in a case which had been sent to an institution in Holland where the patients are taken charge of by farmers, and where no education is attempted, the patients being chiefly employed in manual labour.

SIR FRANCIS CRUISE thought Dr. Bernard's paper one of great importance in directing attention to the mental condition and proper mental training of children. He had seen children of two and three years of age whose mental condition pointed to insanity, and who ultimately proved to be insane. He was strongly convinced that a great many children are insane.

DR. LITTLE hoped Dr. Bernard's paper was only a preface to a work on the subject. No advance would be made if they were content to deal with generalities. Undoubtedly, a great deal could be done in the training of children who had a tendency to grow up imbeciles. About fifteen years ago a boy came under his notice who was growing up an imbecile. He was sent to an institution near London, and is now an officer in the army.

SIR CHARLES CAMERON referred to the work being done by the London School Board in the training of children with impaired intellects. The children are examined by the Medical Officers of the Board, and special classes have been formed for the purpose of teaching those with mental defects.

DR. BERNARD, replying, said the perceptive faculties do not begin to dawn until the dentition period, and the infant could be ruled and controlled in the same way as an animal. The moment the child is born the doctor should ascertain if it is physically normal. Then the cry, and the movements of the hands and legs, should be watched. Certain rules of guidance should be laid down for the mother and nurse, and so the infant automaton could be taught to observe order.

The Section then adjourned.

SECTION OF MEDICINE.

President—JOHN W. MOORE, M.D., President of the Royal
College of Physicians of Ireland.

Sectional Secretary—R. TRAVERS SMITH, M.D.

Friday, March 9, 1900.

The PRESIDENT in the Chair.

Exhibits.

DR. CONOLLY NORMAN—Cases illustrating the relations of insanity to tabes dorsalis.

DR. WALTER SMITH asked if these were looked upon as independent cases of mental derangement complicated by tabes, or as mental derangement a part and parcel of tabes, as he had for a long time regarded tabes dorsalis as a cerebro-spinal condition, and the early development of optic atrophy, ocular paralysis, &c., seemed to point to this idea. He inquired if Dr. Norman had ever seen what he might call cerebral crises—that is, sudden attacks of giddiness, and a tendency to feel as if the whole brain was slipping away.

DR. W. R. DAWSON said that the association existing between tabes and general paralysis showed that the two diseases must have something akin, and that there was no hard and fast line to be drawn between them. Strümpell was of this opinion, and believed general paralysis to be simply cerebral tabes, and this point was borne out by the first of these cases. This patient had at one time excessively exaggerated knee reflexes, which were now completely lost. In cases where tabes complicated general paralysis the tabetic symptoms appeared first with loss of knee reflexes, and as the tabetic symptoms disappeared the knee reflexes recurred. In this case the order of symptoms was reversed.

DR. NORMAN, in reply to the inquiry of Dr. Walter Smith as to association of insanity with tabes in these three cases, said that the association in the first case was probably only an accidental one. It might be that cases two and three would some day develop symptoms resembling those of general paralysis, which would bring them into line with those cases seen every day, in which tabes appeared first and then general paralysis. Of course, tabes dorsalis, like every other nervous disease, probably attacked, pre-eminently, those inclined to nervous failure. Referring to general paralysis associated with tabetic trouble, he pointed out that lesions have been traced in general paralysis in every part of

the nervous system from the cortex to the nerves. From time to time they were wont to see the occurrence of cerebral crises, which were remarkably like the other visceral crises, which tended to show that the cerebral trouble was similar to elsewhere. He got from one to three cases per annum of locomotor ataxy which had gone round the Dublin hospitals, and then terminated with him, and turned out to be general paralytic cases.

Replying to Dr. Dawson, the remissions of symptoms gave the first case a peculiar character, and it was to be noticed that there were remissions in the tabetic symptoms as well, although they were not so well marked. The patient was at one time so tabetic that he could not walk at all, but he was now able to walk about.

A Fatal Case of Congenital Bullous Eruption.

DR. R. H. KENNAN read a paper on this subject. [It will be found in Vol. CXIX., page 241.]

DR. LANGFORD SYMES said this case resembled one he saw some years ago, but the bullæ were not so prominent, and it was more a case of violent purpura than this case. It was in a two or three weeks' old child, which died within four or five hours of its admission to hospital. It had been described by German writers as purpura fulminans, and the blood-stained fluid described by Dr. Kennan seemed to point in this direction, but the case he had seen had not so many bullæ, and had far more purpuric marks.

DR. WALTER SMITH said that bullæ were produced on the skin under such diverse conditions that it was doubtful how far they were to accept a particular generic word such as pemphigus. He objected to placing a skin affection due to a distinct ætiological factor, say syphilis, under the same name as one not due to a distinct cause. He, therefore, objected to a diagnosis such as syphilitic pemphigus or psoriasis, because it was not possible that psoriasis could be both syphilitic and non-syphilitic. They could say a scaly syphilide. He agreed with Dr. Kennan in the non-syphilitic character of his case. They often saw pemphigus in connection with acute dermatitis, such as erysipelas, and even in ordinary scabies he had seen very large bullæ occur, as well as in cases of neuritis, such as herpes zoster and traumatic affections of nerves. In the class due to unknown causes he would rank cases he would like to apply the term pemphigus to. He would be inclined to define pemphigus as "an eruption of abruptly defined, fragile, thin-walled bullæ, upon a non-inflamed surface, symmetrical, and prone to relapse." They might be content to leave Dr. Kennan's case in the class of those due to an unknown cause,

though the big word tropho-neurosis might please some. Even in cases of ordinary pemphigus of a relapsing sort repeated bacteriological examination had revealed nothing definite.

DR. KENNAN briefly replied.

Notes on a Case Simulating a Perforating Gastric Ulcer.

DR. W. J. THOMPSON and MR. CONWAY DWYER described a case of the above. [It will be found in Vol. CXIX., page 252.]

DR. R. TRAVERS SMITH asked if any cause had been found why the coil of the jejunum should have become gangrenous.

DR. CRAIG said that so long as liver dulness was present he would hesitate in giving a diagnosis of perforation into the abdominal cavity, though rupture might take place into an abscess on the posterior wall of the stomach without the occurrence of this sign.

DR. DOYLE said that it had not been proved that there was no gastric ulcer in this case, though there was certainly no perforation.

DR. KIRKPATRICK related a recent case diagnosticated as perforated gastric ulcer which had taken place ten hours before being sent to hospital. Although the liver dulness was apparently normal before operation, which was done two hours after admission, a small perforated ulcer was found on the anterior wall, and the contents of the stomach were found in the abdominal cavity.

MR. CROLY spoke.

DR. LITLEDALE asked if the condition was due to an infarction of the intestine. If there was no adhesion found that might be a possible explanation of the condition. He considered that the mesenteric vessels were not sufficiently examined in such cases.

DR. PEACOCKE saw a case under Mr. Gordon's care with acute symptoms like strangulation. They could feel the coil of intestine through the abdominal wall. After laparotomy an almost gangrenous coil of intestine was found, but no bands or adhesions, and the conclusion came to was that a thrombus had formed owing to syphilitic affection of the arteries.

DR. PARSONS said that some eight or ten years ago he had brought before this Section some cases of perforated gastric ulcer coming under his observation while Resident Medical Physician in Sir Patrick Dun's Hospital. He related again some of the particulars of these cases, and observed that he had then emphasised the necessity of early laparotomy in such cases. Personally he knew no point to which he attached more importance in diagnosis than the abolition of the area of hepatic dulness. One other point of importance was a steady increase in the rapidity of the pulse.

The temperature was of no value, as in the worst cases they found subnormal temperature, and in most cases he had seen there was no febrile reaction.

The PRESIDENT (Dr. J. W. Moore) said he remembered a case in the Meath Hospital in which a perforated gastric ulcer produced all the symptoms of empyema, and tapping confirmed this diagnosis. There was also a discharge of pus from the bronchial tubes. The autopsy showed that the abscess was below the diaphragm, which it had perforated.

Dr. THOMPSON, in reply, said that in the most recent work on the subject the absence of liver dulness was stated not to be an absolute sign, neither was the absence of hæmatemesis.

MR. DWYER said that an examination had not been made, as the specimen had been mislaid, but he had no doubt that it was due to a thrombus owing to the sharply defined nature of the gangrenous patches and the absence of adhesions. Mr. Treves had described many cases of that kind.

The Section then adjourned.

LITERARY NOTE.

MESSRS. REBMAN, Limited, announce the following new books as just issued:—"The International Text-book of Surgery," by British and American Authors, in 2 volumes, edited by A. Pearce Gould, M.S., F.R.C.S., of the Middlesex Hospital, and J. Collins Warren, M.D., LL.D., of Harvard Medical School. An entirely new "Cyclopædia of Practical Medicine and Surgery," in 1 volume, illustrated, by 70 special contributors, compiled under the editorial supervision of Drs. George M. Gould and Walter L. Pyle, of Philadelphia. "A Text-book of Pharmacology and Therapeutics, or the Action of Drugs in Health and Disease," by Dr. Arthur R. Cushny. "The Gross and Minute Anatomy of the Central Nervous System," by Dr. H. C. Gordinier. "Gould's Year-book of Medicine and Surgery for 1900," in 2 volumes (medicine and surgery). The same publishers also announce the following for early publication:—Volume II. of Prof. Deaver's work on "Surgical Anatomy." A new book by Mr. Marmaduke Shield, entitled "The Causes, Consequences, and Treatment of Nasal Obstruction." "A Treatise on the Theory and Practice of Medicine for Students and Practitioners," by Dr. H. W. Syers, of the Great Northern Central Hospital, London.

SANITARY AND METEOROLOGICAL NOTES.

Compiled by SIR J. W. MOORE, B.A., M.D. Univ. Dubl.;
P.R.C.P.I.; F. R. Met. Soc.;

Diplomate in State Medicine and ex-Sch. Trin. Coll. Dubl.

VITAL STATISTICS

For eight weeks ending Saturday, July 14, 1900.

IRELAND.

TWENTY-THREE TOWN DISTRICTS.

The average annual death-rate represented by the deaths—exclusive of deaths of persons admitted into public institutions from without the respective districts—registered in the week ending July 14 in the Dublin registration area and the twenty-two principal provincial urban districts of Ireland was 17·3 per 1,000 of their aggregate population, which, for the purposes of these returns, is estimated at 1,062,188. The deaths registered in each of the past eight weeks, and during the whole of that period, in the several districts, alphabetically arranged, corresponded to the following annual rates per 1,000 :—

Towns, &c.	Week ending				Average Rate for 4 weeks	Towns, &c.	Week ending				Average Rate for 4 weeks
	May 26	June 2	June 9	June 16			May 26	June 2	June 9	June 16	
23 Town Districts	23·3	20·0	19·4	21·0	20·9	Limerick -	25·3	18·2	16·8	15·4	18·9
Armagh -	21·4	28·5	14·3	14·3	19·6	Lisburn -	21·3	25·7	12·8	38·3	24·5
Ballymena	11·3	5·6	11·3	11·3	9·9	Londonderry	17·3	18·8	11·0	20·4	16·9
Belfast -	20·5	19·5	17·9	20·5	19·6	Lurgan -	22·8	27·4	18·2	13·7	20·5
Carrickfergus	17·5	35·1	23·4	5·8	20·5	Newry -	20·1	28·2	8·1	16·1	18·1
Clonmel -	14·6	9·7	19·5	9·7	13·4	Newtownards	5·7	5·7	51·0	11·3	18·4
Cork -	28·4	20·8	24·2	20·1	23·4	Portadown -	12·4	12·4	12·4	18·6	14·0
Drogheda -	22·8	15·2	22·8	11·4	18·1	Queenstown	11·5	28·7	5·7	11·5	14·4
Dublin (Reg. Area)	25·2	21·9	23·1	23·9	23·5	Sligo -	15·2	15·2	5·1	15·2	12·7
Dundalk -	16·8	12·6	25·1	16·8	17·8	Tralee -	33·6	22·4	11·2	16·8	21·0
Galway -	34·0	0·0	3·8	56·7	23·6	Waterford -	29·8	17·9	11·9	13·9	18·4
Kilkenny -	70·8	28·3	18·9	23·6	35·4	Wexford -	27·1	13·5	22·6	13·5	19·2

TOWNS, &c.	Weeks ending				Average Rate for 4 weeks	TOWNS, &c.	Weeks ending				Average Rate for 4 weeks
	June 23	June 30	July 7	July 14			June 23	June 30	July 7	July 14	
23 Town Districts	17·8	20·4	18·4	17·3	18·5	Limerick -	14·0	30·9	18·2	19·6	20·7
Armagh -	0·0	7·1	28·5	14·3	12·5	Lisburn -	4·3	8·5	21·3	21·3	13·9
Ballymena	16·9	33·8	0·0	28·2	19·7	Londonderry	23·6	11·0	29·8	12·6	19·3
Belfast -	18·9	19·8	18·3	15·7	18·2	Lurgan -	31·9	13·7	18·2	9·1	18·2
Carrickfergus	5·8	11·7	5·8	17·5	10·2	Newry -	8·1	8·1	32·2	24·1	18·1
Clonmel -	9·7	14·6	29·2	9·7	15·8	Newtownards	0·0	28·3	17·0	11·3	14·2
Cork -	18·7	21·5	18·7	17·3	19·1	Portadown	24·7	0·0	12·4	18·6	13·9
Drogheda -	7·6	7·6	7·6	22·8	11·4	Queenstown	17·2	23·0	17·2	5·7	15·8
Dublin - (Reg. Area)	17·9	20·0	17·7	18·2	18·5	Sligo -	0·0	15·2	15·2	0·0	7·6
Dundalk -	0·0	41·9	12·6	16·8	17·8	Tralee -	78·3	39·2	11·2	11·2	35·0
Galway -	18·9	49·1	45·3	45·3	39·6	Waterford	17·9	13·9	6·0	17·9	13·9
Kilkenny -	14·2	37·8	37·8	33·0	30·7	Wexford -	18·1	36·1	9·0	18·1	20·3

The deaths from the principal zymotic diseases registered in the 23 districts during the week were equal to an annual rate of 1·2 per 1,000—the rates varying from 0·0 in eighteen of the districts to 11·3 in Galway—the 12 deaths from all causes registered in that district comprising 3 from measles. Among the 108 deaths from all causes registered in Belfast are 2 from whooping-cough, one from diphtheria, 7 from enteric fever, and one from diarrhoea.

DUBLIN REGISTRATION AREA.

In the Dublin Registration Area the births registered during the week amounted to 218—108 boys and 110 girls; and the deaths to 127—65 males and 62 females.

DEATHS.

The deaths, which are 21 under the average number for the corresponding week of the last ten years, represent an annual rate of mortality of 18·9 in every 1,000 of the population. Omitting the deaths (numbering 5) of persons admitted into public institutions

from localities outside the area, the rate was 18·2 per 1,000. During the 28 weeks ending with Saturday, July 14, the death-rate averaged 31·9, and was 3·2 over the mean rate for the corresponding portions of the ten years 1890–1899.

There were 13 deaths from zymotic diseases, or 3 over the number which took place in the preceding week, but 7 under the average for the 28th week of the last ten years. This number includes 4 deaths from whooping-cough, 3 deaths from cerebro-spinal meningitis, 2 deaths from enteric fever, and one each from simple continued fever and diarrhœa.

Tubercular disease caused 38 deaths—namely, 31 deaths from pulmonary tuberculosis, one from mesenteric disease, 5 deaths from tubercular meningitis, and one death from scrofula.

Diseases of the respiratory system caused 20 deaths, which number is 5 in excess of those occurring in the preceding week, but one under the average number for the 28th week of the last ten years.

Convulsions caused 5 deaths—2 in children under one month, 2 in children over one month and under one year, and one in a child 2 years of age. Three deaths were due to apoplexy, and 3 deaths to other diseases of the brain and nervous system.

Five accidental deaths and one death from suicide were registered.

STATE OF INFECTIOUS DISEASE IN DUBLIN.

(1.) CASES OF INFECTIOUS DISEASES NOTIFIED TO THE PUBLIC HEALTH COMMITTEE OF THE CORPORATION.

Sir Charles A. Cameron, C.B., Medical Superintendent Officer of Health for the City of Dublin, has furnished the following information regarding the number of cases of Infectious Diseases in the City of Dublin, notified under the “Infectious Diseases (Notification) Act, 1889” :—

Week ending 23rd June,	...	...	50 cases.
„ 30th June	...	...	38 cases.
„ 7th July,	...	...	61 cases.
„ 14th July,	...	...	66 cases.

Of the 66 cases notified, 21 were erysipelas, 17 enteric fever, 13 scarlatina, 6 measles, 2 meningitis,^a and 7 continued fever.

^a This disease has been placed upon the list of compulsory notifiable diseases by the Public Health Committee of the Corporation, from the 9th of May for a period of four months, upon the recommendation of Sir C. Cameron, C.B.

(2.) CASES OF INFECTIOUS DISEASES UNDER TREATMENT
IN DUBLIN HOSPITALS.

Twelve cases of scarlet fever were admitted to hospital, 9 were discharged, and 62 remained under treatment at the close of the week. This number is exclusive of 19 patients in Beneavin, the Convalescent Home of Cork-street Fever Hospital.

Ten cases of enteric fever were admitted to hospital, and 10 cases were discharged, leaving 40 under treatment at the close of the week.

One case of typhus fever was admitted to hospital, and 3 cases remained under treatment at the close of the week.

Four cases of diphtheria and 12 cases of measles also remained under treatment; no death from the above-mentioned diseases occurred in hospital during the week ending July 14th, 1900.

ENGLAND AND SCOTLAND.

The mortality in thirty-three large English towns, including London (in which the rate was 14·2), was equal to an average annual death-rate of 15·6 per 1,000 persons living. The average rate for eight principal towns of Scotland was 17·2 per 1,000, the rate for Glasgow being 17·4, and for Edinburgh 18·1.

METEOROLOGY.

Abstract of Observations made in the City of Dublin, Lat. 53° 20' N., Long. 6° 15' W., for the Month of June, 1900.

Mean Height of Barometer, - - -	29·878 inches.
Maximal Height of Barometer (1st, at 9 a.m.),	30·426 „
Minimal Height of Barometer (24th, at 9 p.m.),	29·371 „
Mean Dry-bulb Temperature, - - -	57·6°.
Mean Wet-bulb Temperature, - - -	54·2°.
Mean Dew-point Temperature, - - -	51·1°.
Mean Elastic Force (Tension) of Aqueous Vapour,	·374 inch.
Mean Humidity, - - - - -	79·6 per cent.
Highest Temperature in Shade (on 15th),	71·6°.
Lowest Temperature in Shade (on 23rd),	47·2°.
Lowest Temperature on Grass (Radiation) (on 2nd), - - - - -	42·5°.
Mean Amount of Cloud, - - - - -	69·2 per cent.
Rainfall (on 22 days), - - - - -	3·169 inches.
Greatest Daily Rainfall (on 20th), - - -	·521 inch.
General Directions of Wind, - - - - -	W., S.W.

Remarks.

Opening with anticyclonic conditions and fair, though cool, weather, June, 1900, belied its early promise and proved extremely unsettled, showery, and thundery. The mean temperature was, indeed, about one degree Fahrenheit above the average, but the amount of cloud was large (69·2 per cent.). Bright sunshine prevailed for only 158 hours, and torrential showers fell on many consecutive days, being frequently accompanied by thunder and lightning. Midsummer's Day, Sunday, the 24th, was inclement in the extreme, a downpour of rain accompanied by south-easterly to south-westerly gales being experienced in Ireland.

The rainfall in the city (3·169 inches) and in the Phoenix Park (3·700 inches) was much heavier, though not more frequent, than that at Killiney (2·29 inches) and in the Co. Wicklow (Greystones, 1·810 inches; Newcastle, 2·149 inches). This depended on the direction of the prevalent winds (W. and S.W.) and on the fact that the rainfall was made up of "splashes" rather than of "soaks."

In Dublin the arithmetical mean temperature (58·9°) was above the average (57·8°) by 1·1°; the mean dry-bulb readings at 9 a.m. and 9 p.m. were 57·6°. In the thirty-five years ending with 1899, June was coldest in 1882 (M. T.=55·8°) and in 1879 ("the cold year") (M. T.=55·9°). It was warmest in 1887 (M. T.=62·3°); in 1896 (M. T.=61·4°); and in 1899 (M. T.=61·3°).

The mean height of the barometer was 29·878 inches, or 0·039 inch below the corrected average value for June—namely, 29·917 inches. The mercury rose to 30·426 inches at 9 a.m. of the 1st, and fell to 29·371 inches at 9 p.m. of the 24th. The observed range of atmospheric pressure was, therefore, 1·055 inches.

The mean temperature deduced from daily readings of the dry-bulb thermometer at 9 a.m. and 9 p.m. was 57·6°, or 5·7° above the corresponding M. T. for May, 1900. Using the formula, *Mean Temp.* = *Min.* + (*Max.*—*min.* × ·465), the value was 58·4°, or 1·2° above the average mean temperature for June, calculated in the same way, in the twenty-five years, 1865–89, inclusive (57·2°). The arithmetical mean of the maximal and minimal readings was 58·9°, compared with a twenty-five years' average of 57·8°. On the 15th the thermometer in the screen rose to 71·6°—wind, S.W.; on the 23rd the temperature fell to 47·2°—wind, N.W. The minimum on the grass was 42·5° on the 2nd.

The rainfall amounted to 3·169 inches on as many as 22 days. The average rainfall for June in the twenty-five years, 1865–89,

inclusive, was 1·817 inches, and the average number of rainy days was 13·8. The rainfall, therefore, and the rainy days were far above the average. In 1878 the rainfall in June was very large—5·058 inches on 19 days; in 1879, also, 4·046 inches fell on 24 days. On the other hand, in 1889, only ·100 inch was measured on 6 days. In 1887, the rainfall was only ·252 inch, distributed over 5 days. In 1899, 1·643 inches fell on only 8 days.

High winds were noted on only 3 days, and the force of a gale was attained only on the 24th. The atmosphere was foggy on the 5th. Solar halos were seen on the 3rd and 4th. Temperature reached or exceeded 70° in the screen on only 2 days, compared with 17 days in 1887, only 1 day in 1888, and 14 days in 1899. Thunderstorms occurred on the 6th, 12th, 13th, 14th and 19th. Thunder was heard on the 7th, and lightning was seen on the 5th, 11th and 25th. Hail fell on the 14th.

The rainfall in Dublin during the six months ending June 30th amounted to 13·090 inches on 14 days, compared with 11·295 inches on 95 days in 1899, 12·115 inches on 98 days in 1898, 13·950 inches on 113 days in 1897, 7·854 inches on 84 days in 1896, 12·282 inches on 80 days in 1895, 14·361 inches on 109 days in 1894, 9·624 inches on 78 days in 1893, 11·770 inches on 97 days in 1892, 8·748 inches on 77 days in 1891, only 6·741 inches on 67 days in 1887, and a twenty-five years' average of 12·313 inches on 95·4 days.

At Knockdolian, Greystones, Co. Wicklow, the rainfall was only 1·810 inches on 21 days. Of this quantity ·220 inch fell on the 12th and again on the 21st. The total fall since January 1 has been 18·191 inches on 105 days, compared with 19·510 inches on 95 days in the first six months of 1899, 13·500 inches on 88 days in those of 1898, 18·125 inches on 106 days in those of 1897, 7·356 inches on 61 days in the same period of 1896, 14·270 inches on 67 days in 1895, 17·381 inches on 96 days in 1894, and 11·776 inches on 75 days in 1893.

The rainfall at Cloneevin, Killiney, Co. Dublin, amounted to 2·29 inches on 23 days. The greatest fall in 24 hours was ·27 inch on the 24th. The average rainfall for June in the 15 years, 1885–1899, was 1·750 inches on 12·4 days. In 1897, 3·59 inches fell on 20 days, in 1898, 2·03 inches fell on 15 days, in 1899, 2·47 inches fell on 10 days. Since January 1st, 1900, 13·82 inches of rain have fallen at this station on 113 days, compared with 13·10 inches on 97 days in the corresponding six months of 1898, and 13·62 inches on 89 days in the first half of 1899.

At the National Hospital for Consumption, Newcastle, Co.

Wicklow, the rainfall was 2·149 inches on 22 days, compared with 3·748 inches on 8 days in June 1899, 2·459 inches on 14 days in June, 1898, and 4·078 inches on 15 days in June, 1897. On the 22nd, ·336 inch was measured. The maximum temperature in the shade was 69·1° on the 15th, the minimum temperature in the shade was 45·0° on the 1st. At this station the rainfall for the six months ending June 30 amounted to 16·904 inches on 103 days, compared with 18·639 inches on 91 days in the first six months of 1899, 14·918 inches on 88 days in the same period of 1898, and 18·372 inches on 102 days in that of 1897.

At the Railway Hotel, Recess, Connemara, Co. Galway, the rainfall in June, 1900, was as much as 7·676 inches on 22 days, the measurement on the 24th amounting to 2·210 inches. The month was very wet and boisterous at that station, and a south-westerly gale of great violence occurred on the 24th—Midsummer's Day.

TREATMENT OF APPENDICITIS.

M. BROCA (*Rev. Med. de Paris*).—Absolute rest, opium internally, ice over iliac fossa, injections of sweet oil, and rectal feeding in severe cases. M. Broca does not recommend operation until it is plainly indicated. From 1892 to 1895 he operated on all cases of appendicitis on their admission. Of his 67 cases, 45 were cured and 22 died, showing a mortality of 33 per cent. In 1896 the influence of Jalaquier cooled his ardour for operation, and he temporised, even in severe cases, with the result that of 27 cases he lost but 4, reducing the mortality to 13½ per cent. In 1897, following the same practice, he had 28 cases with 3 deaths—a mortality of 10·7 per cent. In 1898 the mortality rose, for of 16 cases he lost 3—a death-rate of 18·8 per cent. The following year (1899) the death-rate fell to 10·8 per cent.—37 cases being treated with a loss of only four patients. The marked increase in the mortality in 1898 M. Broca ascribes to the fact that the patients on admission were in a hopeless condition. He notes that in the same hospital during the same year 28 cases were operated on, the death-rate reaching 50 per cent.

PERFORATING TYPHOID ULCER.

J. E. PLATT says the incision should be made on the right side, the ulcer inverted without cutting away its edges, the abdomen flushed, and a drain inserted.

PERISCOPE.

EPITHELIOMA OF THE FACE IN A YOUNG SUBJECT.

At a meeting of the Société des Sciences Médicales de Lyon, M. Jermier presented a patient with epithelioma of the cheek bone, which necessitated operation. This subject was brought before the medical profession in a thesis by M. Goy (Lyon, 1898), "Epithelioma of the Face in Young Subjects." The disease in the case of this patient commenced as a pimple on the right cheek bone when he was thirteen years of age. From time to time the patient rubbed and scratched the pimple until it became an open sore. This sore gradually became larger, until it was as large as a sixpenny piece. It neither got smaller nor bigger until he was called up as a conscript, when he was transferred to Saint Etienne, and put on a course of treatment (arsenical). He entered the Hotel Dieu de Lyon in July, 1898, with an ulcer on the right cheek bone as big as the palm of the hand. It gave off a disagreeable smell, was sanious, tumid, and rigid. Its borders were hard, with deep overhanging edges. On the suspicion of syphilis the patient was treated with iodide of potassium, hypodermics of calomel, and mercurial frictions. He notably lost flesh; but there was no sign of any improvement in the ulcer. In December an attack of periostitis of the upper jaw bone of the right side was treated by a free section, which extended to the external angle of the eye. To cover the surgical wound, two flaps were taken from the neck, and a third from the temporal region. After fixing these flaps in position there still remained uncovered about the size of a five-franc piece, which was allowed to granulate a portion of the wound and cicatrise. The microscope showed that the ulcer consisted of epithelial cancer. Fourteen months afterwards there was a slight return of the disease, and the contraction of tissue due to the cicatrization had caused ectropium. A plastic operation cured the ectropium, and the hairs which grew on the flaps brought from the neck were destroyed at their roots by electrolysis.—*Lyon Médical*.

CONGENITAL ATROPHY OF THE BICEPS.

At a meeting of the Société Nationale de Médecine de Lyon, M. Bernard presented a soldier having congenital atrophy of the biceps of his right arm. He had suffered from malaria in Crete and Tonkin, for which he received the classic treatment—quinine—and

quickly got well. During his life in the army the lesion was never detected, and it in no way interfered with the movements of the forearm, although the muscle was represented by nothing but a thin tendon. M. Bernard considers that the atrophy is due to a central nervous lesion. Professor Pestort, who also examined the patient, accepts M. Bernard's view as to this opinion.—*Lyon Médical*.

CONDITIONS SIMULATING APPENDICITIS.

DR. JANEWAY (*Medical Record*) at a meeting of the Practitioners' Society, New York, enumerates the following conditions which may be mistaken for appendicitis:—(1) Neuralgia affecting the lower abdominal nerves on the right side; (2) pneumonia or pleurisy of the right side; (3) renal colic of the right side; (4) hydronephrosis of the right side; (5) movable kidney of the right side; (6) perforation of latent duodenal or other intestinal ulcers; (7) tuberculous ulcers of the cæcum; (8) cancer of the cæcum; (9) faecal impaction; (10) narrowing of the hepatic flexure of the colon; (11) typhoid fever in its beginning or during a relapse; (12) follicular tonsillitis; (13) salpingitis; (14) sapræmia. These cases sufficiently illustrate the need of careful thought lest we fall into the error of considering every case with pain and tenderness in the right iliac fossa as appendicitis.

POST-PARTUM HÆMORRHAGES, CURED BY SIMPLE DRAINAGE AND FOLLOWED BY PSYCHOSES.

We can admit, says C. Zalackás, a post-puerperal psychosis. The psychoses following gynæcological operations are very exceptional, and when they exist we usually have to do with women predisposed or already affected with insanity. There is no connection between the psychoses and the operative procedure. Psychic troubles can be produced as well after any other operation as after ovarian castration. The operation never of itself produces madness. There exists a psychic state before operation called phobia, which may persist after the operation. In psychoses at precocious menopause a certain predisposition is at work. The fear of mental perturbation should not deter one from operating when the indications are clearly present.—*Le Progrès Médical*.

OPERATION IN GASTRIC ULCER.

BIDWELL says operation in gastric ulcer should be done (1) in all cases of perforation at the very earliest possible moment—also in subphrenic abscess; (2) in cases of hæmorrhage (*a*) when there is continual oozing of blood, especially if the stomach be dilated, and

(b) in cases of repeated severe hæmorrhage; (3) in cases in which there is severe pain and vomiting unaffected by treatment, and which is producing progressive emaciation; (4) in case of dilatation of the stomach from contraction within or from adhesions outside the stomach.

BATHING IN THE TYPHOID FEVER OF CHILDHOOD.

E. AUSSET (*Journal de Médecine de Paris*) finds a series of cases in children more severe than the adult form. He never finds at autopsy in an infant typhoid which has escaped recognition during life. In forty-five cases there were four before the fifth year, twenty-nine between five and twelve years, and twelve between the twelfth and fifteenth years, with two deaths in each series. The author believes in systematic bathing from the very onset; the temperature should be regulated for each case; the younger the child the less cold is the bath. It is unnecessary to give baths as cold as the pure Brand method calls for.

AN ATAXIC INFANT.

At a meeting of the Société des Sciences Médicales de Lyon, M. Lévy presented (in the name of M. Lannois) a young child of three years and a half old. The infant was born in the Charité. Its antecedents were imperfectly known. The mother had no sign of syphilis, and the labour was natural. Trembling of the muscles appeared at two years of age. The upright position could not be retained by the child unless there was something to hold by. When moving, the limbs were kept widely apart, and were jerked irregularly; at the same time the superior extremities were constantly moving in a tremulous manner. When walking the patient moved as if drunk, the legs wide apart, and feet seemed to clutch the ground as they were placed on it. There was no loss of muscle force, only the troubles of sensation. Divergent strabismus and nystagmus were present on both sides. The knee-reflexes were exaggerated; there were no contractures, and the sphincters were normal. The patient presented all the phenomena of cerebellar ataxy of Babinski. M. Lannois considers the phenomena were due to an arrest of the development of the cerebellum, followed by atrophy during extra-uterine life.—*Lyon Médical*.