cuticle, but holds the deeper tissues until union is secured, despite moderate tension and even suppuration, should it occur. This, however, is not meant to imply that I neglect in the least the additional safeguard of that most valuable adjunct in plastic surgery—the relaxation suture, whether in the form of the corrugated vulcanite disc with silkworm gut or my wire and clip method.

A METHOD OF APPLYING ANTISEPTICS TO THE DEEPER PARTS OF WOUNDS.

By THOMAS H. KELLOCK, M.C. CAMB., F.R.C.S. ENG.,
CAPTAIN, R.A.M.C. (T.)

THE multitude of septic wounds at the present time, which occupy the attention of so many medical men, has very naturally given rise to the use of a variety of antiseptics and many methods of application. To judge of the efficiency of the former due attention should be paid to the latter, and it is obviously unfair to judge of the value of an antiseptic when it is only applied to the part of a wound that is visible. In every wound of any magnitude there must be recesses and deep parts into which, however well such a wound may be drained, the antiseptic applied in this way never reaches and where organisms may collect and multiply unmolested.

The Carrel method of irrigating wounds has done a good deal to improve their treatment and has drawn attention to the advantage of applying antiseptics to the deeper as well as to the more superficial parts of wounds. The method, however, has certain disadvantages. The tubes generally used are perforated with round holes for some distance from the closed end, and if there be much resistance to the exit of fluid at the deepest part the solution finds its way through the openings nearer the surface of the wound and so the deeper parts are not reached.

These tubes, too, act to a certain extent as drains. Pus finds its way into them through the openings and is washed back into the wound again when fluid is allowed to flow. Another disadvantage of the method is the necessity of keeping the patient still. Movement on his part is apt to cause the tubes to be drawn out of the wounds, and it is not at all an uncommon experience to find tubes that were originally inserted deeply into a wound lying only just under, or even on, the surface.

Attempting to improve on this, I have found the following method of much service. Taking a piece of drainage-tube about 8 or 10 inches in length and of calibre about equal to that of a No. 8 catheter, the end is securely closed by a ligature and a small oblique incision made with sharp scissors as near as possible to the closed end. The incision should cut a small valve with apex directed towards the closed end—the more oblique it is the better—and should at its base divide one-third of the circumference of the tube (see figure, A). This incision will be found to act as a most

efficient valve, opening under very slight pressure from within and preventing any regurgitation of pus or of the irrigated fluid. The tied end should be inserted to the very depth of the wound and the free end of the tube left projecting through the dressings; the number of tubes used must depend on the size and character of the wound, but it will generally be found that one, or at the most two, will suffice.

In cases of a perforating wound of a limb or of a wound with two openings widely separated a very useful modification is to ligature the tube at its centre and cut a valve on each side of the ligature and pointing towards it (B). This is passed through the wound and the irrigation done from both ends. In this way any fluid that is irrigated into the free ends of the tube must certainly find its way to the very bottom of the wound and from there towards the surface.

To secure the tubes in position a useful plan is to make a small independent incision through the skin near the edge of the wound just large enough to grip the tube without occluding its lumen, and the free end of the tube is drawn

through this after the tied end has been passed to the bottom of the wound. The free ends of the tubes are left outside the dressing and the irrigating fluid passed into them by means of a glass syringe. Gravity alone will generally suffice, but, if necessary, a little assistance may be given by using the plunger of the syringe. If only a small quantity of fluid be used at a time the dressings on the wound need only occasional changing. I have been using flavine for this purpose. One to two drachms of a 1 in 1000 solution is injected through the tubes twice in 24 hours, and the results in some very septic cases have been most satisfactory.

A question on which I have not yet been able to satisfy myself completely is whether wounds treated in this way should be drained. If free drainage is provided, the antiseptic fluid very readily finds its way to the surface and is absorbed by the dressings, and so is not able to remain in contact with the real surface of the wound. I have sometimes thought that the best results have come about when the drainage is not free. The walls of the wound then remain open and the mixture of pus and antiseptic fluid is brought into contact with the suppurating surface at all points.

Clinical Notes:

MEDICAL, SURGICAL, OBSTETRICAL, AND THERAPEUTICAL.

MEMBRANOUS CONJUNCTIVITIS FOLLOWING A SQUINT OPERATION.

By S. JOHNSON TAYLOR, M.B. EDIN.

RECENTLY I performed an advancement cum tenotomy operation for a convergent squint, in a girl aged 11 years, followed by a complication which I believe to be very rare. I used three stitches for the advancement and one for the tenotomy part, which I did, as usual, by the open method. All went well. Two days later the upper eyelid was a little more swollen and glazed than is usual and the advancement-wound region did not look quite nice; this was soon succeeded by a definite dirty membranous area which I diagnosed clinically as diphtheritic, the tenotomy-wound area quite escaping. I at once had antitoxin injected hypodermically and used quinine lotion freely and frequently, as advised by Sir John Tweedy in "Diphtheritic Conjunctivitis," and slowly the condition cleared up; the stitches became very loose and either were cut out or came away. The sloughy membranous patch involving the tendon and distal end of the external rectus was loth to disappear and did so very gradually, leaving a swollen granulation patch and a convergent squint as marked as before the operation. The general systemic disturbance was very slight. No bacteriological examination was made or was necessary, the clinical condition being quite sufficient to label the disease. The patient had diphtheria last Christmas and two other of the children had it, and the baby with cleft palate has been for some weeks—before the operation—and still is in our isolation hospital as a carrier of the complaint.

The ground behind the patient's house is soaked with sewage, there being no drainage system there. I have mentioned this case to several of my London colleagues, but can find no parallel thereto. The one attack of general diphtheria did not confer any immunity against the local affection. I venture to state that this complication of a squint operation is excessively rare if not unique.

Norwich.

THE CLINICAL MANIFESTATIONS OF A DOUBTFUL CASE.

By M. J. PETTY, M.B. CAMB., F.R.C.S. ENG.

THE following case is recorded as a problem in diagnosis.

Patient, girl, 6 years of age. No evidence of other cases in family. Mother and father both healthy. When patient was 2 years of age a slight fit, with loss of consciousness, occurred; apparently complete recovery. Four months later the child began to drag right foot along the ground, since then progressively getting worse till whole of right side of body was noticed to be affected. Occasional

vomiting occurred. Bowels always regular. A week ago she had another fit, leaving an apparent palsy of all the eye muscles. Vomiting took place occasionally. I was called in to see the child, as it was supposed that symptoms, &c., were due to a tumour near the quadrigeminal bodies.

State.—The child in poor general condition, lying in bed on right side, and complaining of localised pain over left occipital region which had been present for four days. Mental condition quite clear, but speaks in a monotone and answers quickly and rationally. The right hand is held in left hand, and moved about by latter. Head measurement 50½ cm., regular in shape. Examination of cranial nerves: 1. Smell sense delicate—not the slightest impairment. 2. No change in optic nerves and fundi; fields of vision normal. Vision normal. 3, 4, 6. Right side: Pupil dilated and movements in all directions impaired; moves more than left. Slight exophthalmos. Left side: Eye apparently immovable, but on persuasion child moved eye slightly in all directions. Reaction to light good; reaction to accommodation could not be determined. 5. No loss of sensation. 7. Supranuclear lesion on right side; left side apparently normal. 8. Hearing normal on both sides. 9. Taste normal. 11. Unaffected. 12. Deviation to right side when tongue is protruded.

Right arm: Kept flexed over chest. Some active flexion and extension of elbow and distal joints. Grasps hand well but weakly. Tendon reflexes absent. No loss of sensation. Arm, forearm, and hand muscles definitely atrophied. Left arm: Loss of tendon reflexes, but otherwise normal in all its movements, not showing an intention tremor. Trunk: Normal sensation. Abdominal reflexes right side are absent, but are brisk on left side. Sphincters normal. Right lower limb: Definite paresis compared with left. Knee-jerk absent; other tendon-jerks also absent. No ankle clonus. Hallux erectus present. Definite disuse atrophy present; limb is 2.5 cm. shorter than its fellow. No loss of sensation. Left lower limb: Stronger than right. Movements well coördinated. Knee-jerk and other tendon-jerks absent. Ankle clonus absent. Babinski's sign: Extensor response.

No scoliosis. Definite contracture of post-vertebral muscles in occipital region. Wassermann reaction negative (two occasions). Cerebro-spinal fluid not examined. Urine normal. T. about 99°F. at night. P. 120. Heart, lungs, and abdominal viscera normal.

Fits, intense headache, occasional vomiting, and the eye symptoms suggested intracranial pressure, but of this there was no evidence on further investigation. Friedreich's disease, with which much of the condition agrees, is excluded by the absence of definite ataxia and of hereditary transmission.

Buenos Aires.

PETROLEUM IN THE TREATMENT OF PYORRHOEA ALVEOLARIS.

BY H. J. CORIN, L.D.S.

THE treatment of pyorrhœa alveolaris should be undertaken by the dentist only in conjunction with the patient's medical adviser, since pyorrhœa is now commonly acknowledged to be aggravated by, if not actually due to, causes other than dental. It is, for instance, in many cases primarily induced by the absorption of toxins from a septic small intestine, a condition which obviously needs skilled medical attention.

For such a condition purified petroleum is used widely and successfully, both in those cases where surgical interference is necessary, and also in those more common instances where no operation is required. Most probably the therapeutic action of petroleum is known to the vast majority of medical men; but I would like to emphasise the caution given some time ago in this journal after various investigations,¹ that only the purest oil should be administered.

In a case where there was only a slight discharge of pus from the gums, one course of local treatment, the use of an effective mouth-wash and the patient ordered to take petroleum emulsion, produced a complete cure. The patient was unable to come to town for six weeks after treatment, but seen at the end of this period there was absolutely no trace of discharge.

For long-standing cases where there is a large discharge I have found petroleum with guaiacol to give wonderful results in clearing up the conditions quickly, and when there is acute gingivitis petroleum with salol is useful. After my experience I can confidently recommend this method of treatment as effective in many instances where other methods have failed.

Old Burlington-street, W.

¹ Vide THE LANCET, Oct. 2nd and 16th, 1915.

Reviews and Notices of Books.

Organic to Human: Psychological and Social. By HENRY MAUDSLEY, M.D. Lond. London: Macmillan and Co., Limited. 1916. Pp. 386. Price 12s. net.

THIS latest work by Dr. Maudsley is fully worthy of the high reputation that has for long placed him in the front rank of contemporary scientific thinkers and writers on mental processes in health and disease. His first book, the "Physiology and Pathology of Mind," published hard on half a century ago, showed his mastery of all then available knowledge of the structure and functions of the human nervous system, no less than his signal capacity for thinking scientifically. It contained, moreover, the germs of much that its author has since from time to time given to us in his well-known works expanded and illustrated by further experience and mature reflection. Even as Dr. Maudsley's earliest work drew the attention of philosophical thinkers more than that of practising medical men, so probably will this latest book appeal mainly to the scientific student of psychology and sociology whether without or within the ranks of the medical profession: It must, indeed, be considered chiefly as an important contribution to the still vexed controversy between the warring schools of "materialists" or "mechanists" on the one hand, and of "vitalists" or "spiritualists" on the other. Justly pre-eminent though Dr. Maudsley is as an authority on the subject of insanity, and much as his other work owes to his practical knowledge and experience in this special study, his most enduring title to fame will be that of the scientific philosopher who in his day and generation consistently and persistently upheld the principle of the unity of the human organism and its continuity with the rest of Nature's processes.

"A true psychology" (says Dr. Maudsley in the first chapter of the book before us) "must be founded on a clear and distinct recognition of the essential continuity and constant flux of things in which all so-called causes are effects and all so-called effects causes: mind be conceived in thought as a part of the nature which its organisation is in fact. Continuity of motion without end, the perpetual flux which such old Greek philosophers as Democritus, Heraclitus, and Epicurus insisted on, and Lucretius poetically expounded, that is the fundamental fact: the world never the same for an instant, nor two minds ever the same, nor any mind ever the same for a single hour."

Again:—

"Into the composition of every idea, of every mood, of every act, of every reflection, every organ and every gland of the body has entered essentially; the minutely different structures and physico-chemical changes and exchanges of the parts in different persons producing different temperaments, and the signally special organs of male and female bodies different mental qualities requiring their special direction of development and function. To deride the notion of mechanism as anywise applicable to life and life in mind is the natural error of him who, ignorant of the exquisitely subtle physical and chemical processes at work in the simplest vital substance, thinks grossly only of inert matter and its mechanism as exemplified in the structure of a wheel-barrow or a steam-engine."

One more quotation must suffice for the purpose of indicating some of the main lines of thought in this book:—

"That all positive intelligence is sense-evolved, though true, is not the whole truth: motor reactions contribute as essentially as sensory impressions to the construction of a mental organisation and fitly respond to them in the complex composition of ideas and judgments. The old saying, *Nihil in intellectu quod non prius in sensu* might indeed be fitly supplemented by the addition of *et in motu*."

It is clearly apparent that no adequate criticism of the main arguments of this book is possible within the limits of this notice. Even to attempt it would involve a useless plunge into a troubled sea of dispute. But this much may be said in some justification of the high degree of value which we attribute to it. The least biassed thinker on matters psychological, who is dissatisfied with any theory concerning mental phenomena which involves the importation of a purely imagined and unverifiable agency in explanation, but still may find some difficulty in regarding such phenomena as natural products of organisation, is at least likely to discover, after carefully studying Dr. Maudsley's book, that part, if not all, of his difficulty is lessened or removed. The case for