Springer

PLANTS PEOPLE
POSSIBILITIES

Cayman Islands

Author(s): W. Fawcett

Source: *Bulletin of Miscellaneous Information (Royal Botanic Gardens, Kew)*, Vol. 1888, No. 19 (1888), pp. 160-163

Published by: Springer on behalf of Royal Botanic Gardens, Kew

Stable URL: <http://www.jstor.org/stable/4115440>

Accessed: 27-06-2016 06:11 UTC

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at
<http://about.jstor.org/terms>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

Springer, Royal Botanic Gardens, Kew are collaborating with JSTOR to digitize, preserve and extend access to *Bulletin of Miscellaneous Information (Royal Botanic Gardens, Kew)*

“(3.) I have never attempted to propagate it by seed, but always by roots. When a clump or tuft is dug out, it may be divided into as many small divisions of roots as one pleases, and these are put down again in rows, about three feet from one another, and the same interval between each root planted. It will yield a very trifling return the first two years, but by the third or fourth year, when the roots have spread and multiplied, it gives a *good* crop. The plot on which it is planted must be kept free from other grass. When it is seven or eight years old, the roots should be beaten down with wooden mallets, or a plough should be run through them in every direction, and fresh earth thrown over the whole increases the yield. If not treated in this way, it will cease yielding any crop. When grown too old, it must be taken up entirely, re-divided in small bunches of roots, and transplanted to a fresh locality.

“(4.) We bought the grass always in local *hâts* for roofing purposes (*as rope*) before we grew our own, and nowhere in our neighbourhood am I aware of its being cultivated in any but very small patches by a solitary man here or there. It is not cultivated as a source of income or trade, so that I am unable to say where the roots may be bought, or at what price. We got a small quantity of the roots originally from a Hindu village, but by fostering and spreading their cultivation have now a considerable quantity. It should be planted in a *dry* spot, where no water lodges, as experience has shown in one of our stations, where the water oozed up from below and rotted the roots, that it would not grow there. A sloping site is probably the best.

“When we first started the mission here we had to pay Rs. 4 a maund in the *hâts* for the grass, in order to twist it into rope or string, and it was the having to pay so much that led me to try and cultivate it ourselves. The grass runs to seed in the hot months, shortly before the rainy season, but these must be cut off and removed, or the crop will deteriorate.”

Besides a great variety of native names, Bhabur grass has, from the difficulty of exactly ascertaining its affinities, received an almost equal number of botanical ones. Under the name of *Pollinia eriopoda*, it is discussed in the Journal of the Linnean Society (vol. xx., pp. 409, 410), and it is figured and described in Hooker's *Icones Plantarum* (tab. 1773) as *Ischaemum angustifolium*, the name finally assigned to it by Hackel, the most recent monographer of Grasses. To the courtesy of the Bentham Trustees, we are indebted for the use of the accompanying plate.

LII.—CAYMAN ISLANDS.

The Cayman Islands consist of the Grand Cayman, Little Cayman, and Cayman Brac. They are situated in the Caribbean Sea, about 200 miles to the west of Jamaica, and are included under the Government of that Colony. Grand Cayman is 17 miles long, and varies in breadth from 4 to 7 miles. The coast is in some parts bold and rock-bound, but with no elevation exceeding 150 feet. The island is described as well wooded in parts and producing dyewoods and mahogany. The cultivated products of the soil are similar to those of Jamaica, and consist of coconuts, sugar cane, and ground provisions. Amongst the curiosities of Grand Cayman are a cave at Bodden Town, which extends some hundreds of yards under the sea, and a natural cistern of unknown depth, containing clear, sweet water, at a locality called East End. This cistern measures

about 50 feet across, and is situated in the middle of a cliff of flinty rock.

Little Cayman is 9 miles long and about a mile broad. Cayman Brac is 10 miles long, and about a mile broad. Both these islands lie in a north-easterly direction from Grand Cayman, and distant about 70 miles.

The Cayman Islands are so far out of the regular traffic of steamers that they are seldom visited. The present population of Grand Cayman is about 4,000. Little Cayman contains only 35 inhabitants, while the population of Cayman Brac is said to be about 300 whites and 30 blacks.

It is a remarkable characteristic of the inhabitants of these tropical islets that they are a temperate, strong, tall, healthy-looking people, chiefly white or coloured. They are doubtless descended from the original settlers of the last century. The proportion of black people (negroes) is comparatively small.

The present Governor of Jamaica, Sir Henry Norman, has twice visited the Cayman Islands. On the occasion of his more recent visit [May 1888] he was accompanied by Mr. William Fawcett, F.L.S., Director of the Botanical Department, whose mission was specially connected with an investigation into a disease which has existed for some time amongst the cocoa-nut palms at Grand Cayman.

We understand that Mr. Fawcett is at present engaged in the preparation of an official report of his visit. In the meantime the following extract from a letter just received from him will afford a first impression of the vegetable resources of these lonely and little known islands :—

Jamaica, 21st May 1888.

I have just returned from a visit with the Governor to the Cayman Islands. We left Jamaica on 7th May, and reached Grand Cayman on the evening of the 8th. The next morning I went on shore, but the rain came down in such torrents that I could not do much. On the 10th I went about seven miles along the coast to the north, examining cocoa-nut palms at various places that were diseased. The next day I went along the south coast about 15 miles and back the same day, and had several diseased cocoa-nut palms cut down for examination. On the 12th I went inland about 2½ miles to the Sound, a shallow lagoon which runs in from the north side of the island, and on Sunday, 13th, did the same. On the 14th we left Grand Cayman, and touched at Little Cayman, landing for a short time, and then went on to Cayman Brac. There are very few people at Little Cayman, about 40; but at Cayman Brac there are between 400 and 500 inhabitants, living by fishing and cultivation of "ground provisions" (yams, &c.) and bananas. Cayman Brac exports a large number of cocoa-nuts, as much, I was told, as 600,000 to 800,000 annually. The shore is lined with cocoa-nut palms, and there is no disease. A very great number of nuts are, however, destroyed by rats.

Grand Cayman is surrounded by coral reefs, and the shore vegetation is what is usual in such places, but at a short distance from the sea there are signs of an indigenous flora, which would well repay detailed examination. My special work being to inquire into the agricultural products of the islands, I could not devote much time to collecting, and such specimens as I have brought back home have suffered a good deal from want of proper facilities for drying, &c. However, as soon as they are dry I will send them to Kew for determination. I found two orchids, which are well worth cultivating, if you have not already got them. One is a species of *Aeranthus*, like *A. funalis*, but larger and finer, I think. The other is a species of *Schomburgkia*, near *S. crispa*,

but the lip is purplish, not pink red. I enclose flowers only for the present, and if you like I will send some of the living plants, as well as dried specimens.* From what I saw of the interior the soil appeared to be formed by the disintegration of limestone, this red clay and the limestone rock being of the same character as is found to such an extent in Jamaica. Wherever, therefore, it is of any depth it is suitable for cultivation. Mangoes grow well, but they are said not to be so good as some of the Jamaica fruit. Oranges, both sweet and bitter, and a few lemons are produced, and great quantities of limes are exported pickled. Yams, cocoes, sweet potatoes, cassava, pine apples, melons, sugar canes, bananas, guinea grass all grow well. I took a few good pine apple suckers with me, which they were glad to have. The sugar cane looked poor, but they said that it was due to long-continued drought, and that canes grow sometimes from 12 to 15 feet in height. There are a great number of papaw trees, some with thick trunk and several branches; the fruit is usually large. There is a good deal of logwood, fustic, mahogany, and hard-wood timbers, which would appear to prove that the island is something more than a mere coral reef. The people are excellent shipbuilders, and use their native mahogany and other hard timbers in the construction of schooners up to 50 tons, but the planking is now at least imported from the States. These hardwooded trees grow on the north side, but as it was doubtful whether I could go and return in one day, I had to abandon a projected visit. I was sorry, also, not to see the cultivation there, as I heard that the soil was the richest in the island and more cultivation carried on than elsewhere. *Furcræa cubensis* grows, and I believe some attempt has been made to extract the fibre. The Jamaica Keratto (*Agave Morrisii*) is plentiful on all the islands. I saw two trees of coffee growing not far from the sea, and though they were close up to the house, and not pruned, the crop was not a poor one, considering the previous drought. This was probably the first attempt to grow coffee in these islands, and I think that the result justifies planting. The guava and the coco plum grow wild, as in Jamaica.

In Little Cayman there is mahogany, but no cedar (*Cedrela odorata*); and in Cayman Brac there is plenty of cedar and scarcely any mahogany. In Little Cayman there is no logwood nor fustic; at least I was told so by the inhabitants, and I met with none. On Cayman Brac I found fustic growing in the woods, but logwood was said not to grow. The interior of Cayman Brac, within the fringe of coral beach, where the cocoa-nut grows, is limestone, disintegrated into hollows which make walking very difficult and dangerous. Wherever sufficient red clay exists provision grounds are made, where the usual "ground provisions," bananas, and pine apples are grown.

There are considerable deposits of phosphates in these islands, which are being worked at the Grand Cayman.

At Grand Cayman the disease in the cocoa-nut palms appeared somewhere about 15 years ago, and spread throughout the whole island, killing off every single tree. Seed was imported from the mainland, and the inhabitants have been persevering ever since in their efforts to re-establish the former flourishing cocoa-nut walks. Sometimes the trees die while they are quite young, and sometimes a crop is produced

* From an examination of these flowers at Kew, it appears that the *Schomburgkia* is *S. Thomsoniana*, Reich. f., Gard. Chron., 3rd Ser., vol. ii., p. 38 [1887]. The locality of Professor Reichenbach's plant is not stated. The other orchid is probably a new species of *Dendrophylax*, near to the Jamaica plant *Dendrophylax* (*Aëranthus*) *funalis*, Grisebach Flor. Brit. W. Indies, p. 625, but the flower is larger and the spur much longer.

and then the tree is attacked. The disease is quite independent of wind, travelling as often against it as with it, and attacking often the finest trees in a walk. The outer leaves first show signs of the disease, gradually turning yellow; but I invariably found that it was the bud leaves which were affected first at the apex, and as the disease gained ground the outer leaves lost their green colour. Sometimes on cutting the stem right through just above the ground it appeared perfectly untainted and smelt and tasted sweet. At other times, when the disease had made more progress, though the stem section looked white, it smelt and tasted sour, and this occurred even when only the upper portion of the "cabbage" was discoloured. The discolouration was a purplish black, eventually becoming quite black. There was a sour smell as of fermentation in the "cabbage," and, considering all the circumstances, I believe that the disease is due to a bacterium. I had no suspicion of this before starting, as I was told that the disease was the same as the Jamaica scale insect; but even if I had taken material for the purpose of examination, there was not sufficient time to do anything. I have brought back small pieces in alcohol (diluted), but have not yet examined them under the microscope. The disease is evidently not caused by an insect, but after the disease has taken hold of the tree, and decay has set in, various insects may be found, such as ants, a small beetle (called the cocoa-nut fly), and white larvæ, which may be the cocoa-nut fly larvæ. I did not see a trace of the Jamaica beetle or the scale insect.

W. FAWCETT.

LIII.—VALONIA IN CYPRUS.

The following correspondence has passed between this establishment and the Crown Agents for the Colonies on the the subject of acorns shipped to this country by the Government of Cyprus as Valonia :—

ROYAL GARDENS, KEW, to THE CROWN AGENTS.

Royal Gardens, Kew,

GENTLEMEN,

June 4, 1888.

I HAVE the honour to acknowledge the receipt of your letter of May 25th, with enclosures (herewith returned), and accompanied by three samples of acorn-cups sent from Cyprus as Valonia.

2. This important tanning material is in large use in this country, as much as 30,000 tons having been imported in 1885. It consists of the acorn-cups of the Valonia Oak, *Quercus Ægilops*, L., and the commercial supply is derived apparently exclusively from Greece and Asia Minor. Valonia from these two sources is commercially distinguishable, and they are produced by distinct local forms of the Valonia Oak.

3. The acorn-cups sent from Cyprus are not Valonia at all, and do not in any way resemble it. One of the samples, in fact, consisted merely of the acorn-cups of the Holm or Evergreen Oak, so commonly planted in this country. I am afraid there is not much prospect of finding any commercial use for these acorn-cups; but Mr. W. N. Evans, of 66, Stackpole Road, Bristol, takes much interest in possible sources of new tanning materials, and I think if samples were sent to him he might be disposed to see if any use could be found for them.

4. In writing to the Colonial Office, April 20th, 1881, I pointed out that Gaudry, in his report on Cyprus to the French Government, expressed his astonishment not to meet with the Valonia Oak in the