The Journal of the American Medical Association

Published Under the Auspices of the Board of Trustees

VOL. LX, No. 22

CHICAGO, ILLINOIS

NOVEMBER 30, 1912

ARTERIOSCLEROSIS *

THOMAS D. COLEMAN, A.M., M.D.

Visiting Physician to the Augusta City and Lamar Hospitals
AUGUSTA, GA.

Arteriosclerosis is a penalty of our present civilization and is increasing in frequency so rapidly that any study or contribution leading to its better understanding is well worth while.

For a profitable consideration of the subject, certain fundamental facts of the physics of the circulation and the maintenance of cell life must be recalled. The proper nutrition of the different organs of the body is dependent not only on the functional activity of their own cells, but also on the vital activity of the cells making up the cardiovascular mechanism. Not only must both cell elements remain structurally and functionally intact, but there must be a certain intravascular tension in the arteries, arterioles and capillaries.

I believe no physiologist holds that tissue nutrition can be maintained in the absence of tension. This neither negatives nor loses sight of the principle of osmosis, which, in this connection, might stand for chemical tension. A concrete illustration of this may be found in the kidneys. As long as they function normally, even excessive blood-pressure does not cause albumin to be forced through the epithelium lining the tubules in appreciable amount; it is only when the vital activity of the lining cells is interfered with through long-continued strain, toxins or irritants, that albumin and possibly blood appear in the urine. Following and perhaps concomitant with this, connective-tissue changes take place, and the sclerotic condition develops.

A point quite puzzling and one requiring much study is: Why is sclerosis sometimes localized and sometimes general? As a working hypothesis it might be assumed (1) that there is a local defect, either congenital or acquired, in the organ or organs involved; or (2) that the specific toxin has a selective influence which acts on the vasomotor or trophic centers, regulating the blood-supply and nutrition of the part involved, the toxin or irritant acting on the vasomotor mechanism and secondarily on the cells lining the arterial and capillary systems and producing sclerotic changes in the vessels and throughout the areas supplied by them.

Excessive muscular work, either general or local, may impair the integrity of the vascular mechanism. Even in this case it is likely that toxins resulting from tissue destruction may be operating. As to how these act, beyond certain generalizations, it is impossible at this

time to determine with any degree of accuracy. It has been conclusively shown that local sclerosis, without marked sclerotic change in other parts, occurs in the vessels of the brain, and that to this condition is due the fact that so many more ruptures occur in this portion of the arteriocapillary tree than elsewhere; but why? Is it because this portion of the organism is more actively employed during the entire cycle of life or because the vessels in this region are more vulnerable through constant use and grow old while the other parts of the body are yet young?

Certain it is that apoplexy is relatively frequent here, while it is relatively infrequent in other organs, irrespective of age. The importance of this phase of the subject is apparent when one reviews the number of sudden deaths, the frequent paralyses that make the further life of the sufferers a burden to themselves and the state—all the result of sclerosis not infrequently limited to the cerebral arteries. If the preapoplectic symptoms could be interpreted with any degree of accuracy, how many of these tragedies of existence might be averted? Neither the trained tactile sense nor the sphygmomanometer is a sufficient danger-signal. I have had the opportunity of being with a number of patients immediately before, during and after the rupture of cerebral vessels, and beyond some increase in vascular tension and irritability of temper or depression of spirits, there was nothing to give warning of the impending catastrophe.

Again, the vessels of the kidneys are especially liable to undergo sclerotic change producing commonly chronic interstitial nephritis. In these cases also it is desirable to know why the local sclerosis of the renal vessels occurs at times without appreciable change elsewhere. The liver vessels undergo sclerotic change and the same problems confront us; from what we know of the functions of this important gland we should expect to find sclerosis more commonly present in it than the kidneys, and yet I believe that this is not the case. Sclerotic change may occur in any portion of the arterial tree, and atheromatous patches are frequent in the larger trunks after middle life, the coronary arteries being especially subject to sclerotic degeneration. Coronary sclerosis alone has frequently been encountered.

It may be maintained that all these organs are more or less constantly active and more liable to degenerative change. It is not so clear, however, why the arteries of the extremities are prone to undergo sclerotic change, for their activity is not so great or so constant as in the organs mentioned. Yet sclerosis occurring in the arteries of the extremities and the head is common, and it is in the former that we make such estimates as we may of the degree of the sclerosis and the amount of pressure existing within the arteries.

* Read in the Section on Practice of Medicine of the American Medical Association, at the Sixty-Third Annual Session, held at Atlantic City, June, 1912.

ETIOLOGY

Admitting, then, that it is impossible at this time adequately to explain localized sclerosis, it is nevertheless profitable to inquire into its general causation, its method of manifestation and its course and management. Age is a certain producer of sclerosis, but a fixed span of years cannot be given. After middle life a certain degree of sclerosis is apt to be found in the majority of persons. Though the expression be trite, nothing is truer than the axiom that "a man is as old as his arteries," some men of 40 years being in reality 60, 70 or 80. The life of an engine is dependent on the metal which goes into its construction and the care that is taken of it; the same is true of the more perfect physical mechanism; and Osler has aptly observed that a man often takes less care of his body than he does of his motor. It is undeniably true of the body as of the machine that it is better to wear out than to rust out, but in purposive design neither extreme is desirable.

Aside from age, the unailing producer of sclerosis, we have other causes operating; chief among these is the high pressure at which the human machine is run in our modern civilization. The average man or woman lives in an atmosphere of work and worry, taking insufficient thought of either physical or mental needs. One could follow up the myriad avenues of life indefinitely, but they all lead through ways more or less unwholesome and end in the one meeting-point, premature decay. Man presents a pitiful spectacle racing at top speed to gain wealth or position, staggering under the delusion that old age will bring rest and enjoyment, and becoming enlightened, alas, too late, to find that his rest is disturbed by discomfort and that enjoyment is a memory only.

In a practice of more than a score of years I have seen many instances of men who have slaved through life and gained their golden fleece, and who, if their bodies were not broken by ill health, found themselves so enslaved by the habit of work that rest was positive punishment. Such men are like the old blind treadmill horse which, having earned its freedom, when turned out to pasture each morning as the sun rose, walked in a circle till the night.

Next to the stress and strain of modern life, and undoubtedly augmented by it, come the acute infections and intoxications as causative agents. These act as irritants to the cardiovascular apparatus, impairing the vitality of the cells forming their structure and inducing sclerotic change, which makes the vessels less elastic and interferes with the normal metabolism of the organism.

Syphilis is a common producer of sclerosis, and I believe that, as our knowledge increases, it will be proved to be a much more frequent cause than is realized by our profession to-day.

Next come poisons, under which may be included lead and other mineral poisons, alcohol, tobacco, tea, coffee, etc. Billings has shown the causative rôle played by lead, and others believe that mercury when taken in large quantity and for a long time produces it. Alcohol may be classed as an indirect causative agent, in that when used to excess it produces digestive disturbances, which in turn result in auto-intoxications, which are admittedly a prolific cause of the disease.

It would be amiss if in this connection I did not cite Cubot's scholarly paper on alcohol as a producer of sclerosis: in a long series of cases he found sclerosis much less frequent among a series of hard and chronic drinkers than among a similar series of professional men. It should be borne in mind, however, that his work does not

prove that alcohol does not indirectly produce sclerosis; it does prove that sclerosis is less frequent in alcoholics than in those who live constantly under high tension. In this connection Osler calls attention to the fact that not infrequently persons who are abstemious in the use of alcohol are "exceedingly careless and indulgent in the matter of eating." Other authorities are equally suggestive on this score. It is often the case that men who are innocent of one excess go to another. Again, sclerosis is found in those whose occupations demand great and continued physical strain, and in young men who keep up high and long-continued tension in athletic sports.

SYMPTOMS

The effect of arteriosclerosis, by whatever cause it may have been produced, is the disturbance of function; the parts involved become less active and atrophy more or less gradually, according to the speed of the circulation. Sclerosis may involve only the smaller arteries, in which the symptoms will vary with the part or parts affected. The first symptom may be sudden death from the rupture of a cerebral vessel or thrombosis of a coronary artery.

In the kidneys the sclerotic change affects not only the vessels but the interstitial tissue and parenchyma of these organs as well, and is evidenced clinically by changes in the urine. A common form of nephritis following sclerosis is the contracted kidney, in which the blood-pressure is usually high; the apex-beat of the heart is displaced downward and to the left; there is a "thudding first sound" and an accentuation of the second, and more or less nervous and digestive disturbance, with the appearance of albumin and casts in the urine.

Sclerosis occurs in the splanchnic vessels, manifesting itself during life by irregular and otherwise unaccountable attacks of abdominal pain. Advanced sclerosis of terminal larger vessels may result in necrosis or gangrene; and sclerosis of smaller ones, in hemorrhagic infarcts. Finally, it must be borne in mind that sclerosis of the vessels of all the parts mentioned may occur without "any change in the palpable arteries." Since many of the symptoms of the early stages of the disease are indefinite, obscure and not pathognomonic, it is plain that its early diagnosis is at all times difficult and frequently impossible.

It is this phase which makes the critical study of arteriosclerosis all the more imperative, for it is only in its beginnings, whatever be the etiology, that any hope of cure can be promised. After sclerosis is well established a return to the normal should not be expected. It is of the utmost importance, therefore, that the general practitioner be on the alert to discover the earliest recognizable symptoms. These are left ventricular hypertrophy, digestive disturbances, more or less irritability of temper or other nervous phenomena, and the presence of albumin and usually casts in the urine, although cases may occur with any of these absent.

The blood-pressure is estimated through the medium of the touch in palpable arteries and by the employment of an instrument of more or less precision, the sphygmomanometer. This instrument should be more generally employed in all physical examinations, as its evidence is of no small value; it is, however, subject to considerable error and should not supplant the touch of the skilled examining finger. If time permitted, it would be profitable to go into these defects in detail. Hoover, in an excellent contribution to this subject, makes the following observations:

Whether there were really different endovascular pressures in the several arteries or different degrees of hypertonus in the arterial walls which gave the results, the deduction remains the same, namely, that the blood-pressure apparatus applied to the accessible arteries does not give infallible information as to the aortic blood-pressure; as a consequence it is not safe to rely on routine measurements from the area as a guide to the employment of the vasodilators in treating patients with a high arterial pressure. I do not wish to discourage the blood-pressure apparatus. It is a very valuable instrument, but its chief value consists in teaching us how to palpate the arterial pulse. No instrument can take the place of the finger, nor can any instrument teach us so much about the pulse as can be learned by palpation.

Daland has shown that the mere application of the pneumatic cuff in certain neurotic subjects often increases the blood-pressure as much as 30 or 40 mm.

PREVENTION AND TREATMENT

The widest field for good lies in the prevention rather than in the cure of arteriosclerosis. All causes which we know operate to produce the condition should be curtailed or eliminated as far as possible. There are two parties in this problem, the physician as the educator, and the individual who is willing as far as may be to profit by the advice. The individual should as closely as he can approximate the normal life; and excesses of all kinds, mental as well as physical, should be avoided. The pace that kills and the strenuous life should be shunned. Food should be taken in sufficient quantity, and of a quality calculated to maintain the body in a state of physiologic equilibrium; beyond this it is unwise to go constantly or at frequent intervals. Alcohol, coffee, tea and tobacco should be used moderately, if at all.

The exciting cause must be treated in sufferers from toxic agents in the blood, such as lead-poisoning, diabetes mellitus, syphilis, tuberculosis, gout, rheumatism and the acute infections. It goes without saying that the age and general condition of the patient should be considered. What would be moderate in the young would be immoderate in old age; we should not put new wine into old bottles.

The treatment of arteriosclerosis may conveniently be divided into hygienic, dietetic and medicinal.

Everyone requires a certain amount of exercise; the kind and amount should be adjusted to meet the demands of the individual. When active exercise for one reason or another cannot be taken, much good will come from massage properly applied. It quickens the circulation and increases the elimination of waste products. In dilating the vessels of the skin, it to that extent diminishes the work of the heart.

Baths properly administered are of distinct value, including hot baths, hot packs, vapor-baths, tepid baths and carbon dioxid or the so-called Nauheim baths. Cold baths are rarely indicated. All have for their object the increase of blood through the skin, and also excretion by the skin, thereby diminishing the work of the heart and of the kidneys.

The physical and mental activities of the patient must be carefully directed. In some cases curtailment alone is sufficient; in others total abstinence from all work, physical and mental, is imperative. It is frequently surprising to see the good that immediately follows after the patient has made a complete surrender in this direction. Many an old boiler will last for years when run under diminished pressure.

Time will not admit of going into the matter of diet in detail; suffice it to say that the individual capacity will have to be studied in each case. It is a fact that the

majority of persons eat too much, and nearly all too rapidly. In mild cases cutting down the diet may be sufficient, eliminating such articles as are apt to overburden the alimentary tract and the kidneys. In general, meat should be indulged in sparingly. The sugars may be borne well by some, but are apt to lead to gastrointestinal fermentation. Milk, variously modified, eggs, fresh vegetables and fruit are more or less staple articles, as is bread, and may be taken by most patients. In advanced cases a rigidly restricted diet is at all times imperative. In all cases the amount of alcohol, tobacco, tea and coffee allowed should be moderate, and doubtless the best rule in many cases is "not at all." Since we have seen, however, that a large degree of sclerosis is present in most persons after middle life, it is not wise or expedient to be extreme in these matters, until our knowledge becomes more exact.

Individual instances often shatter one's theories. I have seen men, in apparently robust health and advanced in age, who had throughout their adult lives been hard drinkers, excessive users of tobacco, or immoderate tea and coffee habitués; and yet sclerosis in them was in abeyance. The total abolition of these creature comforts is often unwise and their regulation must be left to the wisdom of the physician.

As to the use of mercury in syphilis, I should be willing to run the risk of producing arteriosclerosis by the use of mercury rather than to run the surer risk of having the sclerosis develop from the effects of syphilitic infection. In all luetic cases the iodids are of value; and in many cases which are not luetic their rôle is distinctly good. Their employment may be more or less empirical, but in a large number of cases I believe they tend to hold down tension and discourage sclerotic change.

Nitroglycerin acts quickly and may be employed for a long time without harm, its chief drawback being its evanescence. It rapidly causes dilatation of the peripheral vessels and if pushed sufficiently far will cause flushing of the face and headache. It not only relieves the overburdened heart, but increases the activity of the kidneys. It may be administered in appropriate cases in doses of from 1/50 to 1/100 grain every two or three hours for long periods without injurious effect. Nitrite of amyl pearls may be broken and inhaled in urgent cases, but are not a substitute for nitroglycerin. The more lasting effects of the nitrites make them preferable; of these sodium nitrite in 1-grain to 3-grain doses repeated three times a day or every four hours, may be given.

ABSTRACT OF DISCUSSION

DR. LOUIS FAUGERES BISHOP, New York: I believe that the causes of arteriosclerosis can be divided into two classes, one in which the affection is due to certain poisons like lead and possibly other metals, to syphilis and other infectious diseases, and a larger class, in which the arteriosclerosis is compensatory. Ninety out of 100 cases of high blood-pressure are due to auto-intoxication or to the circulation in the blood of amino-acids which are produced by the breaking up of proteids in the intestinal canal and a failure on the part of the liver to exclude these toxic substances from the circulation. I think that the high blood-pressure is caused by an attempt on the part of Nature to force out of the system through the kidneys the products of protein decomposition, and that the heart is damaged as well as the kidneys. My observations are founded on the examination of about 50,000 urinary analyses among the well-cared-for classes in New York City. A study of these well-cared-for classes gives a better picture of the causes of arteriosclerosis than what we can get in hospital cases. Examination of patients on a

restricted diet are misleading. Observation of the urine, covering a series of years, shows first an excess of indican; later one finds as well a faint trace of albumin and casts. As the case goes on the patient develops neurasthenic symptoms and then arteriosclerosis. I have followed many of these cases in private practice and I have come to certain definite conclusions. Arteriosclerosis is a very common affection among physicians. I have on my records sixty-five physicians with arteriosclerosis and in nearly every instance there was indican in the urine, a high blood-pressure, albumin and casts, and other features pointing to amino-acid poisoning.

The important point regarding these cases of arteriosclerosis is, of course, the treatment. Are the amino-acids coming from meat decomposition always the cause of the trouble? I do not believe that this is the cause in every case. Among the other possible causes may be overindulgence in eggs or fish.

Another very important thing in the treatment of these cases, I believe, is the proper administration of castor oil; these patients on beginning treatment are given 1 ounce of the oil for three days, then the same amount once a week and then the same amount once a month; the results obtained are often surprising. A third important measure in the treatment of these cases is physical labor. I have seen patients with arteriosclerosis who have had slight strokes of paralysis, with a blood-pressure of 230, and who, under the regimen mentioned in regard to the proteid diet and with the employment of physical exercise, have gone on for years.

Dr. G. W. McCaskey, Fort Wayne, Ind.: There are three factors to be considered in every case of so-called arteriosclerosis: First, the sclerotic changes in the blood-vessels proper; second, the cardiac sclerosis; and third, the factor which is clinically most important, because most amenable to treatment, the arterial spasm. So far as the etiology of arteriosclerosis is concerned, I think that two distinct causes should be recognized, the first being toxic, and the second mechanical and due to physical overstrain from various causes. The toxic causes may be derived from the intestines or may be the result of metabolic disturbances of one source or another, perhaps various products of the amino-acids. Severe physical strain or the overexcitement of the cardiovascular apparatus resulting from emotional excitement or mental stress undoubtedly produces arteriosclerosis.

It is important that we should have clear-cut notions of the therapeutic indications in these cases. We ought, in other words, to distinguish clearly between the organic changes of the blood-vessels and heart which cannot be removed, and the arterial spasms which are very largely removable. Rest in bed with thorough elimination, carefully regulated diet, from which the purin bodies should be largely if not completely removed, and the use of vasodilators constitute the proper treatment.

As vasodilators I commonly use the spirits of glonoin, and begin in hospital cases with 1 drop every hour, increasing every day or every alternating day to 2 drops per hour and pushing the dose in this way up to the limit of tolerance, or until satisfactory results have been obtained.

Dr. Henry O. Marcy, Boston: Arteriosclerosis is largely the result of the undue pressure that is brought to bear on the human machine which we should all learn how to care for and protect. To me the wonder is that the modern business man in our large cities does not break down earlier. In the whirl of business care, his every activity is in high tension. He takes little time for meals, pampers his appetite with every delicacy, uses coffee and alcoholic stimulants freely, smokes excessively, exercises little, sleeps badly and wakes to repeat the round. Over such a man at home the physician has little control. Send him to Carlsbad and note the change. Diet simple, largely milk, the waters in sufficient quantity to act on the excretory organs; no alcohol, no tobacco, nothing except as ordered by the physician. The psychotherapeutic effect of his hours of hot mud-baths and repose four or five thousand miles from his business cares renovate him and make him enthusiastic over his cure, but he returns only to plunge again into the same round of busi-

ness dissipation which leads to physical ruin. Why call the results by such meaningless names as "arteriosclerosis"? The duty of the physician is to sound the danger-signal before it is too late; to teach hygienic living, abstemious habits, the simple life. We of the profession know that under wise guidance the ills of life would be reduced one-half, the production of the race doubled and longevity increased at least one-third.

Dr. James M. Anders, Philadelphia: Dr. Coleman spoke particularly of the localized variety of arteriosclerosis and of the great difficulty in finding its precise cause. I think that the majority of authorities are agreed that it is due most commonly to syphilis, and this also accords with my own personal experience and observation. To my mind the treatment or management of arteriosclerosis often savors of a comedy of errors. If the modern conception of the histologic changes of the condition be correct; if, primarily, the media is the seat of either productive or degenerative lesions, and to compensate these changes, the intima becomes greatly thickened, then surely, two things must follow: First, the lumen of the vessel must be narrowed, and second, the tension in the vessels must be raised to a corresponding degree. One is tempted to ask, Why attempt to interfere with this compensatory mechanism of Nature? I believe that the trend of the profession has been and still is in the direction of interfering too much rather than too little with this wise conservative process. There is, however, a class of cases made up of instances of incipient arteriosclerosis, in which the blood-pressure is considerably elevated, associated with a marked vasoconstriction due largely to nervous causes. Here absolute bodily and mental rest, arterial relaxants and liberal feeding will often effect an apparent cure, provided that the causes, both near and remote, can be successfully combated. It should not be forgotten that the principal aim in all cases must be to ascertain and then to combat causes.

Dr. DeLancey Rochester, Buffalo, N. Y.: If anything was brought out clearly in the previous discussion of this subject in this Section several years ago, it was that the chief etiologic factors of arteriosclerosis are syphilis and hard work. Under the circumstances, to place these people at hard work is decidedly contra-indicated. Any exercise should be undertaken only under careful observation. I agree with what has been said regarding the reduction of the proteids in the diet, especially if there is a decided impairment of renal function. Elimination of waste products is of value in the treatment of this disease. I agree with Dr. Anders that arteriosclerosis, when localized, is largely a condition due to syphilis. Since the use of the Wassermann reaction we have found more evidence of syphilis as an etiologic factor in arteriosclerosis and changes in the circulatory apparatus, especially of heart and aorta, than ever before.

Dr. M. A. Conn, Brooklyn: I wish to cite a case just now under my care. A man, aged 27, presented himself April 2, 1912, with the following history: While walking in the street with a friend twelve days previously he suddenly collapsed and was picked up unconscious and limp and pronounced dead, being refused admittance into the offices of two successive nearby physicians on that ground. The third physician finally allowed the dying man to be stretched on the floor while a hypodermic was being administered and the ambulance telephoned for. After remaining in the hospital one week, the patient was discharged as cured and was permitted to walk home without any warnings or restriction whatever. The man, a plumber by trade, big and well nourished, looked sallow and anxious, with dark rings under his eyes and he breathed rather laboriously. A careful examination revealed the following condition: pulse high-tensioned; mitral and aortic systole feeble and muffled; the characteristic blue line about his gums; traces of albumin, some hyaline casts and a slight excess of indican in the urine, and some moist râles at the base of the left lung. On applying my sphygmomanometer, I was not a little amazed to find that no less than a pressure of 280 mm. Hg was required to stop the radial pulse. As a hectic history, was positively denied and no signs or symptoms pointed in that direction, the

conclusion was reached that the case was one of chronic plumbism producing arteriosclerosis and interstitial nephritis, the feeble myocardium giving rise to the pulmonary edema and to the frothy, pink expectoration which occurred the night prior to the patient's sudden collapse and continued for two days thereafter. He was informed of the extreme gravity of his condition, was put to bed on a strictly milk diet, and, after a thorough elimination by bowels, kidneys and skin, was given potassium iodid to remove the lead from the system; doses of nitroglycerin were gradually increased. Seeing, however, no improvement, subjective or objective, even after doses of 1/20 grain of nitroglycerin three times a day, I discontinued its use and on the suggestion of Dr. G. R. Butler, who saw the case with me, began the use of 1/2-grain doses of erythrol tetranitrate three times a day with the simultaneous administration of 5 drops of the tincture of strophanthus three and four times daily and also short steam baths, with patient in bed, once every one or two days for a week or so. The patient is temporarily improving, is allowed some outdoor exercise and a little more liberal diet, with little fluid, however. The blood-pressure is still above 240 mm.; some albumin and casts are still to be found in the urine, but there were no attacks of angina pectoris recently, no dyspnea or polyuria and the whole of the Stokes-Adams' syndrome is absent.

DR. MARVIN L. GRAVES, Galveston, Tex.: Until we know more regarding the etiologic factors concerned in the production of arteriosclerosis we can do but little in combating it. A great deal of consideration should be given to the character of the life of the individual in ascertaining its relation to high blood-pressure or arteriosclerosis. Dr. Bishop believes that these arteriosclerotic patients with high blood-pressure should be urged to take exercise, but I think a discriminating judgment is absolutely necessary in treating these patients. For instance, arteriosclerosis occurring in a laboring man who has been at active physical exercises all his life should be treated differently from that in a man who has been leading a sedentary life. Under many conditions I do not think it would be advisable to prescribe large amounts of physical exercise for these patients who are afflicted with arteriosclerosis. To order a man who leads a sedentary life, and who eats too much of all kinds of foods, not only the proteid foods but all kinds, to take physical exercise indiscriminately would tend only to work harm. Such men should be kept as much as possible in the open air with properly graduated exercise; the total amount of food they were in the habit of consuming should be cut down, especially the proteids. In doing this, many of these individuals would be relieved. I think that a sharp distinction should be drawn between toxic conditions which produce high blood-pressures and changes in the arteries which cause arteriosclerosis with its increased blood-pressure.

Too much stress is laid on the reduction of the blood-pressure. I have seen, in patients of 45 and 60 years of age, an unduly high blood-pressure—one of 150, one of 200 or even one of 225 mm. Hg—which has been reduced by the enforcement of a rigid diet and the use of nitroglycerin or sodium nitrite, as well as an active purgation; under such treatment one also sees patients who are made worse and not better. I do not believe that we should always try to reduce the blood-pressure to what we consider to be the normal. It is impossible, in my opinion, to reduce to the normal, with safety to the patient, a high blood-pressure which has been maintained for a number of years, especially with arteriosclerosis.

DR. TOM A. WILLIAMS, Washington, D. C.: Arteriosclerosis is a condition which can be prevented but which, when it has occurred, cannot be cured. Many of the early symptoms are cerebral. The guide in the treatment of individuals afflicted with arteriosclerosis should not be what could be accomplished in lowering the blood-pressure *per se*, but the study of the state and condition of the individual which determined his optimum pressure. Naturally, with a very high optimum pressure, time of life will be short. Of the means of treatment at our disposal, the most important relates to the diet.

I believe in a diet which is similar to the Chittenden standard; that is, that 50 gm. protein per day is ample and should not be exceeded in these cases. It is wonderful how well these patients will do when the proteins are cut down, and more especially the proteins that are taken at the evening meal before the lowered oxidation of sleep. Beyond abundance of succulent fruits and vegetables, I do not believe that purgatives should be resorted to as eliminants. They interfere with osmosis and muscle tone of the bowels and hence favor the absorption of substances which make for increased blood-pressure. Both as revulsants and eliminants I favor the use of hot baths and massage, and no expensive apparatus is needed. The efficacy of active exercise in these cases is exaggerated, but not contra-indicated in those cases of arteriosclerosis which might be described as being in the presclerotic stage. It is a very important thing for the patients themselves to appreciate their condition. What is impending unless forestalled should be impressed on them by the physician in order that they may readjust their lives and do what is best for their greatest efficiency in the long run, even though it might result in some immediate loss of comfort or money. These principles of practices have been of great value in my hands, and have made unnecessary recourse to expensive measures like d'Arsonval currents, water cures and the like.

THE MENACE TO THE YOUNG CHILD OF THE COMMON INFECTIOUS COLD*

THOMAS S. SOUTHWORTH, M.D.

NEW YORK

Many of my colleagues are doubtless convinced of the truth contained in my title, and would accept the text without the discourse which follows, for it is scarcely possible that my individual experience has been exceptional or that it has not been duplicated by scores of others. Nevertheless, it has seemed to me proper to put my views in concrete form in the hope that there may result a wider realization and discussion of this problem in all its bearings.

The observations of recent years have taught me to respect the common infectious cold as a far from trivial affliction and one capable of inflicting serious injuries on its victims. Ever since the initial intensity of the influenza pandemic of 1889-90 waned, we have experienced recurring waves of infectious catarrhal colds which have been spoken of generically as influenza, or grip, or simply as colds. In the absence of laboratory confirmation of the exact bacterial cause, the classification of such endemic or epidemic affections has not been possible, and the resulting uncertainty has found expression in the loose terminology adopted alike by the medical profession and by the laity.

Mention should properly be made in this connection of the careful work in this country by Wollstein, but since her researches demonstrated the presence of the Pfeiffer bacillus as a causative or complicating factor in only a certain proportion of the inflammations of the respiratory tract among the infants studied, they threw but little light on the etiology of the cases under discussion, save by indicating that such cases may be due to several diverse forms of microorganisms. Nevertheless, since it is well-nigh impossible under the ordinary conditions of general practice to differentiate true influenza from its imitators, and since clinical observation has demonstrated the unmistakable infectiousness of such colds, we are forced to consider them as a whole for the very practical purpose of inquiring whether much

* Read in the Section on Diseases of Children of the American Medical Association, at the Sixty-Third Annual Session, held at Atlantic City, June, 1912.