JOURNAL OF THE ROYAL MICROSCOPICAL SOCIETY.

FEBRUARY 1883.

TRANSACTIONS OF THE SOCIETY.

I.—*Observations on the Anatomy of the Oribatidæ.*

By A. D. MICHAEL, F.L.S., F.R.M.S.

(Read 10th January, 1883.)

PLATES I. AND II.

IN preparing the work upon the British *Oribatidæ*, which I am now writing for the Ray Society, it became desirable to deal, as far as I was able, with the leading features of the anatomy. It was

EXPLANATION OF PLATES I. AND II.

PLATE I.

- FIG. 1.—Alimentary canal of *Nothrus theleproctus*. × 50. *a*, œsophagus; *b*, ventriculus; *d d*, cæca; *e*, small intestine; *f*, colon; *g*, rectum; *i i*, anal plates; *j j*, preventricular glands.
- „ 2.—Alimentary canal of *Hoplophora magna*. × 50. Same lettering.
- „ 3.—Alimentary canal of *Damæus geniculatus*. × 35. Same lettering, and *c*, ingluvies; *h*, labium; *l*, maxillæ; *m*, palpi.
- „ 4.—Part of the alimentary canal of *Leiosoma palmicinctum*. Same lettering, and *n n*, supposed ducts from preventricular glands.
- „ 5.—Ending of trachea in the acetabulum of the third leg of *Damæus geniculatus*. × 100. *a*, acetabulum, removed from the ventral plate; *b*, coxa; *c*, femur; *d*, trachea, ending in a small air-sac in the acetabulum.
- „ 6.—Two tracheæ of *Damæus geniculatus*, uniting into a single trunk before their attachment to the acetabulum of the second leg.
- „ 7.—Air-sac near mouth of *Nothrus theleproctus*. × 350.
- „ 8.—A portion of the left side of *Leiosoma palmicinctum*, the dorsal shield and all the internal organs, except the super-coxal gland, having been removed. *a*, wall of rostrum; *b*, first leg; *c*, second leg; *d*, ventral plate; *e*, super-coxal gland.
- „ 9.—The super-coxal gland of *Leiosoma palmicinctum*. × 350. *a*, gland; *b*, globular body (vesicle?).
- „ 10.—Mandible of *Oribata globula* (ordinary type in the family). × 180.
- „ 11.—Mandible of *Pelops phæonotus*. × 600. Type of the genus *Pelops*.
- „ 12.—Mandible of *Serrarius microcephalus*. × 250.
- „ 13.—Serrated end of same mandible. × 500.
- „ 14.—Two teeth of same mandible, highly magnified.

[PLATE II.

Ser. 2.—VOL. III.

B

a subject upon which there were practically only three existing authorities. The first of these is Dujardin,* a keen-eyed observer, who saw a good deal, but unfortunately, as far as the *Oribatidæ* are concerned, drew conclusions from what he saw which have formed a stumbling-block for most later writers. The second is Nicolet,† and this may be treated as the only substantial work upon the subject. The third is Claparède,‡ whose work is excellent as far as it goes, but it only deals with one very exceptional species of this family, and that with a view to the development rather than the anatomy.

When I first commenced, my idea was simply to verify Nicolet's work before repeating his account in my own book; and to ascertain that it was correct, not only for the one or two species he had described, but also for others. As I advanced, however, I found so much variation in different forms, and so many points upon which I was not able to coincide with Nicolet's descriptions, that I was led to devote the greater part of my leisure during the summer and autumn of 1882 to the investigation: it has been

PLATE II.

- FIG. 1.—Male reproductive system of *Nothrus theleproctus*. × 50. *a, a*, lobes of the testis; *bb*, flat portion, possibly vesicula seminalis; *cc*, vasa deferentia; *d*, ductus ejaculatorius, with penis, penial skeleton, and sclerites, &c.
- „ 2.—Penis and penial skeleton and sclerites of same species. × 600. *a*, penis.
- „ 3.—Penis of same species, side view. × 1500.
- „ 4.—Semen of *Damæus geniculatus*. × 1800.
- „ 5.—Female reproductive system of *Oribata lapidaria*, taken when the creature has just emerged from the nymphal stage. × 40. *a*, ovary; *bb*, oviducts; *c*, vagina; *d*, ovipositor; *e*, genital plates; *f*, anal plates.
- „ 6.—Female reproductive system of *Cepheus tegeocranus*, mature, showing the oviducts full of eggs. × 60. Same lettering.
- „ 7.—Female reproductive system of *Damæus geniculatus*, mature, ovary nearly exhausted. × 50. Same lettering: and *g*, globular expansion of oviduct; *hh*, copulative suckers (so called); *ii*, tendons to genital plates; *jj*, muscles attached to same tendons.
- „ 8.—Portion of the ovary and portion of oviduct of same species, up to and including the globular expansion. × 120. Same lettering.
- „ 9.—End of trachea in acetabulum of leg of *Leiosoma palmicinctum*. × 180. *a*, edge of acetabulum; *b*, coxa; *c*, trachea; *d*, muscles of leg.
- „ 10.—Anal sac of *Nothrus theleproctus*. × 100. *aa*, sac; *bb*, genital plates; *cc*, anal plates.

* "Premier Mémoire sur les Acariens," Ann. des Sci. Nat., 3rd ser. iii. p. 5. Journal de l'Institut, 1842, p. 316.

† "Histoire Naturelle des Acariens, qui se trouvent aux Environs de Paris," Archives du Muséum, vii. (1855). Paris.

‡ "Studien an Acariden," Zeitschr. f. Wiss. Zool., xviii. (1868) p. 446.

suggested to me that the results may be of interest to one of the biological societies, I have therefore described some of them in the present paper, instead of waiting to include them for the first time in the future treatise. I do not propose, in these pages, to give any exhaustive account even of such portions of the anatomy as I am acquainted with, that would be hardly fitted for this Journal, and would certainly be too lengthy, but I shall confine myself to such portions as seem to me to be undescribed, or to vary substantially from Nicolet's account, and to such other parts as are necessary to the comprehension of the novelties. It is only fair to Nicolet to point out, that, in his beautiful work above quoted, he states, that, in consequence of the minute size, and the hard, opaque, chitinous, exo-skeleton of these creatures, he found the internal anatomy extremely difficult. Claparède gives a similar reason for scarcely touching upon the internal anatomy.

There cannot be any doubt that Nicolet and Claparède were right as to the difficulty. The largest specimens are under a millimetre in extreme length, and many of the species which I have dissected are not above half that size; they are possessed of a chitinized cuticle which is nearly as hard and as brittle as glass, and which is usually quite opaque at all times except immediately after the ecdysis. I first tried observations at that period, but I soon found that I could not obtain much information this way, for, although, now that I am acquainted with the organs, I can frequently recognize many of them through the dorsal plate during its short period of transparency, yet the view is too imperfect, and the organs too much hidden by one another, for original inquiry. I endeavoured to stain them whole, but entirely failed in getting any stain to penetrate the chitin of the exo-skeleton, or even to run in at joints, stigmata, &c., although I tried the air-pump, hoping it might assist. Finally, after a not very satisfactory attempt at section cutting, I determined to face the difficulty of the small size, and to rely entirely upon actual dissection—these notes are the result.

I soon found, that, in order to produce a successful dissection, the creature must be in good, healthy, condition; it must not have been kept long in confinement, and it must be dissected immediately after death.

The dissections have, in each instance, been frequently repeated, i. e. upon a considerable number of specimens of each species, as, in consequence of the difficulties, the naturalist would be too liable to error in judging from one or two instances.

All the figures are made from actual dissections, they are not calculated, or put together, from the result of consecutive sections. The preparations from which the drawings are made, are, in almost all instances, now in my possession, stained with logwood, and

mounted for the Microscope, so that their appearance can easily be verified by any one interested in the subject, although of course different opinions may be entertained as to the correctness of the conclusions which I draw from them.

The dissections have chiefly been made under a Stephenson binocular, with powers varying from an inch to $\frac{1}{2}$ in. and low eye-pieces; they have, however, in cases requiring it, been verified and examined under amplifications running up to over 1000 diameters.

I think I may say that the tendency of my mind has been a desire to confirm previous writers, and not to upset their statements. Where I am not able to agree with them, it is that I have failed to obtain the same result, and that the actual organs before me do not seem to me to agree with the accounts.

The principal differences which I find in the internal organs are in the respiratory system, and the reproductive system. I find differences of lesser importance in the alimentary canal; and I believe that I have traced some secreting organs not before recorded; these are the points upon which I intend to touch; and, finally, I shall mention two matters in connection with the exoskeleton which seem to me to be worthy of notice.

It is necessary to describe the alimentary canal in order that the varieties, &c., which it displays may be understood, although Nicolet figured the greater part of it correctly in one species.

The Alimentary Canal.

In the *Oribatidæ* the alimentary canal is somewhat short and simple, that is to say, there is a comparative absence of convolutions in the hind-gut, and the very numerous cæcal prolongations of the mid-gut found in many allied groups, as the *Aranea*, *Picnogonida*, &c., are reduced to two, which, however, are usually of great size and importance. The proportions of the parts of the canal, and the size and arrangement of the cæca, vary greatly in different species, but, as far as my experience goes, the divisions of which it is composed are always the same. I say "as far as my experience goes," because I have not dissected anything like the whole of the species.

The canal is composed of the cesophagus, the ventriculus, a short small intestine, the colon, and the rectum, terminating in the anus and anal plates; of these the two first are almost horizontal, and are placed in a straight median line near the dorsal surface; the small intestine turns downward and slightly forward, and the colon and rectum are more or less perpendicular, and lie beneath the ventriculus, or almost so; the result of this is, that, when the dorsal shield is removed, the cesophagus and the ventriculus, with its

cæca, are usually the only parts of the canal seen. The canal is, of course, firmly attached round the cavity of the mouth, and at the anus, but in other parts it seems to float freely in the general body-cavity, with only very slight attachments, if any. If the œsophagus be cut away from round the mouth, and the rectum from round the anus, a hair may easily be passed under the canal, and its whole length be drawn out on the hair without further injury.

Taking *Nothrus theleproctus* as a convenient type :—

The œsophagus is a long, thin, almost straight, tract of the canal, extending from the mouth to the ventriculus, and having its lowest point at the former and its highest at the latter. It has thin walls, capable, however, of considerable expansion and contraction. The cavity of the mouth being larger in diameter than the lumen of the œsophagus, the latter necessarily widens somewhat as it approaches the former, and the widened portion might, not unfairly, be termed the pharynx; posterior to this the œsophagus continues of almost even circumference for the greater part of its course through the cephalothorax; near its posterior extremity it widens, in some species this enlargement is considerable, and then forms an ingluvies or crop, the jabot of Nicolet; this writer correctly gives the ingluvies as very much developed in *Damæus geniculatus*, plate I. fig. 3, c; in *Damæus clavipes* it is even more developed and is almost as large as the ventriculus. I cannot say that I have ever seen it so large in other genera, but it is no doubt quite distinguishable in many others. Nicolet also, in the same drawing, depicts the œsophagus as being constricted at short intervals by circular bands of muscle, so that it presents a moniliform appearance. I have not been able to detect this moniliform effect in any other species which I have dissected, but the bands of muscle are usual. In the ingluvies of *Damæus geniculatus* these circular bands of muscle are beautifully seen after the preparation has been stained with logwood, and indicate that it is an expansion of the œsophagus, not a separate stomach. Nicolet further states that an air-bubble is invariably found floating in this ingluvies; my own observations do not quite confirm this; a large air-bubble is certainly to be seen very frequently floating in the canal, but it appears to me that it is not by any means invariably present, and that, when present, it is more frequently in the ventriculus than in the ingluvies; it is doubtless due to the creature's living chiefly upon liquid materials derived from vegetable substances, and absorbed by a sucking process, which accounts for the presence of liquid in considerable quantities in the fore- and mid-guts, and for those parts not being quite filled with it.

The œsophagus continues the whole length of the cephalothorax, and passes a very short distance into the abdomen; it is sharply

constricted by a circular muscular arrangement at the point where it enters the ventriculus, forming a very perfect valve, which prevents any food material from returning from the ventriculus to the œsophagus; this may often be seen very well by the air-bubble, which, carried by the liquid in which it is floating, may be observed to pass down from the œsophagus to the ventriculus, but it is always stopped at the valve whenever it has a tendency to return. The œsophagus enters the ventriculus about the centre of the anterior end, which possibly it may not be proper to call cardiac in a creature which is not known to possess any heart.

The *ventriculus* is the largest and most important portion of the canal; it is a wide sac, occupying from half to about two-thirds of the length of the abdomen, near the dorsal surface (plate I. figs. 1, 2, 3, 4, b). In *Nothrus theleproctus* the sac is of a shape approaching pyriform, the narrower end being the anterior one, this however is rounded, although smaller than the hinder part; the posterior end has a tendency to a blunt median point. The walls of this viscus are thick and muscular, more so than any other part of the canal except the rectum, and it is here that digestion doubtless chiefly takes place, a food mass may, most frequently, be seen occupying the pyloric portion. From the widest part of the ventriculus, i. e. about two-thirds of the length from the anterior end, a large cæcal diverticulum arises on each side, and, in the present species, after standing outward (laterally) so as to form a shoulder, proceeds almost straight backward. These diverticula are longer than the ventriculus itself, often more than half as long again, and nearly half its diameter; they are of almost even size throughout, so that they present a sausage-like appearance; their posterior ends are rounded.

There are probably greater differences between the ventriculus and cæcal appendages in different species and genera of the family than are found in any other part of the canal; this is especially true of the size and form of the cæca. In *Hoplophora magna* they have become very small in proportion to the ventriculus, which is widest anteriorly, and prolonged almost to a point behind. The cæca are globular, and are attached to the ventriculus by short peduncles (plate I. fig. 2, d). I mention this species, as Nicolet figures it without cæca, which is not correct in the specimens which I have dissected. In the case of *Damæus clavipes*, the cæca are not any longer distinctly visible, but are mere enlargements of the outer posterior angles of the ventriculus. *Damæus geniculatus* is an intermediate form in this respect.

Nicolet treats the cæca as being simply diverticula of the stomach, without any special office, and this probably may be the most obvious and natural suggestion; as far, however, as my own judgment goes, I doubt the correctness of the view. In the very

numerous instances which I have examined, to the best of my ability, I have not ever been able to detect food in the cæca, either in specimens dissected out or in those observed in life where the transparency of the chitin enabled me to do so, which was frequently the case; whereas it is rare to find the ventriculus or colon without food contents. In addition to this, the structure of the walls of the cæca, particularly near their distal ends, is different from that of the ventriculus, they being thicker and much more glandular, and the lumen smaller in proportion, often very narrow. These considerations lead me to infer that their function is secretory, and not simply that of an extension of the stomach, nor would there be anything extraordinary in this being so, as a very similar arrangement seems to exist in *Apus*, *Limnadia*, &c., the ends of the cæcal diverticula of the mid-gut being differentiated to form glandular organs, and the same arrangement, carried to a greater extent, prevails among the *Malacostraca*, &c. If these two large diverticula have, as I suppose, the office of secreting fluids necessary for digestion, it would explain to a great extent the apparent absence of the so-called hepatic tubes, an absence observed by Nicolet, who remarks that he was not able to detect any liver. In connection with this subject, however, must be taken the observations which will be found below as to the presence of diffused follicles, supposed by some writers to have a hepatic function, over the surface of the canal.

The *small intestine* (plate I. figs. 1, 2, 3, 4, *e*). This portion, forming the commencement of the hind-gut, may possibly be considered as simply a portion of the colon, from which it is not divided by any valve; but as, although short, it is always present, it may be convenient to treat of it under a separate heading.

The entrance to the small intestine from the ventriculus is always closed by a very efficient valve, which may occasionally be seen to open in order to give passage to the balls of digested or partially digested food. The small intestine proceeds from the ventriculus at a point lying to the left of the median line and near the pyloric end, but not actually at the end; the entrance is situated either at the edge of, or slightly on the under surface of the stomach, but its position varies a little in different species.

The *colon* (plate I. figs. 1, 2, 3, *f*), in all species in which I have examined it, is an elliptical enlargement of the hind-gut, very considerably smaller than the ventriculus, and usually clearly defined, particularly at the posterior extremity, where it is provided with a constriction or valve which usually completely closes the entrance to the rectum; it ordinarily turns downward, or slightly forward, so as to be brought more or less under the ventriculus. In a dorsal view when the notogastral shield has been removed without disturbing the position of the canal, only the anterior end

of the colon is seen. In my figures of the alimentary canal the colon and rectum are extended into a horizontal position so as to be seen.

Nicolet in his drawing of the alimentary canal of *Hoplophora magna* omits this portion of the canal altogether, but he gives one enlargement of the hind-gut almost as large as the ventriculus; this enlargement I presume he intends for the rectum. I can only say that I have not found such an arrangement in any of the specimens of this or any other species which I have dissected; indeed Nicolet's canal differs so materially from what I have met with, that I should think that we were dealing with different creatures, were it not that the species is an extremely well marked one. The walls of the colon are far less thick and less muscular than those of the ventriculus or rectum.

The rectum (plate I. figs. 1, 2, 3, *g*) is usually pyriform, sometimes very small where it arises from the colon, as in *Hoplophora magna*, sometimes larger, as in *Nothrus theleproctus*, but always distinctly divided from the colon by a sharp constriction; in some species, as *Oribata punctata*, it continues small for some little distance, gradually enlarging until near the anus, at, or within a short distance of which, it is closed by powerful sphincter muscles; the actual anal end of the rectum is attached round the opening on the ventral surface which is defended by the anal plates; a series of longitudinal muscles also arise from the posterior parts of the rectum, and are inserted in the exo-skeleton near the anal plates; they doubtless assist in supporting the rectum and holding it in its place, and probably also assist in defecation.

The rectum itself is muscular, with the bands of muscle arranged circularly; this is very clearly seen in *Nothrus theleproctus*, where, as in the last-named species, the constriction is some little distance from the anal plates; the portion of the canal between the two is comparatively very thin and delicate.

The Accessory Glands.—There are two conspicuous glands, not I believe mentioned by any author, which I propose to call the "pre-ventricular glands"; they frequently show like two black spots through the dorsal surface, when at all transparent, particularly in the nymphs, but they are seen equally well in the adults when the chitin is not too opaque, and they appear to me to be always present. When dissected out they are found not really to be black but to vary in colour from deep yellow to dark brown. These glands are shown in plate I. figs. 1, 2, 3, 4, *j*, and are seated on the ventriculus at its extreme anterior part; one on each side of the cesophagus, just where it enters the ventriculus. The edge of each gland is on the dorsal surface of the ventriculus; so that, from the dorsal aspect, they look globular, and often are so, but more frequently they are somewhat flattened and show a tendency

to a bilobed form, in which case they extend down the side of the ventriculus as far as their size renders necessary; they are composed of large, loosely aggregated cells, and are easily broken up if disturbed in most species. Two similarly placed glands appear to exist in such Vermes as *Prorhynchus fluviatilis*.

It was some considerable time before I could trace the ducts from these glands at all to my own satisfaction, they are usually so very delicate that it is extremely difficult to detect them, and, although I thought that they followed the course which I now suppose to be correct, yet I could not feel any certainty. After trying numerous species I experimented on *Leiosoma palmicinctum*, in which I was pleased to find the ducts apparently more distinct; they seem, as far as I can at present judge, to run in an almost straight course along the surface of the ventriculus, which they appear to enter just above the cæca (this is shown in plate I. fig. 4, n), where they are delineated as separated from the ventriculus by dissection. If this course of the duct be correct, the office of these glands is doubtless the secretion of some fluid useful in digestion. The point cannot yet be considered as decided, as the glands have some attachment to the outer wall of the body. I have since found that they are equally well seen in *Notaspis lucorum*. Probably, if Burmeister's views were to be followed, these glands, from the place where their ducts seem to discharge, should be called pancreatic, but as the Oribatidæ are vegetable feeders the function would not be analogous. The glands and ducts possibly form the homologues of one pair of the anterior cæca of the ventriculus present in so many of the Arachnida.

The dorsal part of the anterior portion of the ventriculus and of the whole of the cæca is usually covered with a thickish layer of brown follicular-looking cells, which sometimes entirely cover the dorsal surface of the ventriculus. I cannot say that I have ever succeeded in detecting any ducts from this mass; they would doubtless be very fine, but I think that they are identical with those which many authors have considered as having a hepatic function. Thus Megnin, in his able treatises on the *Gamasinæ*,* and on the *Sarcoptidæ* of mammals, says that he regards this brown granular substance as being analogous to that which coats the hepatic tubes of insects, and as being in fact the liver. A similar organization in many annelids was pointed out long since by Quatrefages,† who assigned it a similar office. Claparède speaks of the ventriculus, &c., and coating of cells as the "lebermagen" in the *Hydrachnidæ*, and that most careful anatomist, Cronberg,

* "Mémoire sur l'organisation et la distribution zoologique des acarïens de la famille des Gamasides," Journ. Anat. et Phys. (Robin) 1876, p. 315. "Monographie de la tribu des Sarcoptides psorique," Revue et Mag. de Zool., 1877, p. 157.

† "Mémoire sur quelques Planariées marines," Ann. des Sci. Nat., 1845.

expressly states his complete adherence to Claparède's opinion, extending it to *Trombidium*.* On the other hand Leuckart † considered that a similar tissue in the leach was not hepatic, and Ledig in his histology agreed in this with Leuckart, and the more recent investigations of Professor E. Ray Lankester ‡ would seem to point rather to a blood-elaborating function for this "botryoidal" tissue.

I have not been able to find any salivary glands nor any appendages to the hind-gut.

The Reproductive System.

This, next to the alimentary canal, is the largest and most important set of organs in the body; the general arrangement is very similar in the two sexes. When the notogastral shield has been removed, the genitalia may be seen lying at the sides of the canal, and in some instances, in the female, when the oviducts are distended with eggs, they seem to have usurped the place of almost all the other organs, and to have pushed them out of position.

The Male Organs of Generation.—These consist of a large central testis, or more probably one on each side, coalescing in the median line by being imbedded in, and united by, a flat mass, which appears to perform the double office of increasing the quantity of the secretion and acting as a vesicula seminalis; two vasa deferentia, a ductus ejaculatorius, a penis with its accessory organs, and three pairs of copulative suckers.

The *testis*, treating the whole as one organ, is very large, sometimes appearing to half fill the body and force its lobes up to the notogaster; it usually extends the whole width of the body, and forms a saddle-shaped mass, which underlies the ventriculus in the centre, there constituting an almost flat layer of considerable thickness, deeply indented, both anteriorly and posteriorly, as though it were two paired organs which have met and united; no sign of suture or demarcation is however visible there (plate II. fig. 1, *a, b*). In addition to this, each side has a tendency to be bilobed, and the anterior lobe, which is part of the flat mass, in some species, or at some periods, rises much nearer to the dorsal surface than the posterior one; the whole varies somewhat in form in different species, in *Nothrus theleproctus* the anterior lobes are large and rounded, in *Oribata lapidaria* they are smaller and squarer. Along each side of the flat part already described, and partly imbedded in it, but not reaching its anterior edge, runs a raised, rounded portion, which is oval in *Nothrus theleproctus*, but varies a little in different species. I have not ever succeeded in

* "Ueber den Bau von *Trombidium*," Bull. Soc. des Nat. de Moscou, 1879.

† 'Die Menschlichen Parasiten,' vol. i.

‡ Quart. Journ. Micr. Sci., 1880, p. 317.

finding any demarcation on the inner side, where they join, nor in removing one from the other without tearing both to pieces. The substance of both is white, soft, and glandular, but the rounded lobes stain more deeply with logwood, and then exhibit a mottled appearance; these lobes are probably the true testes, the flatter central portion would seem to be more or less hollow, and to serve as a vesicula seminalis, as well, probably, as an organ secreting some fluid to increase the quantity of the spermatie material; at all events the walls are very thick and glandular.

The coalescing of the testes into an unpaired organ is of course common amongst the *Arachnida*, e. g. *Phalangium*; it also reminds one considerably of the arrangement in *Homarus*, and in *Pleuroma* and other free-living *Copepoda*, &c., among the *Crustacea*, and in insects a very close approach is to be found among the *Lepidoptera*, as in *Pontia brassicæ*, &c.; but if the flatter central portion acts also as a vesicula seminalis this would apparently be a departure from the crustacean type. Nicolet draws the testis in *Damæus geniculatus* as being four oval, free bodies, on each side, not imbedded in any flat central portion. I am not able to account for this; certainly, when they are seen through the dorsal surface in transparent specimens, or seen after the notogastral shield has merely been removed, they might be mistaken.

It may of course be suggested that the raised oval portion is the true testis, and that the central portion has some hepatic or other function, and not that of a *vesicula seminalis*; but the absence (as far as I have been able to ascertain) of any ducts except the vasa deferentia, and the fact that they lead out of the flat portion, and the intimate way in which the whole is fused into one mass, would seem to negative such a view.

The *vasa deferentia* are substantial tubes of moderate length, not longer as a rule than the testis itself; there is one on each side, springing from the anterior part of the flat portion, close to the raised oval (plate II. fig. 1, *c*), and passing downward, and slightly forward, until they join, and open into the ductus ejaculatorius; they have a great resemblance to the paired *vasa deferentia* proceeding from the unpaired testis in the above-quoted case of *Pontia brassicæ*. As a rule they are retracted towards the surface of the testis. There is but little variety in these organs in the species which I have dissected.

The ductus ejaculatorius.—The two *vasa deferentia* unite, as before stated, at their distal extremities, and are continued by an azygos duct considerably larger in diameter, but short; it is usually slightly invaginated, like the finger of a glove, and wrinkled longitudinally, so that it is capable of comparatively considerable extension and contraction. It is shown in plate II. fig. 1, *d*.

The penis and accessory organs.—Taking *Nothrus theleproctus* as an example, the penis is a small chitinous organ, broader near the distal than the proximal end, but almost pointed at each, the distal end being bulbous (plate II. fig. 3). Near this end there is a chitinous process to which the retractor and extensor muscles are attached. This penis, when retracted, rests in, and is protected by, a second chitinous piece (plate II. fig. 2), which is concave, like half a tube (cut longitudinally); the distal end, however, is closed by a semicircular return, or turning-up, of the half-tube, so as to form a shallow pocket, in the edge of which is a notch, evidently as a guide to the penis when protruded. The walls of the proximal portion of the half-tube are turned outward and expanded, so as to form pyriform blades, doubtless with the object of affording a firmer attachment.

The whole arrangement, when not in action, is retracted within a membranous sheath, which is retained in an open or distended condition by two curved chitinous pieces (plate II. fig. 2). The penis is, of course, in communication with the ductus ejaculatorius. This intromittent organ and its adjuncts bear a striking resemblance to those of *Carabus glabratus*, &c., in the *Coleoptera*: the drawing of the penis of one would almost serve for the other; the membranous tube is Burmeister's præputium, and the chitinous pieces are his horny ridges, or bones, distending the same. The whole of the male organs above described, particularly the testis, vasa deferentia, &c., have a marked similarity to those described by A. Cronberg for *Nesæa coccinea*.*

The copulative suckers, genital plates, &c., have already been well described by Nicolet.

As far as I have been able to observe, the spermatozoa are not mobile or flagellate. I have succeeded in obtaining them from *Damæus geniculatus* by breaking up the testis, and also by pressing the part which appeared to me to act as vesicula seminalis, almost immediately after death; they are small, elliptical, highly refractive, bodies, about the 1-10,000th of an inch in the long axis; they are figured in plate II. fig. 4, highly magnified. They closely resemble, except as to size, the semen of *Tegenaria guyonii* Guerin, one of the spiders which were shown to me by F. M. Campbell, of Hoddesdon; these were non-flagellate, although some spiders appear to have flagellate semen. Blane figures the semen of *Phalangium cornutum* and other species as non-flagellate.†

The Female Organs of Generation.—The general arrangement and position of these, as before stated, greatly resemble those of the male; it is, however, unnecessary to say that the differences are numerous and important. The organs consist of the ovary,

* 'On *Eylais extendens*,' Moscow, 1878 (in Russian).

† Bull. Soc. Vaud. de Sci. Nat., xvii. pl. 6, fig. 23.

the paired oviducts, terminating in an unpaired vagina, the ovipositor, and the copulative suckers and genital plates.

The ovary, or ovaries, for it is doubtful if they should be treated as paired or azygos, consist of a large central sac or gland, which underlies the ventriculus, in much the same position as that occupied by the testis; this I take to be the true ovary, from the walls of which the eggs are differentiated; it will be found fully developed, even when the creature has only lately emerged from the nymphal skin, and when there are not any eggs in the oviducts or other genital organs. At this period the walls of the ovary are cellular, the ova being more or less rudimentary (see plate II. fig. 5). At a later period, when the eggs have become developed, and when their formation and deposition is in active progress, the ovary may be seen to contain numerous eggs in a considerably more advanced state (plate II. figs. 6, 7). This central ovary appears to me possibly to consist of two paired organs, which have coalesced in the central line, much in the same manner as the testes. It is true that the ovary of *Oribata lapidaria* (plate II. fig. 5, a), *Oribata globula*, &c., is an elliptical organ, giving little, if any, indication of a dual origin, particularly when immature, but in *Cepheus tegeocranus* (plate II. fig. 6, a), in the mature ovary, there is a decided indication of a paired origin, there being a marked central constriction, and an approach to the outline of the coalesced testes. In *Nothrus theleproctus*, when the eggs are developed, the ovaries, or portions of an ovary, from the two sides have so slight an attachment to each other that they easily separate, and then appear as paired structures, one attached to each oviduct.

The oviducts.—These, like the vasa deferentia, are two paired ducts, arising, one from near each end of the central ovary; they are long, membranous tubes, which, before the eggs have commenced to mature, appear empty; they are then usually corrugated and plicated organs, of about even dimensions throughout; they are shown in this condition at plate II. fig. 5, b, but it must be understood that this and all the other figures of ovaries are drawn from preparations dissected out and partly extended; when *in situ* they are so much doubled backward and forward that a drawing of them in their actual position would not give much information. As the eggs are formed, and after they have increased a little in size in the ovary, they pass into the oviducts, where the principal growth takes place. The oviducts seem to perform the function of a uterus, and the passage of the egg through them is consequently very slow; the eggs will sometimes be found small in the portion of the oviduct between the ovary and the globular enlargement mentioned below, and large in that between those parts and the vagina; at all other times, or in other species, the eggs all appear about the same size, i. e. usually all fully grown. The oviducts with the

eggs mature are shown at plate II. figs. 6 and 7, *b*; their size then is quite disproportionate to that when the eggs are still in the ovary. The whole oviduct is greatly expanded, and each egg is lodged in a pocket, or chamber, formed by the distension of the elastic walls of the duct, and the flattening of the corrugations. This chamber usually follows the egg in its progress along the duct; in *Damæus geniculatus*, and probably in some other species, the earlier chambers do not appear to follow, or to subside after the passage of the egg, but persist to some extent; this may be due to the egg remaining an unusual time there while acquiring its chitinous shell.

In *Oribata globula*, *Damæus geniculatus*, and some other species, there is, on each side of the central ovary, and joined to it by a short portion of the oviduct, which is of equal length on each side, an almost globular extension of the oviduct, containing a globular body, which has the appearance of an ovum which has not yet assumed the oval form or the chitinous shell, and which almost entirely fills the chamber; these, like the central ovary, are developed when the creature first emerges from the nymphal skin, and while the other portions of the oviducts are unexpanded and empty. Both the central ovary, and certain large cells lying on or in the exterior walls of the two globular extensions, where the latter exist, stain deeply with logwood, &c., whereas the other portions of the oviduct scarcely take the stain at all. Whether these globular chambers be simply uterine, each containing an ovum, which here attains full size, or whether they are, or also function as, spermathecae, or glands secreting a vitelline substance, or one destined to form the egg-shell, is a question which I have not been able to decide to my own satisfaction; some light may be thrown on the question of whether they function as spermathecae when we know how copulation takes place, a matter which is not recorded, and which I have not succeeded in ascertaining; but it is certain that in *Damæus geniculatus*, the egg of which has a very hard chitinous shell, and other species having this globular enlargement, the eggs in the ovary, and between it and the globular body, are small, white, and soft, and show large clear nuclei and one or more round, distinct nucleoli, yelk-division not having more than commenced, even if it has commenced at all; but that, immediately they have passed this point, they have attained their full size, and the shell has become harder and darker, and yelk-division has largely progressed. This would seem to indicate that the globular body is an ovum and the enlargement a special point in the duct functioning as a uterus in these species. The only difficulty of this view would seem to be that the globular expansions, presenting similar appearance, are present in specimens just emerged, which have not any other eggs in the duct. A portion of the ovary with the part of one oviduct up to and including the globular chamber, is shown filled with eggs at plate II. fig. 8.

The duct, even when full of eggs, continues to be plicated longitudinally, so that the eggs, when *in situ*, often appear to lie at the side of the body, one over the other in a bunch; this however is a deceptive appearance, they really follow each other in single file along the duct. The ova sometimes lie end to end, sometimes obliquely across the duct, sometimes almost side by side (through the stretching of the duct); the position of the eggs in the duct is very irregular, I have not ever seen them all following one another end to end at regular distances, and regularly increasing in size from the first to the last, as figured by Nicolet, although the end to end arrangement is not unusual in the species he figures.

These oviducts are what Nicolet calls the ovaries, and what I term the ovipositor is named oviduct by him; I regret using the words in a different sense, but in the first instance it seems to me that his term is scarcely correct, although very natural, he not having observed the central ovary; and in the latter case, although the organ is unquestionably an oviduct, as far as the derivation of the word goes, yet it appears convenient to distinguish it by a name different from the duct between the ovary and the vagina.

The *vagina*, if this be a correct name for an organ which probably does not receive the male one, is a short but wide azygos duct, into which the oviducts lead; it is manifestly the homologue of the ductus ejaculatorius, and is of similar consistency to the oviducts of which it forms the continuation. It appears to vary but little in the different species which I have investigated; its office is to conduct the eggs from the oviducts to the commencement of the ovipositor. The egg does not remain in the vagina for any length of time.

The *ovipositor*, or external vagina of Burmeister, has been described by me in a former publication, and one form of it is well figured by Nicolet. I therefore shall not repeat it here; the same thing applies to the genital suckers and plates.

Nicolet figures the female reproductive organs of *Damæus geniculatus* more on the insect type than the crustacean, i. e. without the central ovary above named, treating my oviduct as the ovary, and drawing it as diminishing to a point at each side, with the eggs regularly increasing in size from the distal point to the vagina. I have not found any specimen in this condition.

It is necessary here to notice a remarkable error, as to the reproductive organs of *Oribatidæ*, for which Dujardin is responsible,* but which has somehow retained a place in modern English works of authority.† The usually keen-sighted French naturalist started with the initial error that the *Oribatidæ* were viviparous: he, having this idea in his mind, did not consider that they could

* "Premier mémoire sur les Acariens," Ann. des Sci. Nat., 3rd ser. iii. p. 5.

† Rymer Jones' 'Animal Kingdom,' 4th ed. p. 309.

have an ovipositor, and as he found the organ he took it to be the penis—a curious idea, for although that organ certainly attains a great length in some of the *Acarina*, e. g. *Proctophylodes* (*Dermaleichus*) *glandarinus* Koch, yet a male organ longer than the whole body, and thicker than the leg, would be an anomaly. Having settled this to his own satisfaction, Dujardin declared the opening below the hinder pair of plates to be the vulva, and concluded that the *Oribatidæ* were hermaphrodite; in this view Dujardin entirely forgot that the animals required an anus, and these errors still survive although Nicolet and Claparède have exposed them.

Respiratory Organs.

Nicolet* describes the respiratory system of the *Oribatidæ* as consisting of two conspicuous stigmata, one on each side of the posterior portion of the cephalothorax, each stigma being funnel-shaped, and opening, by a minute circular aperture, into an air-sac placed transversely in the body and bent upon itself in order to reach the stigma, and of four larger and two smaller tracheæ on each side, the longer ones being distributed two to the dorsal and two to the ventral surface, and forming many convolutions, and the two smaller being allotted to the cephalothorax; and he states that this arrangement is general to all the *Oribatidæ*, and only varies in other families of *Acarina*. I confess that until I began actually to dissect out the respiratory organs I never for an instant doubted the entire correctness of this description, and for a considerable time after my dissections had commenced, I thought that I must have missed the arrangement described by the French naturalist in consequence of the difficulty of the investigation. I found, however, that I could not, in any instance, trace the tracheæ up to the position described, nor could I find any air-sac in this place from which they originated, although I found a sac or tube, apparently glandular, not far off, but with liquid contents instead of air and not connected with the tracheæ. In the course of this investigation I have dissected over fifty specimens belonging to numerous species: I find, with sincere regret, that although many parts of Nicolet's descriptions are correct, yet I am not able to agree with him in other important particulars.

The breathing organs in the adults are usually, as he correctly says, tracheæ (I omit the air-sac question at present); these tracheæ are very long, and so delicate that the slightest attempt to move or separate them, even with the finest badger-hair, usually breaks them, and as they are, as Nicolet says, considerably convoluted, and are interlaced amongst the other organs, I have not found it possible to ascertain with certainty exactly how many there are.

* Loc. cit., p. 410.

I incline to think that it varies. I have also found that instead of being similar in all species the system differs materially.

To commence with the more ordinary arrangement, such as that found in *Oribata globula*, a very good species for seeing the tracheæ, *Damæus geniculatus*, &c. The main tracheæ are simple tubes, never anastomosing, and usually without any branches or dichotomous or other furcations or divisions; indeed I doubt if such ever exist. These tubes are much convoluted, or serpentine, and interlaced amongst the other organs. One large trachea on each side, which I will call the great dorsal, winds above the alimentary canal, very close to the notogaster, until near the posterior margin, when it takes a deeper course, following the alimentary canal, and becomes very difficult to trace. Another large trachea on each side, which I will call the great ventral, passes in a serpentine line along the ventral surface and is in connection with the sexual organs, winding about the oviduct and ovary in the female. Either one or two large tracheæ on each side proceed more along the lateral edges of the abdomen, between the two before described; there are also two or more shorter and finer unbranched tracheæ allotted to the cephalothorax on each side, one pair being distributed to the great muscles of the mandibles. The tracheæ are of almost even diameter throughout the principal part of their length, towards their ends they usually diminish and end in a point; this is not, however, invariable, as the large tracheæ, on the contrary, sometimes enlarge at their distal ends so as to form small bulbs.

The large tracheæ above described approach very near to each other at their origin, and if one does not actually follow them to their commencement (a matter of no slight difficulty), but is satisfied with their appearance after the removal of the notogaster, they unquestionably do look as if they were proceeding to the old so-called stigma (my pseudo-stigma) which seems to be the natural place for them to go to; if, however, the adipose tissue and muscle be removed, and the tracheæ followed to their origin, it will be found that they turn away from the pseudo-stigma, and end separately in the acetabula of the legs. The tracheæ are often enlarged at the commencement so as to form a small air-sac, which either wraps round or is attached to the inner side of the acetabulum: this is seen in plate I. fig. 5 and plate I. fig. 9.

One main trachea only usually proceeds from the acetabulum of each leg, but in some cases two proceed from that of the second leg, and more than one of the small cephalothoracic tracheæ from that of the first leg. In these cases they often join before reaching the stigma and form a very short joint trunk (plate I. fig. 6).

In consequence of the interlacing of the tracheæ among the organs it is far from easy to ascertain which trachea proceeds from each acetabulum, but it appears to me, as well as I could trace

them, that the great ventral trachea proceeds from the acetabulum of the fourth leg, the great dorsal from that of the third leg, the lateral trachea or tracheæ, which wind among the organs and cross right over the body, from the acetabulum of the second leg and the small cephalothoracic tracheæ from that of the first.

I think it possible that the two small tracheæ belonging to the cephalothoracic system find their stigmatic opening at the base of the maxillæ. I should be extremely doubtful about this last point were I speaking only of the species whose tracheæ are described above, as in them these tracheæ, wherever they open, are extremely fine and delicate; but in *Nothrus theleproctus*, &c., referred to below, the air-vessels in this situation are more powerful, and seem as though they were proceeding to the mouth, and it must be remembered that this is the position of the principal stigma, or stigmata, in *Cheyletus*,* *Trombidium*,† *Myobia*,‡ the *Hydrachnidæ*, &c.

In other members of the family, such as *Nothrus theleproctus*, the respiratory system above described is greatly modified: the origins of the tracheæ are placed in the same situations as in the species above described, but the nature of the air-vessels is very different; we no longer have long winding tracheæ of small diameter and very delicate walls, we have instead much shorter organs, thicker and less regular in form, and of a stouter and different texture, by which they may be easily recognized when the body is opened, as they have a thick, silvery appearance not easy to describe. Under a low amplification they assume that slightly iridescent appearance characteristic of a lined object under a power insufficient to resolve it; using a higher power regular cross lines are strongly developed, and a still greater amplification will show an object thickly and regularly covered with small circular bosses which will remind the observer of such a diatom as *Pleurosigma formosum*: this may, however, possibly be a deceptive appearance.

The tracheæ or air-vessels proceeding from the acetabula in the species now being described, are very much shorter and thicker than in the form before referred to, and are not convoluted or interlaced between the organs to the extent which we find in the longer and finer tracheæ of *Oribata*, &c.; at their extremity furthest from the stigma (the blind end) they are usually suddenly diminished in diameter and carried on for a very short distance in a blunt point.

In this form the air-vessels in the cephalothorax, which may

* Fumose et Robin, "Mémoire anatomique et zoologique sur les acariens des genres *Cheyletus*, &c.," Journ. Anat. et Phys. (Robin) 1867, p. 563.

† Pagenstecher, 'Beiträge zur Anatomie der Milben,' Leipzig, 1860, Heft 1, p. 19.

‡ Claparède, "Studien an Acariden," Zeitschr. f. Wiss. Zool., 1868, pl. 37.

possibly arise from oral stigmata, assume greater importance and become some of the largest; a short wide trunk leads into a chamber, from which proceed two cæcal prolongations of unequal length (plate I. fig. 7); these share with the other tracheæ the characteristic of being short, and of the structure described above.

It is very interesting to find this slightly developed and almost rudimentary condition of the tracheæ in the genus *Nothrus*, when we remember that the nymphs and larvæ all through the family usually have the tracheæ in a rudimentary state, and that in this genus the want of hard chitinization in the integument of the adults, their great resemblance to the nymphs, their carrying the cast skins in some species, and other indications, seem to point to creatures where the adult shows less progress from the nymphal stage than in other genera.

In the genus *Hoplophora* I have hitherto failed to discover any tracheæ whatever; Claparède was greatly surprised at the same thing,* he found, beneath Nicolet's stigma, which he calls *peritrema*, three very minute sacs filled with air, which he says are not longer than the width of the stigma, and which, from his drawing, cannot be above the 1-250th of a millimetre in length (his entire creature being over a millimetre); he considers these to be the entire respiratory arrangement of the creature, and to be modified tracheæ, equivalent to the so-called lungs of spiders, scorpions, &c. Although he did not find any leaf-like arrangement within the sacs, or any blood-vascular vessels in connection with them, it must, however, be remembered that Claparède was doubtless relying upon the statement of Nicolet, whose book he quotes, that the tracheæ in other species arose from this so-called stigma.

Claparède expresses his astonishment at the extremely rudimentary condition of these respiratory organs, and he well might do so; one naturally is diffident of questioning a conclusion on this subject arrived at by a man so extremely well acquainted with spiders' lungs as Claparède, but I cannot think that a sufficient respiratory system is shown here; it seems to me far more likely that these minute sacs are connected with the functions of hearing or smell performed by these pseudo-stigmata as suggested by me in an earlier number of this Journal, and it seems to me that there are other means to be found by which aeration could take place in *Hoplophora*. We know that in many soft-skinned acari, as *Tyroglyphus*, *Sarcoptes*, *Dermaleichus*, &c., respiration is performed by the general body-surface without special organs; now in *Hoplophora*, in consequence of the movable ventral plate, so different to that of other *Oribatidæ*, its opening and closing must have a bellows-like action, and great quantities of air must be drawn inside the carapace

* "Studien an Acariden," Zeitschr. f. Wiss. Zool., 1868, p. 512.

and over the delicate lining membranes through which aeration of the blood may well take place.

Before leaving the subject of the respiratory organs I will say a word more as to my not adopting the view held by Nicolet and Claparède, and generally received, that the two organs which look like stigmata on the cephalothorax really are so. One hesitates greatly to attack a conclusion supported by so much authority, but Nicolet is really the only observer who has ever investigated beyond the external appearance; Claparède only dissected *Hoplophora*, and there he did not find anything in connection with the so-called stigmata except the minute air-sacs referred to above. The matter is one of fact; of the tracheæ large enough to be traced I have not ever succeeded in finding one attached to the so-called stigma, or proceeding from any air-sac attached to it, and I have found them originating in other places, as detailed above. Again, the pseudo-stigmata are quite as highly developed in the larvæ and nymphs as in the adults, although the tracheal system is quite rudimentary in the immature forms as far as the observations of other arachnologists and of myself extend, a circumstance (as to the absence of tracheæ in immature forms) also observed by Pagenstecher in *Ixodes* and *Trombidium*, and by Kramer in *Gamasus*, *Tarsonemus*, &c.; and again the extremely small internal opening of the pseudo-stigmata, which is almost or quite filled by the peduncle of the so-called protecting hair (my pseudo-stigmatic organ), is against the hypothesis of their being true stigmata. For what it is worth, it may also be mentioned that in *Pygmephorus spinosus*, a parasite of the mole, which possesses organs much resembling the pseudo-stigmata of the *Oribatidæ*, Dr. Kramer, its discoverer, could not trace any tracheæ to these organs; * although doubtless the weight of this is lessened by the fact that, in consequence of want of specimens for investigation, he was not able to trace the tracheæ to any stigma.

The Super-coxal Glands.

I have said above that Nicolet describes an air-sac which I cannot find, but that I do find a sac, which I believe to be glandular, not far from Nicolet's position, although not attached at the point he names. This sac I propose to call the super-coxal gland, and I have very little doubt, from Nicolet's drawing, that this is the organ he saw and supposed to be connected with what he and others imagined to be the stigma. When the dorsal exo-skeleton of the cephalothorax, and the adipose tissue which underlies it, has been removed, what appears to be the enlarged, blind end of a fine

* "Zwei Parasitische Milben des Maulwurfs," Archiv f. Naturgesch., 43rd year, Bd. 1, p. 257.

colourless sac, may be seen on each side of the body, the seemingly blind end being nearest to the eye; the sac descending obliquely downward and slightly forward, and being attached close to the acetabulum of the coxa of the second leg; a closer examination shows that this is not the only attachment, but that the lower end is apparently bifurcated, and that the second branch is attached much nearer to the centre of the body, and higher in level than the coxal branch. On dissecting out this sac, and carefully extending it, a matter by no means easy, it will be found that what seemed to be the blind end was not the end at all, but that the whole organ is an elongated, sausage-shaped sac, bent upon itself in the middle and taking a single turn, so that the two halves cross, but for some distance the two limbs of the horseshoe (if I may call them so) lie over each other, or are so closely pressed against one another as to appear one; it is only toward the ends that they stand free from each other when *in situ*. The general position of the organ in the body is shown at plate I. fig. 8; and the organ unfolded and showing so much of its minuter structure as I am acquainted with at plate I. fig. 9. The walls of the sac are colourless but highly granular, and apparently of considerable thickness, and, it appears to me, decidedly glandular. Within the sac, i. e. either on its inside surface or in the substance of the wall, and near to the centre (long axis) of the sac, is a double row of highly refractive points, arranged at regular distances, those of the two rows not being opposite, but alternate. A fine distinct double line may be seen uniting these alternate points so as to form a double zigzag line all along the organ, the refracting points being slight prolongations of the zigzags. Three possible explanations of this structure would seem to offer themselves. 1. That the double lines are tubules as in the nephridia of Vermes. 2. That the space between the fine lines is really the zigzag lumen of the sac, possibly running between cells projecting alternately from the opposite walls, and having a cuticularization at the refractive points. 3. That the whole (zigzag and points) is a cuticular strengthening on the interior wall of the sac, the cuticularization being greatest at the refractive points which may be intercellular.

Connected with this tube is a globular body, which has thick but less granular walls, and which is probably hollow; both the gland and this body, when pressed, discharge liquid. There is not any sign of air in either.

What is the office of this organ? I cannot, of course, pretend to do much more than guess, but if a suggestion be allowable, I would point to the nephridia (segmental organs) in Vermes, and the green gland in *Astacus* and other crustacea, and the coxal glands in *Scorpio* and *Limulus*, as those where the analogy may be sought. The resemblance to the former, in particular, is very

considerable in the general form of the organ, and to a lesser extent in the minuter structure; if the double lines be tubules they would be analogous to those in the nephridia.* The sac above described (super-coxal gland) would correspond with the gland in the nephridium, and the globular body with the vesicle.

The position would seem to correspond fairly with that of the green gland and the coxal glands, in the former of which Will and Gorup-Besanez say that they found guanin,† and Huxley says that if this be so there can be little doubt that it represents the kidney, and its secretions the urinary fluid;‡ while Ray Lankester is inclined to look on the latter as homologous with the nephridia.§

I cannot say that I have succeeded in finding any actual opening of the super-coxal gland to the exterior, but the extremely small size of the creatures and of the gland, and the position of the latter, make it very difficult, and indeed Professor Lankester does not seem as yet to have been more successful in this particular respect with the larger forms which he has been dealing with.

In *Hoplophora magna*, about in a similar place to the above-named super-coxal glands (as far as the peculiar form of *Hoplophora* will allow), I find, on each side of the body, a pyriform sac, having the smaller end toward the legs, and the larger, blind end, toward the centre of the body; this sac has walls less finely granular than the glands in *Leiosoma*, &c., but very glandular; I have not found in it either the refractive points or the zigzag markings; it is not bent nor doubled on itself nor twisted; there is a globular vesicle, darker in colour than the sac, on each side of it. These organs require further investigation, but it would appear probable that they are the homologues of the super-coxal gland and vesicle, although presenting considerable differences.

The Exo-skeleton.

The only two points relative to the exo-skeleton which I intend to refer to are:—1. The tectum; 2. The mandible of the species which Nicolet calls *Leiosoma microcephala*.

The *tectum* is a part which owes its name to Nicolet; he alone has pointed out the existence of such an organ; as far as I know it does not exist in any other creatures, and it will be seen from the following remarks, that, in my opinion, it does not exist at all.

Nicolet's statement, when treating of the cephalothorax, is as follows:—"In a large number of individuals, forming the first division of the *Oribatidæ*, the upper part of the cephalothorax is

* See A. G. Bourne, "On the structure of the Nephridia of the Medicinal Leech," Quart. Journ. Micr. Sci., 1880, pp. 283-302.

† Gelehrte Anzeigen d. K. Baierischen Akad., No. 233, 1848.

‡ 'The Crayfish,' 1880, p. 83.

§ Proc. Roy. Soc., xxxiv. (1882) pp. 95-101.

dominated, and sometimes entirely hidden, by a lamellar and tectiform expansion of its base, which advances forward, following its declivity (that of the rostrum), and assuming a form more or less triangular, according as the former is more or less angular, the sides of which expansion are raised oblique projections, often prolonged beyond the front (of the expansion), and always terminated by two setiform hairs. This apparatus, of the functions of which I am ignorant, but which I consider as a protecting organ, and to which I have given the name of tectum, extends from the base of one stigma to that of the others. The lower face of this organ, where it is opposed to the upper surface of the cephalothorax, is not always free in all species; there even exist some in which it is adherent all its length, and then the tectum is only distinguished by its lateral wings, which in that case are usually more developed. In other species this same tectum presents itself as two sub-parallel blades, united by their inner edges, truncated and rounded anteriorly, and through which the body of the cephalothorax may be seen; in this last case the tectum has not any lateral expansions. If I notice these different modifications of the tectum it is because this eminently variable appendage is the best specific distinction that the *Oribatidæ* of the first division present."

Nicolet founds upon variations of the tectum, not only specific distinctions, but even those of sub-families. He describes it in several places, and evidently regards it as a chitinous shelf standing free in the air in many species, except that its basal edge is attached to the hinder part of the dorsal surface of the cephalothorax. This shelf, he says, has turned-up lateral edges in most species; these he sometimes calls the raised borders and sometimes the lateral wings of the tectum; for distinctness I shall give them a name, and I speak of them as the lamellæ.

I have always had great difficulty in distinguishing the species that Nicolet said had free tecta attached by their base only from those which he said had tecta attached by their whole under surfaces by this means of differentiation, but I presumed that Nicolet had satisfied himself that there was such a thing as a detached tectum, and it never struck me to doubt the existence of such an organ until I came to dissect for the purposes of the Ray book; had I doubted it, and trusted to inspection of the living creature, or of dead or mounted examples, I should probably have still considered Nicolet to be right, for certainly in such species as *Cepheus tegeocranus* (*vulgaris*), which is the very type of Nicolet's free tecta, it looks so very like what he described, that I not only should have been, but was, deceived; when, however, I came to dissect the organ away from the cephalothorax in this type species, I found, to my amazement, that there was not anything to come. I then passed a hair under the long projecting ends of the lamellæ, and

found that there was not any difficulty in carrying it back as far as the point where the tectum was supposed to commence, but there it stopped, and that nothing would get it any further, whereas it ought to have passed equally easily into the supposed space between the tectum and the dorsum of the cephalothorax (or vertex as Nicolet calls it). It then struck me that amongst all the very large number of specimens of *Oribatidæ* which I had examined I had not ever seen one where any dirt had got into this space, although it gets into every other place where there is a small hole or depression, and this would be a receptacle just fitted for it; but I thought that possibly a thin liquid, such as alcohol, or even water, might run in where a hair or solid matter would not pass. I accordingly tried, but could not get any to run under the supposed tectum; by these methods, but more especially by careful and frequently repeated dissection of various species, I at last became convinced that the tectum of Nicolet did not exist, and that the appearance of it in *Cepheus*, &c., was an optical delusion. In *Oribata punctata*, &c., the greatly enlarged and horizontal lamellæ somewhat simulate a tectum; this form is Nicolet's sub-parallel blades.

How does the appearance of *Cepheus*, &c., arise? It seems to me that the explanation is as follows:—The lamellæ are real and existing organs, easily seen by even the most superficial observer, and quite easy to get away, by cutting or breaking, but instead of being the upturned edges of a special, detached, horizontal, chitinous organ, they are simply outfoldings of the cuticle of the cephalothorax itself, just as the apodemata which serve as points of attachment for so many of the muscles, are infoldings of the same cuticle; in this manner it is natural that the base of each lamella should be thicker than its summit (or edge), which indeed does not include the true cuticle at all, but only the chitinous secretion from it, and the lamella, being a fold of the cuticle, does not spring sharply at right-angles from the surface of the cephalothorax, but rises in a curve, produced, so to speak, by the dragging up of the cuticle from each side; thus each lamella has a more or less triangular transverse section, the sides being curved and giving considerable extension to the base, *particularly on the inner side*. The dorsal surface of the cephalothorax, from which these lamellæ spring, is convex, and the broad inner base of the lamella filling up the depression caused by the lower part of the convexity, causes the whole space within the lamellæ to appear, and really to be, higher in level than the other parts outside the lamella. Again, the ends of the lamellæ usually rise forming projections, sometimes very short, but often of considerable length, which stand quite free. At the point where the lamellæ cease to be attached by their lower edges, and rise to form the free projections, the two lamellæ are

most frequently joined by a transverse ridge of the same nature, which, in some species, as *Oribata setosa*, *O. orbicularis*, &c., is large and conspicuous, almost as much so as the lamellæ themselves; in most species, however, the cross ridge is much smaller, often a mere line, and it has then the appearance of being the anterior edge of the supposed tectum; as it does not rise sharply on its posterior edge, but slopes up gradually, it would seem to be a natural result of the folding of the cuticle to form the projecting end of the lamella.

The last point I shall refer to is the exceptional mandible of one single species. The ordinary mandible of the family is a chelate mandible which, in all genera, except *Pelops* and the instance I am about to mention, is broadest in the middle and is terminated by a powerful chela (plate I. fig. 10).

In *Pelops* there is marked difference in the shape of the mandible (plate I. fig. 11): the base is broad, almost quadrangular, then the organ suddenly narrows and is continued as a long thin rod, terminated by a very fine chela; this doubtless fits it for exploring small holes. In every recorded instance the mandible is chelate, and this is given as a leading characteristic of the family.

One of Nicolet's species is *Leiosoma microcephala*; in external appearance it is but slightly distinguished from other species of the same genus. It is rare, and I had only two or three specimens; it is my habit to draw the mandible, but this necessitates breaking up the creature, and as they usually vary but little I did not like destroying a specimen; from what I could see, however, I suspected that there was something unusual in the trophi, and I sacrificed one of the three; to my astonishment I found a mandible (plate I. figs. 12, 13, 14), totally different from that of all other *Oribatidæ* which I know of; having somewhat the form of the *Pelops* mandible, but longer in the shaft (or thin portion), and curved instead of straight, but *not chelate at all*, on the contrary, devoid of any movable joint, and regularly serrated along the distal portion for a considerable length so as to serve as a sawing instead of a seizing or tearing organ. It seems to me that with this essential variation it is not possible to leave it in the genus *Leiosoma*, and I propose to create a genus to be called *Serrarius* for its reception.

Finally, I wish to express my thanks to Mr. Charles Stewart, of St. Thomas' Hospital, both for his encouragement to persevere in what I feared might prove to be investigations too difficult for me, and for his able opinions as to the results I had obtained; and also to Mr. E. Bostock, of Stone, and Mr. C. F. George, of Kirton Lindsey, for supplies of living specimens which enabled me to continue the research in London.