difficulties in the way of differential diagnosis of intra-thoracic tumors may be recognized. Moreover, the diversity in the peculiarities of each case, the multifarious character of the pressure symptoms and physical signs, and the absence of a precise order of phenomena peculiar to tumors in this situation, may render a positive diagnosis in the early stages very difficult."

In this case, there had been none of the causes predisposing to disease of the coats of the arteries, but there was offsetting this, the history of laborious occupation, of violent exercise, and of obscure traumatism of seven or eight years previously; pain was nearly absent; this, too, in spite of the evidences of a tumor, as shown by the limited prominence in the left sternal region; pulsation and thrill, murmur, and lateral expansion were absent, important diagnostic factors in aneurismal growths, and the recurrent laryngeal gave no direct sign of involvement, neither hoarseness nor aphonia having existed. The variation in the radials implied obstruction, or pressure from some one of several intra-thoracic causes, not from aneurism alone.

Mediastinal new growths, the majority of which are lympho-sarcomatous, oppose in their earlier stages many obstacles to a conclusive diagnosis. Broadly stated, there are no symptoms or physical signs, nor any precise order of phenomena peculiar to, or definitely diagnostic of intra-thoracic new growths.

The wider the departure from the usual concurrent signs of intra-thoracic tumors; the more diffused the signs of obstruction, and of disturbed mechanism referable to the mediastinum (when thoughtfully considered in connection with the history of the case) the firmer the foundation upon which to rest a diagnosis, of the class of intra-thoracic new growths of which one case is typical.

We were, in considering this case, mindful of the fact, that these growths increase from without inwards, that the slight external tumor might be but a surface indication of extensive deposits, an opinion confirmed by the large area of percussion dullness: that the physical signs of collapse of pulmonary lobules, upper left front, and the evidence of venous stasis in face and neck were conjoint witnesses inseparably connected and associated with the cervical glandular enlargements, an aggregation unimpeachable and conclusive.

Let us now turn to the consideration of the later symptoms found in the rapid growth of the tumor, and in the evidences of obstruction, declared by the external signs, and the local mechanical interference with the functions of the thoracic organs.

At an interval of four weeks the following conditions existed:—

The face, neck and left arm were œdematous and dusky: the neck broadening to its base was much enlarged, irregular in surface, and the supra-clavicular depressions were obliterated; anteriorly, from the sternal notch upwards, the growth masked the outlines of the larynx. Respiratory movements in the left upper chest were absent; the heart's apex was displaced downwards and outwards into the seventh interspace, outside the nipple line; to rapid breathing, there had been added orthopnea, of the most pronounced type, the patient sleeping much of the time with his head bending forward towards his drawn-up knees.

The tumor before noted as a small elevation had so increased in size as to involve the costal cartilages and the ends of the second, third and fourth ribs, and

measured three and one-half by four and one-half inches, with an estimated elevation of one and three-fourth inches; it was a firm, resisting, non-heaving, non-expanding mass, free from pain and not tender on palpation or percussion; there was flatness on percussion, from the left apex to the third interspace and thence to the right of the sternum. Vocal fremitus was absent in this region; respiratory murmur other than transmitted could not be heard; the voice sounds were not distinctive; urgent symptoms, referable to obstruction of the trachea, as well as of the superior vena cava, became more constant, and Dr. Porter's opinion was sought, with reference to surgical measures to secure relief. He decided that the growth in front of the larynx and the evidence of obstruction beneath the sternum precluded tracheotomy, and rendered any operation impracticable.

The evidences of extreme malignancy were found in the processes destructive to life, and rapidly advancing during the few days previous to the death of the patient. Briefly these proofs were, glandular enlargements, cervical and axillary, particularly on the left side, causing obstruction and so great increase in size of the left arm that it was nearly twice the size of its fellow; cyanosis of the face and neck, taking the place of the previous lividity and dusky hue; persistent dilatation of the superficial veins of the neck and face; pressure signs referable to the trachea and left primary bronchi, and in the condition of the heart, displaced downwards and outwards, with marked signs of cardiac failure; most significant of all, a tumor, increasing rapidly along the lines of least resistance, and involving, while not destroying, a portion of the anterior chest wall. We easily recognize in these physical signs the classical phenomena of obstruction, from a quickly developing intra-thoracic new growth.

The rational and physical signs accompanying mediastinal tumors, are complex and varying, and modified as such signs must be by reason of the changed anatomical relations of the structures involved, and the inconstant position of new growths in this region; careful consideration will, nevertheless, reveal the co-existence and close association of diagnostic factors of these three classes, namely, disorders of circulation, disturbances of innervation, respiratory phenomena.

IMPERFORATE HYMEN WITH RETENTION OF THE MENSTRUAL FLUX.¹

BY W. H. BAKER, M.D.

FROM the fact that in the practice of gynecology for twelve years past I have met with but one case, the report of which I wish to present to you this evening, assures me that as a class, they are infrequent.

We often see cases where there is more or less complete atresia of some part of the vaginal canal, as a result of a severe labor, or following some sloughing of the parts after some of the exhausting fevers; but a complete closure of the outlet of the vagina from a congenital malformation, with a well-formed vagina, uterus, and ovaries, occasioning a distention of either the vagina, uterus, or Fallopian tubes, or of all, is of much more rare occurrence.

The report of the case is as follows: Mrs. A. L. was referred to me by Dr. Towne, of Manchester, N.

¹ Read before the Obstetrical Society of Boston, February 13, 1886.

H., who had made the above diagnosis, and desired me to operate. She was a Swede, twenty-one years of age, and had been married six months. From the age of fourteen, she had menses of menstruation, but there had never been any flow. These symptoms varied in intensity, and consisted of pain in the groins, feeling of fulness, and weight in the lower abdomen, sometimes so severe as to render her unconscious for a time. Since her marriage there had been frequent attempts at intercourse, accompanied with pain on her part, and but little satisfaction on the part of her husband. Her general health appeared to be good.

Physical examination showed the external organs of generation well developed, with the exception that the hymen was imperforate, and the meatus urinarius was so much enlarged that it readily admitted the forefinger. It, however, occasioned no incontinence of urine; this latter I also considered a fault of development, rather than a dilatation from the unsuccessful attempts at coition. With one finger in the rectum and one in the bladder, an indistinctly-fluctuating tumor was found to nearly fill the pelvis. By recto-abdominal palpation, no evidence of a distention of the uterus or tubes could be established.

June 24th, 1885. The patient being etherized, and a more careful examination being made, mainly for the purpose of determining whether the tubes were distended, a free incision was made through the hymen, which was about 3-16 of an inch in thickness, and forty ounces of thick, chocolate-colored fluid removed. The uterus and tubes were then found unaffected, and the vagina was thoroughly cleansed with a solution of corrosive sublimate, 1 to 2000. This canal was filled with iodoform gauze, and the patient put to bed.

Subsequently, the temperature was never but once above the normal, which was on the evening of the second day, when it scored 99.5°. The iodoform dressings were changed each second day, which not only insured the cleanliness of the vagina, but promoted the healing of its walls, which were much thickened and granular, bleeding easily, but also in their application and removal, insured against any closing again of the hymen. She returned home in two weeks, and from letters received both from Dr. Towne and her husband, I am assured of her health and happiness. She menstruated but once after the operation, becoming pregnant almost immediately.

By the kindness of the late Dr. J. B. S. Jackson, my attention was called, some years ago, to a most remarkable specimen in the Museum of the Medical Improvement Society, which was removed *post-mortem*, the case having been under his care, and seen by several of the most prominent surgeons of this city practising thirty years ago. Both the uterus and vagina were dilated in this case to such an extent, that the fundus of the uterus reached the umbilicus; the Fallopian tubes also were greatly distended, feeling like sausages on either side of the uterus. Although the condition was recognized by all, yet non-interference was advised, and the patient succumbed to her disease.

In the treatment of the class of cases under consideration, I am sure that it is essential to determine whether the tubes, as well as the uterus and vagina, are distended by the retained menstrual flux, for if so, it will not be safe to make a free incision and empty the uterus and vagina, for either septicæmia will occur from the tubes, or there will be great danger of their rupture occurring from the increased peristalsis of

their walls, excited into action by the contracting uterus. If, then, such condition be present, it would be best first to remove the distended tubes by laparotomy, and then make the free incision through the hymen, and empty the uterus and vagina. If, however, the retention of the fluid does not extend to the tubes, the case may be treated simply by evacuating the vagina, or vagina and uterus, and keeping the walls clean and the opposing walls from uniting by adhesive inflammation.

RECENT PROGRESS IN THE CARE OF THE INSANE.

BY WALTER CHANNING, M.D.

INSANE CRIMINALS.

"THE Connection between Crime and Insanity"¹ is the name of a new book by Sander and Richter, reviewed by Dr. Pelman, who remarks that it has been an accepted fact among physicians who care for the insane, that criminals, who become insane while undergoing sentence, are not fit subjects for insane institutions. Nevertheless, in prison they occasion great trouble, and cannot be prevented from outbreaks of insanity. Whatever plans have been proposed for their provision, the opinion has been pretty unanimous to keep them out of asylums.

Dr. Sander, in his book, takes the opposite ground, and says that, in his opinion, special institutions for the criminal insane are not necessary. Dr. Sander's field of labor is at the great asylum at Dalldorf, where the observations embodied in his book have been made, and they certainly demand serious consideration.

From outside observation of affairs at Dalldorf, Pelman thinks the opposite opinion from that of Dr. Sander would be formed. And this would be strengthened by the communication of Sommer, in 1884, which described the trouble that had arisen from bringing insane criminals together at Dalldorf, in consequence of which it had been necessary to obtain the services of two prison officials, who, to a certain extent, acted as overseers of the division devoted to the care of this class.

Dr. Richter, in the first part of Dr. Sander's book, gives a history of 190 criminals treated at Dalldorf from 1880-1883. The number among these who made attempts to escape is considerable, while not a few succeeded, and others misbehaved themselves. The impression made on Pelman by reading the report of these cases, is that they are not fit subjects for a lunatic asylum.

Sander found that 75 per cent. were in an insane condition when they were sentenced. In only thirty-eight cases were the patients' condition discovered at the time of trial, so that the chance of being unfairly sentenced was only one in three. Out of fifteen paralytics, four only were discovered to be such!

There were six times as many persons among the insane in the institutions of Berlin, who had undergone criminal prosecution, as among the generally healthy population of Prussia, which, Sander thinks, demonstrates the close connection between insanity and crime.

Only eight per cent. were more dangerous and troublesome than the ordinary insane, but the women are left out in this estimate.

¹Die Beziehungen zwischen Geistesstörung und Verbrechen. Von Dr. W. Sander und Dr. A. Richter, Berlin, 1886. Besprochen von Dr. Pelman. All. Zeitschrift f. Psychiatrie. 22tes. Band, 4tes Heft.