medicine seems to have set it to rights." "But it has had none of the medicine." "Then what have you been doing with the child?" "Why, sir, don't you recollect you told me that a friend of yours recommended that sick children should be rubbed all over with salad oil, and that I might try it if I liked, but that you had no faith in it? Well, sir, I did oil it, and the child has been improving ever since."

Bronchitis.—Last January a desperate case came under my care, which in spite of active treatment became rapidly worse. As a last resource I smeared it all over with salad oil, and, to my utter astonishment, there was a marked improvement in the breathing in less than twenty minutes. In a few hours the bronchitis entirely subsided.

Another case—double capillary bronchitis, neglected for several days—came under my relative's notice. He considered it too far gone for medicinal measures, and therefore ordered it to be oiled every four hours. The next day the symptoms had diminished in severity, and on the morning of the third day the child was sitting up in bed taking food, and to all appearance convalescent.

Convulsions.—In these cases the effect of oiling is sometimes truly surprising, the fit ceasing before the completion of the operation, and not subsequently returning. A patient informs me that whenever she observes the symptoms which used to precede convulsions in her boy, she at once oils him, when a calm sleep follows, from which the child wakes up refreshed.

Diarrhœa.—Some time since my cousin had an uncontrollable case of diarrhœa under his care in a child of seventeen months. I advised him to give oil a trial; but he said it was too far gone for anything to be done. I saw the little sufferer myself a day or two afterwards, and ordered it to be oiled every six hours. There was a marked improvement immediately after the first application. By the next day the prostration was gone. Previous to this attack the child was a "puny little thing;" but now (oiling three times a week having been persisted in up to the present time) it is "a splendid boy."

Enlarged liver in a rickety child, with bronchitis supervening.—This case had been under the care of a well-known hospital physician, who gave it up, saying that nothing more could be done for it. I ordered it to be oiled every six hours. After each application a calm sleep followed. In about seventy-two hours the bronchitis began to give way; and a few days afterwards the liver was observed to have diminished in size. The child has not since ailed.

Such is a sample of a few of the cases which have come under my notice. Did I not confine these notes to infantile disorders, a long string of other complaints, which are to be benefited or cured by oiling, might be added. It seems as though the good old Greek and Roman practice will have to come up again. Such is progress!

Kentish-town, Dec. 11th, 1869.

CASE OF PARTIAL PLACENTA PRÆVIA

VERSION; RECOVERY OF MOTHER AND CHILD.

By DR. JASPER CARGILL.

On the morning of the 7th August, 1868, I was summoned to attend Mrs. D—, aged forty-six, who resided nine miles from my residence. She was the mother of a large family, and had never had a bad labour. She wanted three weeks of her full time.

I was informed that, while in her bath on the morning of the 2nd August, Mrs. D— was alarmed by a sudden gush of blood, without any uterine pain whatever. She felt faint, and went to bed. All that day she continued to lose blood at intervals. On the 3rd, she felt better, and there was very little bleeding. On the 4th, slight uterine pains came on, and gushes of blood during each contraction. This continued during the 5th and 6th. Very early on the morning of the 7th, she was seized with strong rigors, and the loss of blood was alarmingly great. A neighbouring druggist was consulted, who at once advised my being sent for.

I arrived at 10.30 A.M., and found Mrs. D— in a very critical state. Her countenance was pinched and anxious; pulse at wrist hardly perceptible. She was in a profuse cold perspiration, and her breathing short and quick. The bed

and bedding were saturated with blood. The pains, which had been constant and feeble, had entirely ceased. Having removed the napkins, I found the vagina filled with clotted blood. On introducing my hand, I felt the os uteri high up, and fully dilated. The membranes were entire, and the child's head presenting. The placenta was attached to the posterior edge of the cervix, and a portion of it detached and curled up, so that the lip of the os could be felt beneath it. I at once gave half a drachm of the liquid extract of ergot, hoping to induce pains, and perhaps bring the head within reach of the short forceps. Having waited ten minutes, I repeated the ergot, by which a few feeble pains were excited. At this period a gush of blood warned me of the danger of any longer delay. Having repeated the ergot, I at once turned. I found no difficulty in reaching the feet; one of which I seized, and turned during the flow of waters. I delivered the child as quickly as I could, and lost no time in removing the placenta. The uterus, on being relieved of its distension, readily contracted. The child (a boy) was still-born for a few minutes, and required the usual means to restore it. When I left the house in the afternoon, both mother and child were in a favourable condition. The mother recovered rapidly, and the baby is now a fine-grown child. The placenta was of the usual size, and the detached portion thick and congested.

Dr. Tyler Smith, in his "Manual of Obstetrics," writes, with reference to placental presentation, "In my opinion, turning is the great operation in placenta prævia when the child is living and viable."

On the 11th of September, 1863, I treated a case of placenta centralis by version, and saved mother and child. I am, therefore, fully persuaded that turning is the operation in these cases whenever the conditions of the case render it possible.

Linstead, Jamaica, 1869.

A Mirror

OF THE PRACTICE OF MEDICINE AND SURGERY IN THE HOSPITALS OF LONDON.

Nulla autem est alia pro certo noscendi via, nisi quamplurimas et morborum et dissectionum historias, tum aliorum, tum proprias collectas habere, et inter se comparare.—MORGAGNI *De Sed. et Caus. Morb.*, lib. iv. Proœmium.

ST. THOMAS'S HOSPITAL.

CASES OF FRACTURE AND DISLOCATION.

(Under the care of Messrs. SOLLY, SIMON, and SYDNEY JONES.)

We continue this week our account of interesting cases of fracture and dislocation which have occurred at this hospital during the past few months, and for notes of which we are indebted to Mr. F. Churchill, surgical registrar.

Treatment of fractured femur in infants.—The difficulty in obtaining sufficient surface for the proper application of the long splint in cases of fracture of the thigh in infants, and the want of sufficient purchase for the fulcrum action of the perineal band, owing to the excessive rotundity and polysarcia of limbs and trunk, render it necessary in some cases to resort to a method different from that usually adopted in the treatment of fracture of the femur in adults. Some cases have recently occurred at the above hospital of fractured femur in infants where the limb was encased in plaster of Paris; but this did not sufficiently control the upper fragment, and the lower fragment being firmly fixed, very great displacement, and consequent distortion, of the limb was the result. It was, therefore, decided to keep the leg at complete rest by laying it on a firm and smooth bed, and covering it with as many sand-bags as were necessary to prevent all movement of the leg, and very satisfactory results were thereby obtained.

CASE 7. *Ununited fracture of femur in an infant.*—M. J. O—, aged ten months, admitted (under the care of Mr. Simon) with fractured femur about the centre of the shaft.

The limb was at first put up with the long outside splint; but the constant movements of the infant caused considerable displacement. Plaster-of-Paris bandage was then applied; but this did not sufficiently control the upper fragment. The limb was then placed between sand-bags, and others were placed across the limb at the level of the fracture, effectually preventing all movement. The bone, which had for some weeks been in the condition of an ununited fracture, now became firmly united, and he was dismissed cured after being submitted to this treatment for about a month.

CASE 8. *Fractured astragalus*.—H. L.—, aged fifteen, a bricklayer's boy, fell from the top of a ladder, a distance of twenty-five feet, and alighted on his heels. He was admitted (under the care of Mr. Solly) with fracture of astragalus of right foot. There was very little displacement of fragments, and the ankle-joint could be moved without giving him much pain. The limb was flexed upon an outside splint, and an ice-bag applied to the ankle. He was dismissed cured four weeks after, without ankylosis.

CASE 9. *Fractured os calcis*.—J. H.—, aged sixty, fell from a height of about six feet, alighting on the back part of the left heel upon some asphalt at the bottom of a saw-pit, and was admitted (under the care of Mr. Simon) with transverse fracture of os calcis. Crepitus felt distinctly on moving the bone. There was considerable swelling and oedema around the ankle. The limb was carefully bandaged upon a Liston's splint, and ice-bag applied. He was dismissed cured.

CASE 10. *Dislocation of right arm into axilla, of ten weeks' duration*.—J. W.—, aged fifty-five, was walking quietly along the road, when he fell on his right arm. He was not sure whether he fell on his hand, elbow, or shoulder, as he was somewhat the worse for drink at the time. Dislocation was not detected until his admission into hospital (under the care of Mr. Sydney Jones) ten weeks after. Chloroform was administered, and Mr. Jones succeeded by careful extension and manipulation in returning the head of the humerus into the glenoid cavity. He was dismissed cured with a useful arm a week after.

CASE 11. *Dislocation of right hip into sciatic notch; fracture of left femur; contusion of pelvis, and wound of scalp*.—J. S.—, aged forty-three, was working on a scaffold when he fell down a distance of thirty feet on to his back. Admitted (under the care of Mr. Solly) with dislocation of right hip into the sciatic notch, fracture of the left femur at the junction of the lower and middle thirds, considerable bruising and contusion of pelvis, and a wound at the back part of the scalp about three inches long. The dislocation was reduced by Mr. Croft flexing the limb firmly upon the abdomen, rotating the limb downwards and outwards, so that the adductors being brought into play tilted the head of the bone back into the acetabulum. The fractured femur was put up on a long outside splint. He is still in hospital, but quite convalescent, having free use of the right leg, union of the left femur in good position, and the scalp wound is now quite healed.

CASE 12. *Dislocated astragalus*.—J. R.—, aged twenty-seven, was tripped up and fell forwards, twisting the right foot under her. Admitted (under the care of Mr. Simon) with fractured fibula and dislocation of astragalus backwards. There was also partial dislocation of the foot backwards, and the astragalus was felt as a distinct rounded prominence behind the external malleolus, upon which it could be moved independently. The dislocation was reduced after careful extension and manipulation. The limb was laid upon a Liston's splint, and she was dismissed cured three weeks after, without ankylosis of joint.

LONDON HOSPITAL.

MISCELLANEOUS CASES.

(Under the care of Mr. HUTCHINSON.)

Keloid.—Mr. Hutchinson has recently had under his care two cases of extreme interest from a clinical point of view, as they presented typical examples of the two forms of keloid disease associated, the one with the name of Alibert, and the other with that of Dr. Addison. The subject of Alibert's keloid was a girl about five years of age, who had been burnt severely over the right shoulder, the front of the

neck, and the upper extremity, and also less severely over several other parts of the body. The cicatrices resulting from these injuries had been converted into the irregular patches and thickened buttons of indurated tissue so characteristic of this disease. Mr. Hutchinson, in some clinical remarks upon this case, stated that the keloid of Alibert was a fibroid growth of cicatricial tissue, and frequently affected the scars caused by burns and scalds, and those left after an attack of small-pox. It was entirely a local affection; whilst the keloid of Addison, on the other hand, which always grew from healthy integument, and never from a cicatrix, was a constitutional disease, and probably some form of neurosis. There was no analogy between the two diseases, except that each presented similar elevated, smooth, and pinkish patches of indurated tissue. The keloid of Alibert never contracts, and thus differs from cicatricial tissue, which, when formed after extensive burns or scalds, is characterised by this property, and gives rise to great deformity. This form of keloid generally sends out irregular outgrowths from the circumference of its patches, and in most instances is very troublesome to the patient on account of its extreme irritability. It is inadvisable to remove the diseased patch with the knife, as the keloid growth always returns in the scars formed after the operation. Mr. Hutchinson has found from his experience that there is invariably a tendency in this form of keloid to diminish in extent, and in many instances to disappear altogether. In some cases these favourable changes occur more rapidly than in others, but the subjects of the disease may always be assured that sooner or later it will come to an end. The second case in the surgical wards of this hospital is one of Addison's keloid, which has been described also under the names of morphea alba and of scleroma. The patient, who is a lad about eight years of age, presents very markedly in the lower limbs the hide-bound condition caused by the contraction of hard, dry, parchment-like integument. Symmetrically arranged over both scapulae are indurated nodules of the disease. This form of keloid differs from that of Alibert in its tendency to contraction, its evident dependence upon some constitutional affection, and its frequent distribution along the course of nerves.

Ulceration of the palate in young patients.—In November last, a stout, healthy-looking girl, eleven years of age, was under treatment in the surgical wards for ulceration at the back of the mouth, which resulted in destruction of the uvula and the adjacent soft parts. There was also an oval perforation in the soft palate. Mr. Hutchinson stated to his class that as ulceration and destruction of the soft palate in an adult generally indicates tertiary syphilis, so does the existence of a similar disease in a young subject at once suggest the idea of hereditary syphilitic taint. But it has been observed that destruction of the palate at an early age is very rarely if ever associated with the signs of inherited syphilis. In the present patient the teeth were not deformed, there was no keratitis, and periosteal thickening could not be detected. If, as we have good reasons for believing, early ulceration be due to hereditary syphilis, then this lesion characterises a group of cases quite distinct from that in which the ordinary symptoms of inherited disease, such as pegged teeth and keratitis, are presented. The most effective treatment of sloughing of the soft palate in young patients is the local application of nitric acid or the acid nitrate of mercury.

Carbolic-acid dressings.—On Dec. 6th, a young man was admitted into the London Hospital with a severe compound fracture of the humerus immediately below the epiphysal line between the head of the bone and the shaft. There was a considerable projection of bone, and about two inches of the shaft had to be removed by the saw. The wound was then well washed with a solution of carbolic acid, and afterwards constantly dressed with this fluid. Mr. Hutchinson states that there can be no doubt that carbolic acid, when applied locally, will prevent and arrest suppuration. It will not, however, guard against inflammation of the soft parts around the seat of injury. In the present case suppuration had been prevented; but the injured arm had been attacked with diffuse cellular inflammation, extending to below the elbow. Although carbolic acid will not prevent unfavourable changes of this kind in the injured soft parts, there is good reason for believing that it controls the force of the inflammation, and hinders the suppuration which is so frequently the result of this local affection.