under similar circumstances to make a journey of any length particularly undesirable. Some of the more active occupations entail similar unpleasant consequences.

Finally, these cases serve to emphasize the fact that gonorrhœa may be an exceedingly grave disease, which does not exhaust itself in the urethra, and that its possibilities render it an object worthy of study by the conscientious student, and of honest care by the attending physician.

A CASE OF HODGKINS' DISEASE.¹

BY HAROLD WILLIAMS, M.D.

I WAS called, July 10, 1886, to see Mr. D., a young gentleman, twenty-seven years of age, a chemist by profession, and married. He had been ill since May, 1886, and had come to Nantucket in June, 1886, with the hope that a change of climate might prove beneficial. The history of his illness was nearly as follows: In May, 1885, while out walking, Mr. D. suddenly felt dizzy, and his vision became defective. He walked home with difficulty, his difficulty in locomotion being due, he thought, to an inability to hold himself erect. Previous to this he had been perfectly well. There was no history of malaria nor syphilis. A few days later Mr. D. became totally deaf, and, at this time, a swelling made its appearance over the inner maleolus of the left leg. This swelling increased in size; was intensely painful, and was of a bluish color, but subsided at the expiration of ten days. There was no injury. As the swelling at the ankle began to subside, several smaller swellings appeared in both legs and thighs. These smaller tumors also disappeared, and were, in their turn, followed by a swelling as large as an orange in the right buttock. At this time there was paralysis of the left leg and thigh. This swelling in the buttock slowly diminished in size for three weeks, when it again increased, and again diminished. His physician told him that his spleen was enormously enlarged; and one oculist who was consulted said that his impaired vision was due to hæmorrhage, while a second oculist assured him that there had been no hæmorrhage. This takes us up to Christmas, 1885, on which day the patient went out-doors. After this there was gradual improvement, though it is probable that the swelling in the buttock never totally disappeared, and the deafness still persisted, though slightly improved.

On July 10th, when I first saw the patient, he was in bed. He complained of much pain, and was so deaf that I could scarcely make him understand me. His body was well nourished; the lips were red, the finger-nails pink, and there was no appearance of leucæmia. Pulse 89, temperature 102°. There were three tumors: one on the left side of the thorax, bounded by the sternum, the third rib, the posterior axillary line, and the eighth rib. This was the largest, and I considered it the forward extension of enlarged axillary glands. The second was in the neck, extending downward beneath the left clavicle, and upward to a point two inches above it. I could not estimate its lower boundary, because of the pain caused by percussion. The third tumor was in the right buttock, and seemed to lie above the *gluteus maximus*. It was as large as

a very large orange. These tumors were very hard to the touch, and the skin over them was of a reddish-blue color, hot and painful, but movable. There was enlargement of the spleen, huskiness of speech, slight difficulty in swallowing, and defective vision. The patient was taking iodide of potassium and small doses of morphia, to relieve pain. This treatment I continued.

On the following day the patient was much worse, and in the afternoon I found him in a semi-comatose condition, from which it was difficult to rouse him. He had vomited at different intervals throughout the day, and there was much difficulty in swallowing and articulation. Pulse 150, and very feeble; respirations 40. The condition of the tumors seemed, outwardly, the same.

The vomiting, feeble and rapid pulse and respiration, and the disturbance of the functions of the larynx, seemed to me to denote pressure by the tumor in the neck on the deep-lying nerves of the trachea, and I warned the family that I considered the case nearly hopeless. Brandy and coffee were given by the rectum.

Instead of dying, as I expected, the patient rallied, and for three days there was slight amelioration of the symptoms, though the tumors seemed outwardly the same. At this time I prescribed Fowler's solution in $\mathfrak{m}$ x doses, three times a day, to be increased up to $\mathfrak{m}$ xx. After this there was constant improvement, although, as the glands subsided, a condition of extreme leucæmia, with furuncles and a carbuncle, supervened. The patient was of a waxen hue, his lips were colorless, and the nails white. To the naked eye, the blood seemed barely tinged with red, and, under the microscope, showed a marked change in the proportion between the red and white corpuscles. As no proper apparatus was at hand, I could merely estimate by what Osler calls "the rough and ready method," that the white corpuscles seemed nearly as numerous as the red.

In October the patient considered himself perfectly well, and there were no symptoms or signs of his illness, with the exception of the deafness, which, though much diminished, was still a cause of serious annoyance.

This case I consider to have been one of Hodgkins' disease, or pseudo-leucæmia lymphatica, followed by leucæmia.² The diagnosis would seem to lie between lymphatic leucæmia and pseudo-lymphatic leucæmia, and is based upon the pathological condition of the blood, and upon clinical differences. In lymphatic leucæmia there is an increase of the white corpuscles, and in pseudo-lymphatic leucæmia there is little or no such increase. In the present case no such analysis was made, but nothing in the aspect of the patient suggested such an increase until the swelling of the glands began to subside. But, on the other hand, the absence of the external appearances of leucæmia; the absence of hæmorrhages and diarrhœa; and the enormous swelling of the glands, would seem to indicate Hodgkins' disease. In either case the prognosis is unfavorable, and doubly so in a case of such long duration as the present.

I have ventured to report this case, in spite of the imperfection of my notes, because it seems to me to demonstrate the decided advantage following the administration of large doses of arsenic, prescribed as it was in this instance, to a patient who had been ill for a period extending over fourteen months. The idea

¹ Read before the Section for Clinical Medicine, Pathology, and Hygiene of the Suffolk District Medical Society, March 9, 1887.

² Vide case of Fleischer and Ponzoldt, quoted by Osler in *Pepper's Syst. Med.*, Vol. III, p. 633.

was suggested to me by an article by Karnvski,³ who, in 1884, reported three recoveries under this treatment. The arsenic was given in gradually-increasing doses, until the dose of ℥xx t.i.d. was reached, in which quantity it was continued until there was puffing of the eyelids and nausea. Then it was replaced by quinia and iron for a week, when it was again resumed in the same manner.

This treatment has been continued in gradually-decreasing doses up to the present time, the patient now taking ℥iii of the Fowler's solution, t.i.d., every other week. I have not seen him since October, 1886, but in a letter I received from him last week, he reports: "I am perfectly well, with the exception of my deafness, and weigh 170 pounds — not a bad weight for a man of my size."

Primary Cytogenic Anæmia	Leucocytic	{ Splenic Lymphatic Medullary }	Leucæmia.
	Non-Leucocytic	{ Splenic Lymphatic Medullary }	Hodgkins' Disease, Lymphatic Anæmia.

Clinical Memoranda.

A CASE OF ALEXANDER'S OPERATION.¹

BY JOHN B. SWIFT, M.D., OF BOSTON.

Mrs. B. came under my care at the Carney Hospital, March 18, 1886, with the following history: She was thirty-seven years old, born in Canada, and had had one child. The labor was a severe one, and she had been torn. Since then she had suffered from "falling of the womb," the organ coming outside. By various means she had kept it up, so that she could work, until the friction caused by the bandages had rendered the parts sore, and she would be obliged to stop work until the abrasions had healed. Lately, she had been wearing a cup-shaped pessary, but with no better success.

An examination showed rupture of the perineum to the sphincter ani, bilateral laceration of the cervix, a small recto-vaginal fistula, and complete procidentia of the uterus. The cervix was hypertrophied, excoriated, and there was a deep ulceration of the tissues, caused by the rim of the pessary which she had been wearing.

The uterus was replaced and held in position by cotton tampons, wet with glycerine and tannin, and the patient kept in bed until the excoriated mucous membrane had healed. Attempts were then made to keep the uterus in place by various pessaries, but without success, and finally, it was decided to perform the Alexander operation of shortening the round ligaments. The operation was explained to the patient, and she readily consented. Owing to the extensive laceration of the cervix, and the hypertrophy of the uterus, it measuring almost four inches, it was thought best to first repair the cervix, hoping, by this means, to reduce the hyperplasia, and thus lessen the weight, so as to give a better chance for success in the succeeding operation. This was done April 17th, and resulted successfully.

Three weeks later, with the assistance and advice of Dr. F. B. Harrington, the Alexander operation was performed.

¹ Read before the Section of Obstetrics and Gynecology of the Suffolk District Medical Society, February 16, 1887.

³ Berl. Klin. Wochen., 1884.

I will not enter into the details of the operation, as they probably are familiar to you all, but will only say that the ligaments were easily found, and shortened about two inches. By drawing on the ligaments, the uterus readily ascended in the pelvis, and was held in position by Dr. Harrington while the ligaments were being secured. They were not cut, but were folded and tucked into the wound, being fastened by silk sutures passed through the pillars of the ring and the tissues at the bottom of the wound. The incisions through the skin were closed by catgut sutures, and the wounds dusted with iodoform and dressed with sublimate gauze.

As there was no perineum, it was feared that a pessary would not stay in place, so the vagina was packed with cotton pledgets, and a T-bandage adjusted. The catamenia, which had never been regular, came on the second day after the operation, and the packing had to be removed, a pessary being substituted, which proved, in the end, to work successfully.

For three days the patient was nauseated continuously, and vomited everything taken by the mouth. She was nourished by enemata. Hyperdermic injections of morphia and atropia over the stomach were given for the nausea, but whether or not they did any good, I cannot say. At any rate, on the fourth day she could retain food, and from that time on had no further trouble.

The temperature was never above the normal, and, aside from the vomiting, the convalescence was uninterrupted. She was kept in bed for three weeks, and then allowed to sit up, but she did not walk any for another week. At the end of that time the pessary was removed, and she was allowed to go about and do light work. On my visit, the next day, I was told that the uterus had again descended; but an examination showed that what was considered the uterus, was the anterior vaginal wall, which was prolapsed. The uterus was in good position, and held firmly in place.

It was decided not to repair the perineum at that time, but to wait and see if the uterus would remain in place without this additional support. This was explained to the patient, and she was told to return at once, should there be any indication of the old trouble. She came to the hospital last September, about four months after the operation, saying that she was all right, and had been doing house-work. An examination showed the uterus just as it was on her discharge from the hospital, well up in the pelvis, slightly anteverted. The prolapse of the anterior wall, of course, was the same, but she declined the perineal operation, saying that she was well enough as she was.

— Hartmann describes in the *Berliner Klin. Wochenschrift*, 1886, p. 612, the results of some experiments in the self-induction of œdema and albuminuria, by the exclusive use of certain forms of nourishment for a given time; for instance, he ate chestnuts, lentils, peas, cheese, etc., each for a series of days, and with no other ingestum but water. He twice produced general œdema, once with and once without albuminuria; the former, under an exclusive diet of sausages, the latter with bread only, (1,000 grams in twenty-four hours). The uncomfortable symptoms which the enthusiastic experimenter induced all passed away with a return to ordinary methods of living.