56. *Case of Foreign Body in the Eye.* By JAMES DIXON, F. R. C. S. E., Surgeon to the London Ophthalmic Hospital.—The following case of expulsion of a foreign body, which had remained *eight years* in the anterior chamber of the eye, shows that the formation of a fibrinous cyst around such fragments of metal, does not ensure the patient against the recurrence of inflammation, and enforces the propriety of immediately removing them whenever it can be done without serious injury to the organ.*

P. G., aged 35, a shoemaker, slenderly built, and of pale, unhealthy complexion, came to the London Ophthalmic Hospital, January 10, 1848, complaining of intolerance of light, pain, and impaired vision, in the right eye. The cornea was clear; the sclerotic presented a vascular zone; the iris moved sluggishly when exposed to the light; the pupil was drawn a little downwards and inwards. On the lower and inner part of the iris, midway between its ciliary attachment and the edge of the pupil, lay a small, rounded mass, the size of a mustard-seed, which seemed to be a foreign body, thinly coated with fibrin. The patient could still see large letters, but within three days had been able to read small print, with this eye. He stated, that eight years ago, the eye was struck (he supposed with a shot) as he stood a few yards distant from a man who was shooting sparrows. Pain and redness of the eye came on, and he consulted a medical man, who gave him medicine which made his mouth sore. The sight was not much affected at this time, and it remained good and useful—almost as good as that of the other eye—for about three years. Inflammation then returned, and he applied a second time for medical aid; mercury was administered, and at the end of a fortnight he was so much better, that he discontinued his attendance. No further inconvenience was felt until the beginning of the present year. On his first visit to the hospital, he was leeches and purged, and then calomel was ordered night and morning, with opium. Within three days the foreign body was more than half denuded of its fibrinous covering, and proved to be a thin, flat scale, like a fragment of a percussion cap. A fresh deposit of fibrin soon took place, and until the end of June inflammation continued, with varying intensity, but the foreign body was never sufficiently free from fibrin to allow of its being extracted without a risk of injuring the lens, which remained perfectly healthy. During February, fresh deposits of fibrin occasionally took place, and were absorbed again: by the middle of March the fibrinous effusion on the hinder face of the cornea was so extensive as to hide the lower half of the pupil, and fine red vessels were seen passing into this fibrin.

On the 10th of April, a little white elevation, with a dark dot in the centre, appeared in the middle of the fibrinous patch on the cornea, as if the substance of the latter were softening, and about to give way. Three days later a fine black point protruded; it was readily seized, and a minute scale of metal extracted, which, on being tested, proved to be copper. No escape of aqueous humour followed its removal, as it had become completely enclosed by fibrin, which formed a barrier behind, at the same time that absorption of the cornea was going on in front.

June 5. A dense opacity hides the inner half of the pupil, and, except a slight adhesion of the iris, is the only morbid result of the injury. The iris is active, the lens perfectly transparent, and vision good.—*Dublin Quarterly Journal*, Aug. 1848.

MIDWIFERY.

57. *On the Causes of the Endemic Puerperal Fever of Vienna.* By C. H. F. ROTH, M. D.—(*Proceedings of Royal Med.-Chirurg. Soc.*, Nov. 28, 1848.) There are three lying-in departments in the General Hospital of Vienna. To one of these strangers are not admitted. Of the two others, to which only the author's

* The late Mr. Walker observed, that copper was "a substance incapable of solution within the eye."—*Oculist's Vade Mecum*, 1843, p. 325.