peritonitis became general accompanied by tympanites, and the patient died on the 30th of January. On the occasion of Dr. Grichard's first visit on the 23d of January, attention was called to the condition of the husband: he was suffering from facial erysipelas. On making inquiries, it was found that on the 15th of January he had been seized with shivering and fever; the same evening erysipelas began on the nose and then spread to the forehead, scalp, and posterior part of the neck. The attack ran a mild course, and by the 30th of January the erysipelas was quite cured. Three important dates may now be noticed:

Jan. 15, 1885. Onset of facial erysipelas in the husband.

17th. Delivery of the wife.

20th. Onset of puerperal mischief.

The husband, though mostly absent at work, returned home every day; in spite of his erysipelas he continued at work till January 23d, and continued as usual to share the same bed with his wife: it was only after the visit on the 23d that he began to occupy a separate bed, and by this time his wife was already seriously ill. The midwife cannot be suspected of conveying infection, for she had not been in any way exposed, and had arrived too late at the labor to undertake any manipulative interference. An investigation into the antecedent history of the wife proved that she had not been exposed to any other sources of infection; her child remained well, and, though of necessity living in the same room, suffered no trouble in connection with the umbilical wound.

The following deductions seem warrantable:

(1) The husband communicated the infectious germ to his wife; the portal of infection was through the external or internal genital organs.

(2) This single case does not allow of our determining the nature of the troubles of lying-in women described under the name of puerperal fever; but in the present case one common infectious agent or miasm must have been the common starting-point of facial erysipelas in the man and septicaemia of the internal genital organs in the woman.

(3) As a prophylactic, the physician ought to remove every case of erysipelas from the neighborhood of lying-in women, and therefore to forbid all cohabitation before, and for some time after delivery with the husband, if the latter is suffering from erysipelas.

Strict antiseptic precautions ought to be carried out in the surroundings of the woman before, during, and after confinement. The accoucheur and the nurses should not forget that they may be the direct medium of contagion, and should, in consequence, neglect no antiseptic precaution.

A lying-in woman should never be placed in a room previously inhabited by a case of erysipelas, until the chamber has been completely disinfected.

BROW PRESENTATIONS.

Dr. E. BLANC, in the *Arch. de Théologie* (July, 1885, p. 614), analyzes and studies an interesting work which has just been carried out by M. DEVAUS, on delivery in brow presentations; it is proposed in a rapid analysis to call attention to the facts, either new or original, which have been brought forward in this investigation. M. Devaus divides his work into five parts, viz., the history of the subject, the signs and frequency of these presentations,

their mechanism and etiology, their diagnosis, and, lastly, the therapeutical indications to be derived. The historical part it is not necessary to dwell upon, nor upon the signs of these presentations furnished by the anatomical peculiarities of the frontal and neighboring regions; attention may be turned at once to the question of frequency. Cases of brow presentation do not seem to be very common. Heinricius, in the Maternity of Elsinfors, found only 12 instances out of 5000 confinements; Mangiagalli, during a period of five years, collected 64 cases, 17 of which occurred at the Maternity of Milan, but does not say out of what total of labors. Devens, examining the records of the Clinique d'Accouchements at Lyons for the last five years, found 5 cases in 1402 labors.

By what mechanism is the expulsion of the head effected in these cases? It may be described briefly as follows: The superior maxillary bone corresponds to the occiput in vertex presentations, to the chin in those of the face; whatever the position of this bone at the onset of labor, the expulsive forces will always tend to bring it beneath the symphysis pubis, which is to furnish it with a *point d'appui*, while the longitudinal diameters of the head, the naso-bregmatic, the naso-sagittal, and naso-occipital clear in succession the pelvic outlet and the vulva. After the occiput has become freed from the posterior commissure, the mouth and chin will appear in front of the pubis. The deformity of the head in these presentations is considerable; the head compressed in its suboccipito-mental diameter assumes the form of a triangle, the base of which, viz., the occipito-frontal diameter may measure 13, 14, or even 15 centimetres (5.1, 5.5, or 5.9 inches). Such a lengthening of the longitudinal diameters of the head would appear a source of danger to the vulva and perineum, but it is to be noted that at this stage the face rests by the superior maxillary bone or by the nose against the back of the symphysis, while the forehead is free in the vulvar orifice; a naso-occipital diameter shorter than the occipito-frontal clears the vulva, and it is only after the occiput is disengaged from the posterior commissure that the chin appears below the pubic arch. Too great a distention of the soft parts is thus avoided. Devens admits an equal likelihood of these presentations becoming transformed into those of the vertex or face; Mangiagalli objects to this; he compares the fetal head to a triangle having its apex at the forehead and its base formed by the occipito-mental diameter; but one of the sides, viz., the fronto-mental, is the shorter; instead, then, of the base of the triangle engaging parallel to the pelvic inlet, it will be inclined at an angle, the forehead will descend lower in the pelvic cavity than the occiput, and the engagement of the head in the form of a brow presentation is thus rendered possible: clearly, if this presentation undergoes a transformation, the face should succeed to the brow. Belluzi never saw a brow presentation end in a vertex presentation. Which of these views is correct? Probably no distinction should be made according as the head is slightly engaged at the inlet, or has descended well into the cavity of the pelvis. In the first case the position of the head may change, producing readily the presentation of either face or occiput; in the second, the chin being lower than the occiput, a change to the occipital presentation is less likely.

Among the causes which produce brow presentations may be quoted the resistance of the cervix, uterine obliquity, hydrocephalus, thoracic or ab-

dominal dropsy, twists of the cord round the neck of the fetus, small size of the child, and, lastly, pelvic contractions. Observations made by ourselves experimentally, as well as the opinion of numerous authorities, confirm the belief that contractions of the pelvis play an important part in producing the presentations now under consideration.

The diagnosis is not difficult, as a rule; the prognosis is serious for both mother and child, but especially for the latter. M. Devars puts the infant mortality at nearly fifty per cent., while other authors make it only twenty or twenty-four in the hundred. In 9 cases recorded by M. Devars, delivery occurred spontaneously in 3, and in 6 forceps were used; in the 3 cases two mothers and two children died, while in the 6 cases only one infant succumbed.

Regarding treatment, it is to be noted that in most cases operative interference is called for; among the modes of procedure most approved are change of presentation, effected by manual manipulation, version, the use of forceps and of the lever, and, finally, as a last resource, craniotomy. While the head is still at the pelvic inlet, change of presentation is particularly applicable; indeed, it may be attempted even when the head is already in the pelvic cavity, though in this latter case we should endeavor, if possible, to bring down the chin.

BINODIDE OF MERCURY AS A DISINFECTANT IN OBSTETRICS.

This subject was brought forward by DR. E. P. BERNARDY, at a meeting of the Obstetrical Society of Philadelphia, on June 4th (*Medical News*, Aug. 8, 1885).

In three cases, the lochia had become offensive after delivery. The uterus was washed out with biniodide, and within twenty-four hours the offensive odor ceased, and from that time the coexisting symptoms began to improve. The method he had pursued in making the solution was: Take three grains and a half of biniodide of mercury, well triturated in a mortar, and rubbed with one quart of boiling water, slowly added, giving a solution of 1 to 4390. This solution he had found non-irritating, but equally efficacious in its action to bichloride 1 in 2000. Indeed, in one case in which the bichloride had been first employed, the result appeared to be decidedly in favor of the biniodide. Moreover, he points out that, owing to the smaller quantity of mercury in the latter, the risk of salivation is diminished.

CASE OF "MISSED LABOR" WITH CAESAREAN SECTION.

DR. STANLEY P. WADSWORTH reports (*American Journal of Obstetrics*, July, 1885) the following instructive case: Mrs. McD., aged thirty-two, married, had one child thirteen years ago, but had not been pregnant since. Last menstruated in January, 1884; usual symptoms of early pregnancy followed. At end of April had symptoms of miscarriage—pain and hemorrhage—for some days, followed by discharge from vagina "like thick molasses."

May 2. Temperature and pulse raised. Cervix enlarged, tender, and discharging a thick bloody fluid in small quantity; abdomen tympanitic, somewhat swollen and sensitive. In right iliac fossa a tumor was apparent to sight and touch, about as large as closed fist, pear-shaped. The above men-