The Journal of the American Medical Association

Published under the Auspices of the Board of Trustees

VOL. XLVIII.

CHICAGO, ILLINOIS, JANUARY 26, 1907.

No. 4.

Original Articles

GOITER.

WITH PRELIMINARY REPORT OF THREE HUNDRED
OPERATIONS ON THE THYROID.

C. H. MAYO, A.M., M.D.

Surgeon to St. Mary's Hospital.
ROCHESTER, MINN.

The general impression to be obtained from a review of the surgical literature of America would be that diseases of the thyroid gland are greatly on the increase. This is probably not the case, but the public has learned that operations on goiter are not as fatal as were supposed from the results obtained when operations were made as a last resort on patients suffering from this disease and in a moribund state. The fact is that the mortality attending the operation (excluding cancer and advanced cases of exophthalmic goiter) compares very favorably with other major surgery; and, in the hands of those experienced, with much of the so-called minor surgery.

SURGERY OF THE THYROID.

The surgery of the neck never seemed to enjoy the popularity of that of the abdomen, and goiter operations are not even to-day sought by the majority of surgeons. Those who were competent did not sufficiently often encourage operative relief until the absolute necessity rendered the mortality so high as to be almost prohibitive. The lay impression became such that operative measures were permitted only in the most advanced cases, while semi-surgical, medical, mechanical and electric treatment flourished. In most cases death came slowly and was looked on as a relief from suffering, a result which satisfied everybody; but the death which follows operation is a shock to all, and in but few instances is there a consideration of the fact that medical treatment has previously failed. Neither is the urgent necessity of the operation nor the desperate condition of the patient taken into account, but the death is made a debit against the surgical side. Especially is this true in cases of exophthalmic goiter.

Surgery of the thyroid is one of the most satisfactory operations made, giving immediate relief with brief disability. The best known surgeon of to-day is Kocher, and wherever his name is mentioned the first thought connects it with the removal of goiter. It is through his monumental work in this line that his reputation traveled far and wide. Among ordinary goiters, in 70 operative cases before 1850, Kocher placed the mortality at 40 per cent. In 400 operations between 1850 and 1883, it fell to 15 per cent., and since 1883 has dropped to less than 3 per cent., Kocher's own being 0.4 per cent in simple cases. In Riverdin's report of the cases of death in 93 cases, respiratory conditions, suffocation and pneumonia caused 43 deaths, hemorrhage 19, infection 13,

shock and nerve injury 9, cardiac failure 6, while only 4 were from tetany and myxedema.

DISEASES OF THE THYROID.

The diseases of the thyroid gland can be classed in a few groups—functional derangements, inflammations, hypertrophies and tumors. Each of these groups presents material for most exhaustive study, and very much has been written concerning them in monographs and books on the ductless glands and various types of hypertrophies and tumors. The present paper, being on the general subject of goiter, will discuss but briefly much of interest in relation to operative work.

In Europe there are certain localities which present such large numbers of people with goiter that the disease

Fig. 1.—Large cystic goiter measuring 31 inches in circumference, including neck.

is considered endemic, and, while it is said to be due to the use of waters from the mountainous districts, the disease is also endemic in some of the flat countries, as in the Indian Punjab and plains of Lombardy. Hereditary influences may also play a part. In the United States there are no such marked localities. The disease occurs in all regions and climates, among rich and poor, food and water having apparently no definite relation to its cause. Some regions appear to have more than their due proportion from possessing more accurate medical observers. The same may be said of exophthalmic goiter, which is often attributed to shock, with about the same proof that birth marks follow maternal impressions.

THE THYROID GLAND.

Development.—The thyroid gland is developed from a median process of the pharyngeal hypoblast which bifurcates below and forms pyramidal lobes, i. e., the isthmus and part of each lateral lobe, the remainder being formed from lateral outgrowths from the pharynx. In most vertebrates they remain distinct, but unite in mammals, the various portions being fused in the human embryo at about the seventh week. In some cases the fusion is incomplete, and separate thyroid bodies are noticed laterally or below the normal location. In others one or more lobes may be missing. The foramen cecum in the tongue marks the point of inversion and the remains of a disappearing structure, the duct or tract of the gland. It is in these lines of inversion, like branchial cysts, that accessory thyroids are found, and in the upper tract especially are so often the cause of serious conditions, as when found in the tongue and also in the median line of the neck above the isthmus. The so-called pyramidal lobe is present in from 30 to 40 per cent. of all people.

Anatomy.—The thyroid body is somewhat like a horse shoe in shape and rests on the trachea, with one lobe on each side, connected below by the isthmus which crosses the upper tracheal rings. These lobes are about two inches long, being smaller at the upper pole. The right one is usually the larger, and the entire weight is from one to one and one-half ounces. The gland is invested by a thin fibrous capsule which divides posteriorly, a portion lining the posterior and inner surface, while a part passes to the opposite side behind the esophagus. This investment explains the production of pressure from tumor growths on both of these structures and the occasional appearance of tumors between them. Fibrous bands also unite the gland to the trachea, which cause the thyroid to move with it. Trabeculae of connective tissue pass into the gland structure and subdivide into the framework for the alveoli and small vesiculae and serve as a space for the lymph channels which extend throughout the gland, performing the function of the ducts of this organ, as has been pointed out by King, Horsley and Hürthle. They have shown that these channels contain the same colloid secretion as is found in the gland vesicles.

The blood supply of the thyroid, considering its size, is remarkable for its extent and also for the freedom of anastomosis. The superior thyroid artery from the ex-

ternal carotid supplies the upper pole on its inner side, dividing and entering the capsule. The inferior thyroid from the thyroid axis enters the capsule below at the hilus. The main veins are the superior, middle and inferior, although many others seem to develop in diseased organs. The nerve supply is from the sympathetic. In intimate relation with the right inferior thyroid artery is the recurrent laryngeal nerve, which lies in the space between the trachea and the esophagus, and is so often affected by pressure of tumors, by operation or scar tissue as to cause hoarseness. On the left side the recurrent is usually more deeply set and not in such close relation to the artery, as was pointed out by Sifton.

Parathyroids.—In each side of the neck behind or within the investing capsule of the gland are the parathyroids, two small gland-like bodies somewhat resembling in structure the suprarenals. Little is known concerning them, but they appear to be closely associated with the thyroid in function, and it is probable that when the whole of the thyroid body is removed and the operation is not followed by untoward consequences the patients have active parathyroids or accessory thyroids. It is claimed by Humphrey that a fatty degeneration of these bodies is found in exophthalmic goiter. I could not verify the statement. The experimental study of parathyroids would tend to show that they hold a most important position in the metabolism of the body. Their removal causes tetany and, as they are small, they may easily be removed on the back capsule of the extirpated gland. The removal of one or two in an operation for colloid goiter may not prove serious

Fig. 2.—Enucleation of cystic goiter; capsule opened.

but in exophthalmic goiter it may prove most dangerous in adding to the shock. To some extent, we can see how enucleation of encapsulated goiters would seldom cause either tetany or myxedema, as a portion of the gland is left and the capsule protects the parathyroid as well as the laryngeal nerve.

Function and Secretion.—Much has been written on internal secretions and, while our knowledge has continuously grown, there is still very much to learn about the ductless glands. A consideration of the functions of the various structures of the body shows that they are controlled by accelerator and depressor nerves, the stimulating and inhibiting forces. Every muscle action to be effectively controlled is balanced by another which opposes it. In other words, there must be a constant equilibrium of forces. When we consider the effects of

the secretion of the suprarenal gland with that of the thyroid, it would seem that in several respects they oppose each other. Such a fact explains, to some extent, the condition of cretinism, in which from lack of a functioning thyroid there is an arrest of development. Through the unopposed action of the suprarenal bodies there is a capillary contraction with circulatory starvation which affects the cerebral cortex, as well as other tissues. The intermittent feeding of thyroid extract causes increased growth and general improvement in appearance, with considerable mental development. The mental development is only sufficient in most cases to convert a passive, helpless imbecile into an active, restless and destructive idiot.

HYPOTHYROIDISM.

The loss of the function of the gland seems to be cause for more serious consequences in the young than in those who have reached their complete development. Young adults who lose their entire thyroids will retrograde in intelligence, while the older individuals frequently develop myxedema, in which we have the myxedematous condition of the skin and subcutaneous tissues, dry skin, loss of glandular activity, loss of hair and mental dullness. Apparently these cases are often overlooked unless well developed. The thyroid increases in size commonly at puberty and in the latter months of pregnancy. Such enlargement usually subsides after a period. Oliphant Nicholson, of Edinburgh, speaks of the enlargement of the thyroid during pregnancy and of the fact as shown by Lange that those who do not have such enlargement have an increased tendency to albuminuria and eclampsia. Kocher reports 70 per cent. of cachexias in 34 cases of total excision. Garre reports 50 per cent. in 67 cases. Billroth had 52 cases develop tetany, with 9 deaths, and 11 developed cachexia. The London myxedema committee only found 24 per cent. in 224 cases of total excision. Considering the remarkable conditions which occur from a loss of the secretion of the thyroid, we are not surprised at the serious derangements from an oversecretion or an abnormal secretion.

EXOPHTHALMIC GOITER.

In 1835 Graves, and five years later Basedow more thoroughly, described the disease known by their names, or by the term exophthalmic goiter. These conditions may be described by various writers in a somewhat different way, yet they all undoubtedly mean the same con-

dition which is characterized by an excess of the activity of the gland. A better and more expressive term is hyperthyroidism. It occurs much more frequently in women than in men and the greater number are between 20 and 40 years of age. It is not markedly hereditary and, while shock may be a contributing factor, it is difficult to elicit a probable cause in the majority of cases.

There are several symptoms in exophthalmic goiter which alone may occur in other conditions, yet, alone or associated, lead the diagnostician to consider the possibility of exophthalmic goiter. They are the enlargement of the thyroid, tachycardia, exophthalmos, muscular tremor and general nervousness. The gland is usually enlarged, but may not be palpable in exceptional cases.

With the exophthalmos may occur von Graefe's symptom, the lagging behind of the upper lid in looking down, or Stellwag's sign, the retraction of the upper lid. Möbius described an occasional insufficient accommodation without diplopia. Exophthalmos may be absent in 20 per cent. of the cases. The heart action may vary from 90 to 180 a minute, a case of medium severity being about 120 to 140. Systolic murmurs are not uncommon. Muscle tremor is easily elicited even in mild cases by having the hand extended, finger spread, palm down.

Associated symptoms are excessive perspiration, occasional rise of temperature and digestive disturbances, chief of which are vomiting and failure to assimilate. Attacks of such diarrhea are not unusual. The onset of the disease may be rapid, that is, during a few days, or it may be over an extended period of

time before being noticed. Many medical men still consider the disease to be a probable intestinal toxemia and look on the goiter as a coincidence, although all of the symptoms may be produced in healthy individuals by feeding thyroid, and even exophthalmos is now recorded as occurring following the administration of dried thyroid for obesity. The gland in a typical case usually presents the true hypertrophy, a wonderful increase in cell activity and the cell areas in the vesicles are greatly increased in layers and by infolding. The cell form also changes in shape. The gland is dry and hard after removal as compared to the ordinary juicy colloid or adenomatous condition. The association of enlarged thyroid in these cases is not definitely understood.

In our work at St. Mary's Hospital, we have operated

Fig. 3.—Muscle section higher than skin incision. Stab drainage.

on exophthalmic goiters which were hardly palpable and have been surprised at the increased size of the gland over the normal when exposed. After all it is a question of increased secretion, absorption and delivery by the lymphatics, not necessarily the retention and development of a tumor. If the exophthalmic gland was no larger than normal it would be many times more active. It is a question if the failure of lymph absorbents in certain areas may not, following hemorrhage or increased secretion, play a part in the production of some types of cysts. Exophthalmic goiter may occasionally develop in a case of goiter of long standing, and then there are diffuse areas of hypertrophy in the general tumor. The best classification of the exophthalmic type is: 1, vascular; 2, hypertrophic; 3, hyperthyroidism developing on previously existing goiter.

A fibroid condition of the gland may follow recovery from exophthalmic goiter, and the consequent loss of function later may cause myxedema. Ord and Mackenzie tabulated 56 medical cases of exophthalmic goiter which had lasted five years or been fatal. There were 10 recoveries, 11 made incomplete recovery, in 13 there was considerable improvement, 4 were unchanged, and 14 (25 per cent.) were fatal. Others make more favorable reports.

With the general hygienic and therapeutic treatment of exophthalmic goiter we like the action of belladonna. We also make use of the x-ray. From its known action on the lymphatics and the glands it exerts a favorable effect on the over-activity of the gland in exophthalmic goiter, and in some cases seemingly develops something of a capsule and partially changes the character of the tumor. While its effect may not be permanent it is a valuable adjunct in preparing advanced cases for surgical procedure. We have had no personal experience with the use of serum from thyroidectomized animals or milk from thyroidless goats. The cases and reports we have seen have given conflicting and not convincing evidence. J. Rodgers, of New York, is now experimenting with a thyroid serum which it is claimed is especially valuable in exophthalmic goiter. It is possible that much may yet come from autotoxic serum treatment.

There is a type of thyroid, usually enlarged throughout the whole gland, appearing at about puberty and during the succeeding ten years which Bloodgood calls simple goiter. In these cases dilated acini are filled with comparatively normal colloid. Similar conditions often occur in pregnancy. In most of them the tumor disappears without treatment. Obstinate cases can be cured by some external application or treatment with iodine or organotherapy. In fact, it is reported (Bull's system) that Kocher claims 90 per cent. of all goiters can be so improved by medical treatment as to render operation unnecessary.

Typical exophthalmic goiter may also exist as a symmetrical enlargement. The irregular enlargement may be adenomata, cysts or unevenly developed colloid goiter. Malignant disease in the thyroid also occurs in irregular enlargements, and Bloodgood states truly that when the disease has progressed so far that a differential diagnosis from benign tumor is possible it is also probable that it is too far advanced to cure. For patients with irregular nodular tumors early operation should be advised. Sarcomas may be more uniform in appearance and cystic. Most of the bulging tumors, evenly rounded and covering the front of the neck, are the cystic or colloid cystic type, with a heavy capsule, in which may be a thinned-out layer of the remains of the gland.

In some cases pressure atrophy seems to have destroyed all the gland, yet, as complete removal of these encapsulated masses or cysts does not often cause myxedema, it is probable that the system becomes accustomed to the very gradual withdrawal of the thyroid secretion. In three cases we have removed all of such encapsulated growths without cachexia, although it is possible that accessory thyroids were carrying on the function. One of the three was a large single cyst, the other two were cystic colloid, one measuring 22 and the other 19 inches across the tumor from one side of the neck to the other.

Loss of voice from tumor pressure occurs more often in malignant than in other tumors. The adenomatous growths, fetal and cystic, are common and represent a species of goiter frequently found in children and young adults. Cysts may follow hemorrhage, but more commonly develop from adenoma and colloid degeneration. Echinococcus is infrequent, but must be considered. The necessity of a second operation on goiter is rare, according to Bergeat, being required in only twelve cases out of 600.

Wolff states that when one-half of the gland has been excised for hypertrophic goiter the remaining half always diminishes. This is confirmed by Riverdin, Kocher, Poncet and Wöeffler. It has also been our experience. We do not believe in the semi-surgical treatment of goiter, that is, the injection, puncture and drainage methods which are nearly as dangerous as radical operative measures. Exothyreopecty or open displacement of goiter, Jaboulay's method, is merely mentioned.

THE RELIEF OF EXOPHTHALMIC GOITER.

The removal of the cervical sympathetic ganglia, as advocated by Abadie and Edwards and popularized by Jonesco and Jaboulay, we have not considered as effective as the operation on the diseased organ itself. Rehn's report makes it less than half as favorable. Surgeons do not seem to appreciate the effect of this operation and the ligation of arteries, also the experimental work on the parathyroids, in destroying the lymph channels which drain the thyroid. Division of the isthmus for the relief of pressure is not effective, as the pressure is on the sides of the trachea and not on the top except in cancer.

During the past seventeen years we have operated on 300 thyroids, with 11 deaths. Of these 110 were for exophthalmic goiter, with 9 deaths in all and but two in the last 64 cases. Two deaths in the early work were from accepting almost moribund cases as surgical risks. Among the remaining operations, representing cysts, colloids, adenomata and parenchymatous, including 9 malignant, tumors, there were but 2 deaths, 1 on the eighth day from pneumonia, the other from tracheal collapse following removal of carcinoma. One case was a large lingual thyroid, 3 were tumors of the pyramidal lobes, 2 only were accessory thyroids in the superior line of inversion, 2 were incisions and drainage for acute inflammatory conditions, 2 were for sarcoma.

We have been much pleased with the results of the surgery of exophthalmic goiter. Of those who survive the operation, 50 per cent. make an early recovery, 25 per cent. improve of the main symptoms during several months, and 25 per cent., while greatly improved, have occasional temporary relapses of the tachycardia and tremor. The exophthalmos is often one of the last symptoms to disappear.

Anesthetic.—The choice of an anesthetic is more often determined by the idiosyncrasy of the operator than the necessity of the case. There were 13 operations made with cocain infiltration. Of these 3 were completed with chloroform. While there are occasional cases in which local anesthesia may be necessary, we have noticed but little difference in the shock or general condition of the cases from the anesthetic employed. We now use ether anesthesia, preceded twenty minutes by hypodermic of 1/6 gr. of morphin and 1/120 gr. atropin. The table is placed in an elevated slanting position with the head up (reverse Trendelenburg). The tumor is rendered prominent by a roll beneath the neck if it does not interfere with respiration.

Operation.—We prefer the collar or transverse incision. In tumors of medium size this crosses the center of the tumor, in larger tumors it crosses the upper third and the lower flap is split vertically to the sternum if necessary.

The incision is through skin and platysma muscle, the flap being raised to expose freely the muscles covering the gland, namely, the sternohyoid and thyroïd, the inner portions of the sternomastoid and omohyoid. In medium-sized growths muscle separation will permit of the delivery of the tumor. The muscle section of the sternohyoid and thyroïd group, if made, should be near their upper attachment so as not to interfere with their nerve supply and also to break the line of scar formation from that part of the skin. After the removal of the gland, the severed muscles are carefully united by suture. The upper section also permits of early ligation of the superior thyroïd artery which is the key to the situation.

The more firm and rounded tumors, with a strong capsule, can be readily enucleated after incising the capsule and penetrating the gland tissue to the cyst. We usually loosen the gland from the outside of the capsule first. On viewing the tumor, the true capsule has the luster and appearance of peritoneum. If one is not sure, incise between the vessels. Should hemorrhage be severe, lifting up the tumor by its capsule will reduce the bleeding until sutures can be placed for its control. In parenchymatous and colloid tumors we make extirpation of one lobe and the isthmus. After exposing it the upper pole is elevated and the superior thyroïd artery is cut between the ligatures. The lateral veins are clamped and cut, the lower pole is elevated into the wound, incision is made along the outer posterior border, and the capsule is brushed off with gauze to the median line. As a rule, the inferior thyroïd is now ligated close to the tumor as the capsule is brushed back. This leaves the nerve behind and usually out of sight. The isthmus is clamped and closed by suture. If the enucleation method for adenoma and cysts leaves a badly torn lobe it can be removed. Sometimes extirpation of one lobe is made and enucleation of cystic masses is done on the opposite side, after the method of Porta, Billroth and Socin. The resection method of Mikulicz was only employed in one case, but with good result. The ligation of the thyroïd arteries as a method was not practiced. We avoid rough handling of the portion of gland to be left.

Should there be a large area of cut gland exposed it is burned over with carbolic acid, followed by alcohol, or often Harrington's No. 9 solution is applied over the cut tissues to close the lymph absorbents and favor drainage. If there has been but little traumatism we do not drain. Large incisions and large cavities we

drain temporarily. All exophthalmic cases are drained as freely as the most septic wound.

These cases absorb some wound secretion containing colloid, causing rise in temperature and increase of pulse the next day. To delay absorption, especially in exophthalmic goiter, patients are given large quantities of salines by the bowel directly after the operation and for the first two to four days. If not retained by the bowel, they are given subcutaneous administration.

The wound is closed by subcuticular suture, great pains being taken to unite the platysma to prevent spreading of the scar.

OPERATION FOR THE SHORTENING OF THE BROAD LIGAMENTS.

AN ANATOMIC AND CLINICAL STUDY.*

ARTHUR E. HERTZLER, M.D., PH.D.

KANSAS CITY, MO.

The material underlying this study was as follows: Forty adults and a larger number of pelves of newborn infants were dissected. Microscopic sections were made in various planes of the broad ligaments. Serial sections were made of these structures in fetuses of from three to six months, and also of pelvis of younger

Figure 1.

embryos. The whole clinical study is based on 46 cases operated on in the past four years.

Works on anatomy are in accord in describing the broad ligament as formed of two layers of peritoneum. It is described in a recent work by my distinguished teacher, Waldeyer,¹ in a general way as composed of two serous layers, between which are vessels and nerves, many smooth muscle fibers and more or less connective tissue.

This generally accepted opinion becomes questionable when it is considered that the peritoneum, strictly speaking, can be dissected from the pelvis complete, except from the fundus of the uterus, leaving the ligaments intact. This furnishes indisputable evidence that the accepted description of the broad ligaments is inadequate, and that they are true ligaments.

In corroborating this statement the dissector must have a clear notion as to what is comprised by the peritoneum. It is incorrect to regard all the fibers below the endothelial layer of cells as comprising the peritoneum. I have in course of preparation an extensive

* This article was received in July, 1906.

¹ Read before the South Kansas Medical Society, April 18, 1906. 1. "Das Becken," p. 530.