Examination showed a dorso-lumbar kyphosis and a marked deformity of the toes of both feet: they were red, swollen, and two of the toes of the left foot were covered with ulcers. There were also ulcers on the right toes but they were healed. There was some power in the feet, but hardly any in the toes, a bilateral foot drop and a steppage gait were present. The motor and trophic disturbances were associated with marked anesthesia of the toes and feet, the lateral surfaces of the feet and legs being merely hypesthetic; a little higher up there was hyperesthesia. The calf muscles and the large nerve trunks were tender. The ankle reflexes were lost, the patellar was markedly diminished on the left, normal on the right. The plantar reflex was almost absent on the left, totally absent on the right. The cremasteric was absent on the right. The abdominal reflex was lost above and below the umbilicus was very inconstant. The internal organs, genito-urinary apparatus, the cerebral nerves, the serologic and urinary findings were negative. The combination of trophic, sensory and motor disturbances were thus almost exclusively confined to the feet and could be caused either by a pure peripheral lesion, a multiple neuritis, by a posterior root lesion—*tabes*, or by a spinal-cord lesion of the two lowest lumbar and two upper sacral segments. These four segments make up what Minor described as the *epiconus*.

Tabes could be easily excluded, as the lumbo-sacral seat of this disease usually gives pronounced root symptoms (shooting pains, ataxia, genito-urinary disorders) which were absent. *Epiconus* lesion was represented by the bilateral peroneal paralysis, steppage gait, lost Achilles jerks, integrity of the knee jerks, of the genito-urinary organs and anesthesia in the lumbo-sacral segments. The lumbo-sacral kyphosis and the injuries also spoke in favor of *epiconus* lesion, but the alcoholic history, the peculiar amnesia and the tenderness of the calf muscles and nerves of the legs spoke rather for a multiple neuritis. The ulcers and mutilations of the feet and toes of thirteen years' standing must therefore be ascribed to the latter. Such marked trophic disturbances are very uncommon in neuritis where the trophic symptoms are usually rarer than the sensory or the motor ones and do not attain the degree of severity as shown in this patient. In the case of *Lepine* and *Porot* they were equally severe—on the hands—but were of shorter duration, while in this patient they existed many years and were for a long time the only sign of polyneuritis, and for this reason this is a unique case.

CASES FOR DIAGNOSIS

By G. W. Hall, M.D.

CASE 1. The patient, whose case Dr. Hall wished to present for diagnosis, had died and the gross pathology of the brain was demonstrated. He reported a case of a man who came into the hospital in a comatose state which had developed suddenly, three weeks after the onset of loss of memory, some headache, emotional instability, and on one occasion vomiting.

The examination showed all the deep reflexes to be exaggerated, a double Babinski and Kernig were present, and at first a slight hemiparesis on the right side was demonstrated. The pupils were fixed. He ran a temperature ranging from 99° to 102° F. The pulse, at first very slow, 54 to 56, increased to 116 to 124. The spinal fluid showed a cell count of 147 with a negative Nonne and Noguchi on one occasion and slightly positive on another. There was no optic neuritis. On section a glioma in the region of the basal ganglia on the left side was found.

CASE 2. Dr. Hall presented a case of a man who for fifteen years complained of shooting pains in the epigastrium and of vomiting. The pain was worse in the morning and associated with vomiting which at times was quite

severe. Examination showed diminution of both knee jerks with irregular pupils which reacted very sluggishly to light. The spinal fluid showed a cell count of 10 lymphocytes on the first and 40 lymphocytes on the second examination with a faintly positive Nonne and Noguchi. The Wassermann reaction was negative upon the serum, but three plus positive on the spinal fluid. The patient gave a history of drinking liquor to excess for a great many years. Dr. Hall thought the case one of both gastritis and gastric crises of tabes.

Dr. Grinker said that the specimen which was passed for examination reminded him of the fact that in gliomata hemorrhages are not at all uncommon. In his own experience he recalled a case which was almost as interesting as this one, in a patient who was secretary of a labor organization. He received a trauma from a brick which fell upon his head, causing a scalp wound, and what was considered at that time an abscess of the brain was the result. He was treated and a month later developed a rightsided hemiplegia with aphasia. He was in a stuporous condition and the diagnosis in that case appeared to be a cerebral abscess, considering the fact that only a month previously he had sustained the trauma to his head, which was suppurative in character. He was operated upon but the surgeon failed to find the abscess and the patient died. A post-mortem was obtained and a hemorrhage was found which involved the internal capsule, the motor tracts, and extended outward to the speech centers, so that he had a complete motor aphasia. The hemorrhage occurred in a glioma which was traced to the anterior horn, where there were evidences of two old hemorrhages. It appeared that not until the third hemorrhage when the motor areas had become involved were there any symptoms. In examining the history of the case, the only evidence of something wrong was that two or three months previously this secretary, who had always been prompt in collecting the dues for his organization and giving the members credit, had forgotten the names of those who paid him and made mistakes in his books. He had never complained of headache and had no other symptoms of brain tumor. The specimen was interesting because it showed three kinds of hemorrhages of different age; the first was probably two years old, the next one year old, and the last one was quite recent.

Dr. Patrick wished to know whether Dr. Hall had carefully examined for sensory disturbances on the abdomen, as cases with gastric crises usually showed something there when they failed to show anything elsewhere.

Dr. Hall said he had examined carefully and had found no sensory disturbances.

A CASE OF DERCUM'S DISEASE

By Julius Grinker, M.D.

The patient, a man of about fifty years of age, came to the hospital for nervousness and incipient delirium tremens.

He was weak and nervous, complained of failing vision and was short of breath, especially on exercise. He entered the hospital for the first time in 1909 for a pronounced mitral lesion. Even at that time he related that shortly after he had developed heart trouble he had noticed several small lumps on his body which were very painful and tender on pressure and occasionally caused spontaneous pain. In 1906 appeared the first nodule on his left arm. Later a second one developed on the opposite side, but not in symmetrical order. Then nodules appeared on the trunk, abdomen and thighs. These were small tumors under the skin which had the peculiar feel of fatty tumors. No evidence of their existence was seen on the skin itself. In many cases there has been noticed a peculiar whitish spot over the skin