seen in skin lesions of variola, properly prepared, in the early stage of the disease:

1. *Cytoplasmic Bodies*.—These may be small structureless bodies, one to four microns in diameter; homogeneous ameba-like bodies lying in vacuoles; larger reticular bodies eight to ten microns in diameter; small dot-like bodies about one micron in diameter. In general, these bodies are best seen in the edge of a young vesicle. They occur in the cytoplasm of normal cells, degenerated cells and between the cells. They may also be seen in the corium before the formation of the vesicle. The presence of these bodies in the skin before there is any definite anatomic lesion is an important point in diagnosis.

2. *Intranuclear Bodies*.—I have not been able to find intranuclear bodies in the earliest cases. It is suggested by Councilman, Brinckerhoff, and McGrath,⁵ that these bodies represent a further stage in the development of the cytoplasmic bodies. As the presence of these bodies is not necessary to a diagnosis they will not be discussed here. They are best found from the time of disappearance of the cytoplasmic forms until repair has begun. For complete descriptions of these bodies the reader is referred to the authors previously mentioned and to Calkins⁹ and Tyzzer.¹⁰

All cases are not typical and variations of type must be considered in the laboratory just as in the clinic. Departure from the typical may consist in the incomplete development of the lesion (abortive type), or in the diffuse hemorrhage that takes place in some cases before the appearance of any lesion. I have not had the opportunity to study typical hemorrhagic cases, but Councilman⁵ states that in cases where purpura preceded the eruption "there was no doubt about the character of the process. The skin showed the earliest stage of true lesions. In all cases there were parasitic inclusions." While laboratory diagnosis of variola may be sometimes quite as difficult as is the microscopic determination of the malignancy of certain tumors, I believe, however, that with the microscope fewer mistakes are liable to be made by a microscopist of reasonable skill in smallpox diagnosis than in tumor diagnosis.

CONCLUSION.

In conclusion, then, by means of acetone fixation, paraffin imbedding and hematoxylin-eosin staining, we can make a laboratory diagnosis of variola from a small skin snipping in an average time of three hours, based on a specific histologic lesion and on the presence of certain bodies in or adjacent to the lesion.

LARGE CARCINOMATOUS TUMOR OF THE LIVER.

REMOVAL SEVENTEEN MONTHS AFTER NEPHRECTOMY FOR CARCINOMA OF THE LEFT KIDNEY. TEMPORARY RECOVERY.

THOMAS S. CULLEN, M.B.
BALTIMORE.

Although Brunn in 1870 removed a small hepatic tumor, the surgery of the liver does not seem to have attracted much attention until Lin, in 1886, successfully performed a partial resection. Since that date, however, more than 100 resections of portions of the liver

have been reported. Among the exhaustive treatises on the subject those of Terrier and Auvray,¹ in France, and of Langenbuch,² in Germany, deserve especial mention. In America the name of W. W. Keen³ of Philadelphia is intimately associated with the development of the technic of liver surgery. He has not only reported three cases of resection, in all of which recov-

Fig. 1.—Kousnietzoff's blunt needle. This large curved needle has a rounded point. Its shape on cross section is shown in the small sketch below. The straight needle is preferable for mattress sutures.

Fig. 2.—Kousnietzoff's liver suture. The needle is threaded double and is carried in and out through the liver, producing a continuous mattress suture, which is cut up into single mattress sutures before tying.

Fig. 3.—Auvray's liver suture. This is the so-called chain suture, and after insertion is drawn taut.

ered took place, but has also published an exhaustive review of the previous literature. Nor should it be forgotten that a Baltimore surgeon, Dr. Louis McLane

1. Terrier and Auvray: *Les Tumeurs du Foie*, Rev. de Chir., 1898, vol. xviii, p. 403.

2. Langenbuch: *Chirurgie der Leber und Gallenblase*, Deutsche Chirurgie, Stuttgart, 1897.

3. Keen, W. W.: *Resection of the Liver*, Boston Med. and Surg. Jour., vol. cxxvi, p. 405, 1892; *Pennsylvania Med. Jour.*, 1897; report of a case of resection of the liver for removal of a neoplasm, with a table of 76 cases on resection of the liver for hepatic tumors. *Ann. of Surg.*, September, 1899.

9. Calkins: *The Life of Cytoryctes Variolæ*, Guarneri, Jour. of Med. Research, vol. xi, No. 1.

10. Tyzzer: *Etiology and Pathology of Vaccinia*, Jour. of Med. Research, vol. xi, No. 1.

Tiffany,⁴ was one of the first Americans successfully to remove a tumor of the liver.

The Nature of Hepatic Tumors.—Few operators have reported more than one case, and as each has had his own idea of the proper nomenclature of tumors, there is naturally much diversity of description. With the more thorough study of the subject, the varieties will undoubtedly be materially diminished in number. The list of those given is as follows:

Constricted and prolapsed lobe of the liver.
 Fibroma.
 Syphiloma or gumma.
 Tubercle (occasionally single, may reach 3 or 4 cm. in diameter).
 Adenoma.
 Lymphadenoma.
 Cysts: (a) echinococcus cysts; (b) biliary cysts (large and lined by one layer of cylindrical epithelium).

or in the old. They are, however, most common between the twentieth and fiftieth years. As has been pointed out by Keen, hepatic growths are much commoner in women than in men. In 68 cases analyzed by him, 13 of the patients were males and 55 females. He thinks that the relative immunity in men is probably due to the looseness of the clothing worn. In the additional cases which I have collected, 4 of the patients were men and 12 were women.

The symptoms of hepatic tumors are in no way characteristic. Practically, they are those belonging to any growth developing in the right upper abdominal quadrant. When the growth is primary and situated near the lower edge of the liver in a thin patient, it is some-

Fig. 4.—Adenocarcinoma of the kidney (natural size). Gynecologic Pathologic No. 6364. This is a longitudinal section of the hardened kidney. There is a well-defined outer capsule which consists almost entirely of connective tissue. Passing off into the growth from the capsule are trabeculae (a) which divide it into large and small alveoli. The growth presents a decidedly mottled appearance. At b it is finely granular. In numerous places it has undergone necrosis and disintegration as seen at c. The ureter is little, if at all, enlarged.

Angioma: (a) congenital; (b) acquired.
 Lymphangioma cavernosum.
 Cancer: (a) primary; (b) secondary.
 Sarcoma: (a) primary; (b) secondary.
 Endothelioma.
 Aneurism of the hepatic arteries.

Echinococcus cysts, cancers and syphilomata are the most common of hepatic tumors. The growths vary greatly in size, some being not more than 2 or 3 cm. in diameter, while others, as in Pfannenstiel's case, may be so large as to render the abdomen larger in its dimensions than that of a pregnant woman at term.

INCIDENCE AND SYMPTOMS OF HEPATIC TUMORS.

Tumors of this character may be found in the young

times palpable, but if the individual be stout the growth can readily be overlooked.

DIAGNOSIS.

Hepatic tumors are rarely correctly diagnosed until the abdomen is opened. In the cases reported the growths have been mistaken for tumors of the kidney, ovarian cysts, uterine myomata, mesenteric and pancreatic tumors, tumors of the colon, aneurisms of the aorta, growths in the gall bladder and stomach, and abscess of the liver. This is equivalent to saying that in a series of cases they have at one time or another been confused with almost every tumor that may be present in the abdomen. Such a tumor is usually first noticed on the right side, in the region of the lower edge of the

4. Tiffany, Louis McLane: Maryland Med. Jour., 1890, vol. xxxiii, p. 531.

liver. In the rare instances in which the movement of the diaphragm is transmitted to the tumor, we possess a diagnostic sign of much value. Occasionally, percussion or palpation, or both, will enable us to detect a definite connection between the tumor and the liver.

Even after the abdomen has been opened and the tumor exposed, one is often still in doubt as to the exact nature of the growth. If the patient has given a definite history of syphilis, it may be assumed, with a relative degree of certainty, that a gumma is present; on the other hand, if a carcinoma of the stomach be pres-

and after the abdomen had been opened the general picture suggested an inflammatory process of the liver. Furthermore, on careful examination of the left renal region no evidence of a return of the former growth could be detected. The internal organs were in good condition, and in the liver no other nodules could be found. The absence of any inflammatory reaction at the junction of the tumor with the normal liver substance was the only sign that pointed against the presence of an abscess. In short, it was found to be impossible to determine with any degree of certainty whether

Fig. 5.—Secondary carcinoma of the liver (one-half natural size). The liver is markedly displaced downward. Occupying the greater part of the right upper quadrant of the abdomen is a hepatic growth. Downward this almost reaches the anterior superior spine; medianward it passes nearly to the umbilicus and laterally makes the abdominal wall prominent. The growth seems tense, is slightly lobulated and, although sharply outlined, gradually blends with the liver substance. The edge of the liver forms a prominent ridge along the lower margin of the growth and the gall bladder is seen lying against the under and inner surface of the tumor. It is readily seen that when the cystic duct was severed the greater part of this liver could be rolled out of the abdomen. For a cross section of the hepatic growth see Figure 7.

ent, the surgeon will be justified in concluding that the hepatic growth is a secondary cancerous nodule.

As a matter of fact, it is rarely possible to be able to make more than a tentative diagnosis, until the growth has been removed. In our case the left kidney had been removed nearly eighteen months previously for carcinoma. The symptoms accompanying the hepatic tumor were quite applicable to a case of liver abscess,

the tumor was an abscess or a metastatic carcinomatous nodule. As the remaining kidney was already impaired, it was thought that, even should the swelling prove to be an abscess, an attempt at drainage would be associated with considerable danger of an acute peritonitis, and that even if this complication were avoided, the continual discharge from the abscess wall would militate against the patient's recovery. On the other hand, in

case the growth were malignant, it was evident that the only hope of success lay in its complete removal. Even after the tumor had been sent to the laboratory a provisional diagnosis of abscess of the liver was made after a thorough macroscopic examination, and the histologic revelation of adenocarcinoma came as a great surprise.

EXPERIMENTAL RESECTION OF PORTIONS OF THE LIVER.

The exhaustive articles of Langenbuch² and of Terrier and Auvray¹ give in detail the interesting animal experiments of Glück, Tizzoni, Griffini, Podvisotzky, Ponfick, Tillmanns and Ceccherelli. Ponfick⁵ found that it was possible to remove three-fourths of a dog's liver without permanent injury to the animal, and that in thirty days the organ had regained its normal size. Regeneration was partly due to hypertrophy, but mainly to multiplication of the remaining liver cells. No new lobules were found. Judging from analogy, the surgeon would naturally infer that in man large portions of the liver might be removed and the patient recover;

Fig. 6.—Method of controlling hepatic hemorrhage. The cut surface represented in the upper sketch measures 16x8 cm. Mattress sutures were applied about 1 cm. from the line of section. Each overlapped the other so that every particle of liver substance was controlled. Pagenstecher sutures were used, reinforced by catgut, where necessary. The large vessels were readily picked up with forceps and tied freehand. After all the sutures had been tied and lifted up, *a* and *b* tended to come together. Beginning at *c*, corresponding opposite sutures were drawn across and tied to each other. The approximation was good, but not quite so accurate as one would infer from the sketch.

and the reported cases prove that the suggestion is well founded, as in some instances large areas have been resected and the patients have regained perfect health.

INDICATIONS FOR RESECTION OF A PORTION OF THE LIVER.

It is important to determine (1) whether the growth be malignant or not; and (2) if malignant, whether it is primary or secondary. In the case of metastatic neoplasms, as a rule, operation is contraindicated, inasmuch as they are usually multiple. Primary carcin-

oma or sarcoma of the liver is rare. Should the hepatic growth be primary, the glands along the vessels and at the base of the liver should be carefully palpated to ascertain whether they have become involved. This point was first insisted on by Tuffier.⁶ In cases of lymphadenoma, with secondary nodules in the liver, operation is not advisable. In all cases of liver growths the following points should be carefully considered: (a) Whether the growths be single or multiple; (b) situation; (c) size; (d) whether they are pedunculated or not.

If the growths are multiple and scattered over a wide area, removal is difficult or impossible. On the other hand, provided that they are near the lower edge of the organ, resection is feasible. In fact, in the majority of cases the situation is all important.

The size is of little importance when the growth is pedunculated and situated near the lower margin of the liver. In the case of a large tumor, however, located near the middle of the organ, removal is exceedingly difficult or impossible. If it is located near the edge, amputation is not difficult, but if it is near the middle of the lobe, a very large portion of the liver must be resected, or a cavity is left which may be the source of an alarming or even uncontrollable hemorrhage. When the growth lies near the portal vein or hepatic arteries, operation is contraindicated, since large branches of these vessels are liable to be injured with a resulting necrosis of correspondingly large areas of liver substance and the death of the patient. If the growth be pedunculated, its removal is easier, inasmuch as the liver substance forming the pedicle is much tougher than other portions of the organ. This toughening is due to partial atrophy of liver cells.

METHOD OF PROCEDURE.

After having decided that the growth is to be removed, free exposure of the liver is absolutely necessary. The abdominal incision is lengthened, and may be curved to the right or left as desired. If the liver is freely movable, the greater part can be brought out of the abdomen; but if the organ is fixed, it will be necessary to loosen it. This is accomplished by severing the suspensory and round ligaments and, if necessary, to a limited extent the ligamentous attachments of the diaphragm. Care must be exercised to avoid injury to the pleural or pericardial cavity, and in case of such an accident the opening should be immediately closed with catgut sutures. The mobility of the liver may also be considerably increased by cutting through the cystic duct, as was done in this case, after which two-thirds of the liver was easily brought outside the abdomen. If the liberation of the liver obtained in this manner is still insufficient, portions of the lower ribs may be cut through and turned back. This free exposure of the liver will be found of the greatest comfort to the operator during the subsequent steps of the operation; for not only is he able to control the blood supply much more thoroughly, but he is also working outside the body instead of deep down in the cavity.

TEMPORARY MECHANICAL CONSTRICTION.

Prior to cutting into the liver, a constricting band, if possible, should be thrown around the organ, at least an inch or more from the line of section. Many surgeons have employed strong rubber tubing; or a long piece of gauze twisted in the form of a rope, as used

5. Ponfick: *Lancet*, 1890, vol. 1, 881.

6. Tuffier: *Bull. et mém. de la Soc. de Chir. de Paris*, 1897, n. s., xxiii, p. 70.

in this case, will answer equally well. Even after this has been drawn tight and clamped with forceps, it must be carefully watched throughout the entire operation, as it is liable to slip from the slanting liver substance and thus fail to prevent a free hemorrhage.

THE INCISION INTO THE LIVER SUBSTANCE.

When we remember the great number of blood vessels that are present one can readily understand the dangers connected with the removal of a large segment of liver substance. In Langenbuch's most graphic account of his experiences we are told that during an operation the hemorrhage was so sudden and severe that the abdomen was in an instant full to overflowing with blood and the field of operation was obliterated. Any one who has read his account will certainly never fail to

the tumor out with the thumb nail (Keen). Each method has its advantages and its drawbacks. The knife gives a clean-cut surface, but many vessels are opened and bleed freely. The cautery leaves a charred surface, and vessels that are temporarily controlled may bleed later. The operator hardly feels safe in trusting solely to small plugs of clotted blood to control the small and medium-sized vessels, when he knows that the escape of one of these may mean fatal hemorrhage. On the other hand, nothing checks capillary hemorrhage better than the cautery at a dull red heat. Blunt scissors, the curette and blunt dissection with the finger leave a much rougher base, but injure far fewer vessels.

THE CONTROL OF HEMORRHAGE.

Probably the safest plan is to control the bleeding

Fig. 7.—Secondary carcinoma of the liver (natural size). As the specimen was very soft it was impossible to cut it before hardening. Encircling the entire growth is a definite capsule varying from 1 to 3 mm. in thickness. *a* is the sharp edge of liver substance that passed over the lower surface of the growth. This growth bears much resemblance to the parent kidney tumor (Fig. 4), and in its central portion has undergone complete necrosis. The cut surface of the liver indicates the line of amputation.

have plenty of assistants, and every conceivable requisite is at hand to meet any similar emergency. Now-a-days, fortunately, the picture is not so disheartening, thanks to the improvements in technic introduced by Keen, Kousnietzoff⁷ and Auvray. I shall not attempt to give in detail the various methods employed, but shall only sketch very briefly the procedures that have apparently yielded the best results.

The liver substance may be severed by: (1) The knife; (2) the cautery (at a dull red heat); (3) the blunt scissors or curette; (4) blunt dissection, peeling

before making any incision into the liver or to tie off different segments as they are severed. Langenbuch has pointed out that the liver substance often shows pathologic changes when tumors are present. Thus there may be a hyperemia associated with an increased friability of the organ rendering the application of sutures difficult. On the other hand, there may be an atrophy of the liver cells accompanied by an increase of connective tissue, which renders the tying of ligatures a relatively easy procedure.

Needles.—One of the chief sources of hemorrhage in liver operations has been from the needle holes; in fact, in some cases, the free oozing from these sources has been uncontrollable. Thanks to the ingenuity of

7. Kousnietzoff (Kuznetsoff): *Laitop. russk. Chir.*, S. Peterb., 1901, 6, 888-931. Mikulicz wrote the preface to this article and mentioned the number of cases operated on by him.

Kousnietzoff, we now possess a needle that renders the application of sutures almost or entirely bloodless (Fig. 1). This needle may be straight or curved; it is flat on the sides, and rounded at both ends. It is gently and slowly forced through the liver substance, and, being blunt, pushes the blood vessels to one or the other side. This needle has yielded admirable results in dogs, and Mikulicz, who has employed it six times in the human liver, says of it, with some enthusiasm: "To my mind Kousnietzoff's needles represent the egg of Columbus in the technic of liver surgery."

Cautery.—Keen slowly and cautiously employed the cautery at a dull red heat, and took twenty minutes to cut off the desired segment. He was usually able to see the larger vessels before they were opened. These he tied, both proximally and distally, and then burned through them. If a large vessel was opened before being tied, he plugged the opening with his left index finger, while with the right hand he passed a ligature threaded on a curved needle carefully through the surrounding liver substance; the ligature was then tied slowly by an assistant. In this manner he was able to remove the growth with little difficulty. Keen insists on the use of dull red heat, and points out that, if the tissue is cut through too rapidly, a sufficient eschar is not formed to check the oozing. In some of his work he combined blunt dissection with the finger with the use of the cautery.

Sutures.—Kousnietzoff has devised a rather complicated continuous suture (Fig. 2), which practically resolves itself into a series of mattress sutures. Auvray has advised a continuous chain suture which he has found to work well (Fig. 3), its object being to distribute more equally the strain on the liver substance and so render it less liable to tear. The sutures of Kousnietzoff and Auvray would seem, from the illustrations, to be open to the objection that hemorrhage might occur from the spaces between adjoining sutures. In our case simple mattress sutures were introduced with the blunt needle (Fig. 6), each suture overlapping the other so that no area was left unprotected. After the application of three or four sutures about 1 cm. from the line of section, the liver was severed over a corresponding area, an assistant in the meantime holding the placed sutures taut. The tumor was grasped firmly in order to prevent oozing.⁸

The sutures were now slowly and snugly tied; others were then applied in a similar manner, and the entire liver growth was readily removed in about ten minutes. The mattress sutures were tied after the tissues had been cut, instead of before, because the liver was stretched near the tumor, and there was danger that after tension was relieved they would be too loose to thoroughly control the vessels. Five or six large vessels were encountered. These were readily picked up with forceps and tied.

Suture Material.—In many of the reported cases catgut was used exclusively. In this case fine Pagenstecher thread, reinforced with catgut, was employed. Catgut is, of course, the material generally preferred, but where there are such large vessels and so much ten-

sion, sutures of silk or linen thread are, on the whole, more reliable; although the silk will gradually work its way out.

Treatment of the Stump.—After controlling all bleeding, it remains to deal with the raw area. Where the cautery has been used, one can not expect to bring charred surfaces together and get union. Again, only in very exceptional cases would it be possible to remove a large tumor and have such a satisfactory folding as in this case. As a rule, the cut surface must be left as it is. Some surgeons advocate stitching the omentum over it. Others close the abdomen without drainage. From a careful review of the cases I gather the impression that it is better to drain. Where this is done, bile is often found escaping through the wound from the cut surface of the liver for several days, and if the abdomen has been closed without drainage, this must of necessity flow into the abdomen. Keen has laid stress on this point, but says that the danger from absorption "seems to be more theoretical than practical." In this case an iodoform drain was carried to the under surface of the liver and another to the upper surface along the line of the sutures. Pfannenstiel⁹ has called attention to the possible danger of using iodoform gauze where large areas of raw liver substance are left, and prefers sterile non-medicated gauze for this purpose.

COMPLICATIONS THAT MAY ARISE DURING AN OPERATION.

The operation is sometimes beset with many difficulties. Where the tissue is very friable the sutures may cut through the liver substance. To overcome this difficulty Ceccherelli and Bianchi used decalcified bone plates over the upper and under surfaces and sutured through them. In some cases, after removal of a nodule from the center of the liver, when it was impossible to check bleeding by sutures, the cavity was packed with gauze, held firmly in place by sutures passed through the liver substance.

In other cases, where it was impossible to check oozing, a rubber ligature was firmly secured around the liver, slipping being prevented by means of two hat-pins passed through the organ at right angles to each other. The edge of the liver was then sewed to the peritoneum of the abdominal wound. In some of these cases the growth was cut away and the stump left to granulate up. In other instances the growth was left to slough away, the constricting band being tightened from time to time.

IMMEDIATE AFTER-RESULTS.

Hemorrhage is the most serious complication. It may take place after a few hours or at the end of a day or two. The employment of the suturing method of Kousnietzoff or Auvray or the overlapping mattress suture will undoubtedly render secondary bleeding far less frequent. Where hemorrhage occurs, however, the abdomen should be opened at once, for the patient is not always exsanguinated. If the bleeding points can not be controlled a rubber ligature should be applied and prevented from slipping by means of hat-pins. The stump can then be treated extraperitoneally, the edge of the liver being fastened to the peritoneum.

MORTALITY.

Terrier and Auvray collected 38 cases operated on for the following diseases: Sarcoma, 4; carcinoma (considered primary at the time of operation), 7; sec-

8. In bisecting the uterus there is always free hemorrhage, especially in the case of a large myomatous organ. Dr. Kelly has taught that strong traction on each half almost completely checks oozing. This principle was applied to the distal segment of the liver with very satisfactory results. It must be said, however, that the liver substance removed with the tumor, in this case, was relatively small.

9. Pfannenstiel: Med. Centr. Ztg., Berl., 1898, lxxvii, 177-179.

CASES OF RESECTION OF PORTIONS OF THE LIVER REPORTED SINCE 1899.

Reporter and Reference.	Sex.	Age.	Duration, Nature and Size.	Method of Removal.	Treatment of Liver Stump.	Result.	REMARKS.
Carl Beck: Chicago Med. Recorder, 1901, xxi, 383-386.	Male...	20	Followed kick 5 years before. "Angio-adenoma," the size of man's head occupied the entire left lobe.	Controlled by rubber catheter, held by two hat pins through stump. Half the tumor removed, half left to slough off.	Extra-peritoneal...	Recovery...	Largely an angioma, but with certain adenomatous areas.
Harrison Cripps: Brit. Med. Jour., Lond., 1903, ii, 18.	Male...	60	Pain after heavy lifting 11 years before. Angioma the size of a child's head.	Slightly adh. to abd. wall; incised; escape of 50-60 oz. blood. Tumor enucleated by dissecting; Volkmann's spoon. Bleeding controlled by sponge pressure.	Abd. closed without drainage.	Recovery...	Angioma partly calcified.
Dahlgren: Nord. Med. Ark., Stockholm, 1902, 3, f. ii, No. 14, 1-24.	Female	33	Over a year. Angioma from left lobe, larger than a kidney—pedicle 6 cm. broad; 1.5-2 cm. thick.	Bleeding controlled by continuous catgut sutures. Tumor removed by cautery.	Omentum attached to stump.	Recovery...	Gives good résumé of the literature.
Macnaughton-Jones: Brit. Gynec. Jour., Lond., 1901-2, xvii, 40-43.	Female	27	One year. Angioma. 3½ x 2¼ inches at liver. Irregular spaces lined by endothelium and filled with blood.	Tied by numerous ligatures; little bleeding.	Pedicle packed above and below with iodoform gauze.	Died 18 hours after operation.	Much shock at operation.
Ginlio, Filippini: Policlin., Roma, 1901, viii, Sez. Chir. 222-240.	Female	32	Tumor of left lobe the size of an adult head, 20x15x5.6 cm.	Rubber ligature and sutures.	Attached to abdominal wall.	Recovery...	Angioma.
Ginlio, Filippini: Same reference.	Male...	62	Cancer of the liver involving the entire left lobe.	Rubber ligature...	Attached to abdominal wall and allowed to slough off.	Recovery...	Primary cancer of the liver.
Duret: Ass. Franc. de Chir. Proc. Verb. Par. 1898, xii, 363-371.	Female	46	Six months. Carcinoma; 16 cm. in its greatest diameter.	Cut away as a wedge. Vessels controlled by a few sutures; raw surfaces approximated by continuous catgut.	Iodoform gauze drain; exerting pressure.	Recovery...	Gall-bladder contained calculi.
Jonnesco: Bull. et Mém. Soc. de Chir. de Bucarest, 1902-3, v, 104.	Female	..	Size of a small orange, adherent to the transverse colon.	Tumor resected; a modified d'Auvray ligature employed.	Closed without drainage.	Recovery...	Thought to be syphilitic. Wound opened on 3d day to let out some blood.
Haubold: Annals of Surgery, 1904, xxxix, 240.	Female	26	Eight months. Gumma, the size of a large orange.	Removed by blunt dissection with scissors.	Hot saline. Gauze checked oozing—surfaces approximated by deep catgut sutures. Drainage.	Recovery...	Growth first thought to be malignant.
Léonté: Bull. et Mém. Soc. de Chir. de Bucarest, 1900-01, iii, 161-164.	Female	50	Adenocarcinoma in the right lobe, the size of a lemon. Weight, 843 grams.	Sutures placed; tumor removed.		Died in 24 hours. Anuria from renal disease.	No bleeding.
Thompson: Tr. Texas Med. Assn., Austin, 1899, xxxi, 216-229.	Female	30	Gumma, size of a small orange; occupied parts of both lobes.	Tumor removed with scissors; transfixed by two needles at right angles. Appl. of rubber.	Fastened to abdominal wall. Infection.	Recovery...	Histologic examination showed gumma.
Nicholson: Interstate M. J., St. Louis, 1902, ix, 427-430.	Male...	39	3¼ inches broad; projected 2 inches from surface of liver.	Cut out with scissors; vessel openings closed with chromicized catgut.	Packed with gauze; partially removed 2d day, some bleeding. Removed 4th day; edge of abd'l wound repaired and closed.	Recovery.	
Pfannenstiel: Allg. Med. Centr. Ztg., Berl., 1898, lxxvii, 177-179.	Female	..	Growth noticed years before. About 12 pounds. Abd. before operation, 116 cm. in circumf. (considerably more than that of a pregnant woman at term). Tumor pale white, "alabaster-looking," fluctuating—sprang from under surface of left lobe. On section composed of large and small blood spaces—some spaces as large as a walnut, others as big as an apple. Micro. exam. showed growth to be an angioma.	A furrow made in liver substance both anteriorly and posteriorly at base of growth. Tumor then shelled out partly by cautery, partly by blunt dissection—bleeding vessels controlled by catgut passed with a Deschamps needle; in part by pressure, in part by strong traction on tumor. Some broad clamps applied, stump about 15 cm. broad thus produced—tied with about 20 catgut sutures; tumor cut away with cautery.	Large funnel-shaped stump left, attached in upper angle of abdomen wound. Packed with sterile, non-impregnated gauze. Length of operation 1½ hours. Pulse at end, 120-140. Discharge at first, bloody; later, bile.	Recovery...	Resembled an ovarian cyst, but on vag. exam. uterus and appendages were clearly outlined. The continual traction of tumor had probably rendered the liver substance of pedicle firmer than normal. Very little urine at first, probably from great loss of blood; salt sol. given freely.
Maresch (Hochenegg) Ztschr.: F. Heilk., Wien u., Leipz., 1903, xxiv, 39-50.	Female	5	Three years; at first slowly, later rapidly. Abd. greatly distended. Lymphangioma 20x18 cm.	Punctured, supposedly an ovarian cyst; pedicle about 7.5 x 2.5 cm. Collar of peritoneum drawn back. Tumor then dissected out bluntly with forceps and elevators. Tremendous bleeding from torn vein. Abd. full. Firm pressure made. Hem. controlled with catgut sutures and needle.	Incision 6 cm. long made toward right abdomen wall, 3 cm. below the xiphoid. Pedicle of tumor sewed into this transverse incision. Original incision then closed.	Recovery; discharged on 21st day.	Supposed to be an ovarian cyst.
Terrier et Auvray: Revue de Chir., 1898, xviii, 719.	...	50	Cancer the size of a closed fist, probably secondary to growth in bowel. Adherent.	Chain sutures of Auvray. One vessel tied free-hand.	Packed with gauze...	Recovery...	Fistula still remaining; silk sutures came out gradually.
Barkoff Russ. Chir. Arch., St. Peterb., 1903, xix, 70-72.	Female	40	Ten years; the size of a hen's egg; oval and yellow.	Cut out elliptical depression; approximated with catgut. No hemorrhage.	Closed without drainage.	Temporary recovery.	Echinococcus.
Cullen, Thomas S. . .	Female	30	Recognized less than a week before operation. Adenocarcinoma of liver, secondary to primary growth in left kidney. Growth size of child's head. (Fig. 5.)	Cut away with knife after application of mattress sutures. Raw surface approximated. (See Fig. 7 in text.)	Iodoform gauze drainage.	Patient in good health 8 mos. after operation.	Evidence of return in scar where left kidney was removed nearly 2 years ago.

ondary carcinoma, 1; adenoma, 6; angioma, 4; tumors of an indefinite character, 3; gummata, 9; biliary cysts (non-parasitic), 4.

Of the patients, 32 recovered and 6 died. The deaths were due to hemorrhage, shock or sepsis. Keen has collected 74 cases. Some of these are included in Terrier and Auvray's lists. Sixty-three of the patients recovered and 11 died—a mortality of 14.9 per cent.

The causes of death were: Shock, hemorrhage and exhaustion, in 8 cases; septicemia, in 2 cases; pulmonary embolism, in 1 case.

In this group of Keen's are three of his own cases; all three patients recovered. As mentioned elsewhere, Mikulicz has had six cases with equally happy results.

Patient.—On Jan. 6, 1903, I saw Miss T., in consultation with Dr. E. R. Trippe. The patient was 30 years of age and of very slight build. For six months she had felt a small lump in the left side, but had hesitated to speak of it until a week previously. It was first noticeable in the region of the left kidney and at first had increased in size gradually, but later at a much more rapid rate.

Examination.—On examination the left renal region was found to be filled by a tumor which extended upward under the ribs, and downward almost to the pelvic brim. It was slightly movable.

Operation.—On January 8 an incision was made over the most prominent part of the tumor. This incision extended from the quadratus lumborum behind, obliquely downward almost to the median line just above the pubes. The tumor was everywhere firmly adherent and was liberated with a great deal of difficulty. Only at one point was there any escape of its contents. Here the capsule had broken slightly, allowing the escape of a small amount of friable, carcinomatous-looking material. After liberating all adhesions and tying every bleeding point with catgut the renal vessels were exposed and tied first *en masse*, and then singly. The abdominal cavity was opened in several places as the tumor was intimately blended with the peritoneum. We were, however, able to protect the abdomen from infection. The muscles were brought together with silver wire, the skin with silkworm gut. The patient, although much shocked, gradually recovered. Her temperature rose on one occasion to a little above 100 F. She made an uninterrupted recovery and left the hospital at the end of the fifth week.

Result.—On March 9, 1903, I received a letter from the patient saying she was very much improved; the wound had healed entirely and she was feeling much stronger. I saw her from time to time and she showed continued improvement and regained perfect strength.

Subsequent History.—On June 2, 1904, I received a message from Dr. Trippe saying that the patient was not so well and that her temperature was 102.5. Five days previous to this the patient had noticed a small lump just below the free edge of the ribs on the right side. At first it gave her no concern, but apparently doubled¹⁰ in size within two days and during the following two days again showed marked enlargement.

Examination.—When I saw her it occupied the region between the edge of the ribs and the anterior superior spine. It was round, tense and somewhat movable. We were very careful in our examination, fearing that it might be a friable secondary growth following the removal of the large carcinomatous kidney in January, 1903. Her pulse was rapid; her temperature was 102.5 F.

Operation.—As she was losing ground rapidly an immediate operation was decided on. An incision having been made over the tumor, it was found to be perfectly smooth and free

from adhesions. The incision was continued upward to the free margin of the ribs (Fig. 5) and downward almost to the anterior superior spine. The growth proved to spring from the liver; it was larger than a child's head and was almost globular in form. The edge of the liver appeared as a slightly elevated ridge, forming a crescent over the center of the anterior surface of the growth. Attached to the inner side of the growth was the gall bladder. I carefully examined the liver, but failed to find any secondary growths and also was unable to detect any evidence of metastases in the abdominal organs or any thickening in the renal region on the left side. The right kidney was about half as large again as usual, and was perfectly normal, considering the absence of the other. The urine, however, contained some pus and a little blood. On careful examination it was impossible to determine whether we were dealing with a cystic new growth or an abscess formation. In the former case we were at a loss to tell why there had been as yet no adhesions and why the surrounding liver substance was so firm. On the other hand, it was difficult to account for the elevation of temperature if no abscess was present. After some hesitation we decided to remove the growth together with a small portion of healthy liver substance. The gall bladder was exposed, the cystic duct clamped and cut and the proximal portion of the duct tied off. The lower half of the liver could then be readily drawn out through the abdominal incision. It was everywhere walled off from the abdominal contents by gauze, and a gauze rope was securely fastened around the liver about an inch above the point at which the cut was to be made. A cautery was called for, but—fortunately, as after results showed—the instrument would not work. I then took a long, blunt needle and, starting from left to right, passed mattress sutures through the entire substance of the liver. These were about 1 cm. apart and each one overlapped the other for about 2 or 3 mm. The greater number of sutures consisted of Pagenstecher thread, the supplementary ones being of catgut. After placing two or three mattress sutures I commenced to cut, the tumor at the same time being held taut. The sutures were then tied. This process of applying sutures and cutting and then tying was continued until the entire growth had been removed. Five or six large blood vessels were encountered, which spurted freely. These were picked up with artery forceps and tied freehand with fine Pagenstecher thread. The entire growth was removed without the loss of more than two drams of blood. After the sutures had been tied there was absolutely no bleeding from the surface. After removal of the growth there remained a raw surface fully 16 cm. in length and 6 cm. in breadth. On making traction on the sutures I found that the raw surface could be rolled in on itself so that the two halves formed flaps. These were brought together, until little or no raw substance remained. The ends of the sutures that had already been tied were utilized to bring the opposite sides together. Iodoform gauze was placed beneath the edge of the liver and also on its surface. Although the patient was apparently in a desperate condition when she went on the table, she stood the operation well, and when she was taken from the operating room her pulse was only 104. On the next day her temperature dropped to 100 F. The general condition was excellent and the pulse 84. To look at her one would hardly have realized that any operation had been performed. She made a rapid recovery and left the hospital in about four and a half weeks.

Result.—I saw her on Aug. 4, 1904. She was looking very well and had regained her color; the wound had practically healed. She had little or no pain. On the left side there were areas of suspicious thickening in the old renal scar.

Note of Sept. 7, 1904: On examination some of the liver sutures are seen working their way through a small sinus. They are giving no trouble, however. The patient is up and around, apparently in good health. On the left side along the linear incision, where the kidney growth was removed a year and a half ago, several nodules are felt. These are rapidly increasing in size. One of them is fully 6 cm. in diameter and

10. The tumor had evidently existed for a considerable length of time, but was not recognized until it was suddenly forced from beneath the ribs on account of its increasing dimensions.

has already become slightly attached to the skin. It is bluish in appearance.*

PATHOLOGIC REPORT OF KIDNEY.

Miss T. Pathologic No. 6364. The specimen, roughly speaking, is kidney-shaped, irregular in outline and covered by many adhesions. It measures 20x13x12 cm. Attached to the lower end is the ureter, which is about 4 mm. in diameter. On palpation the tumor seems to be cystic. This condition is more noticeable at the upper pole. Toward the lower end the growth is firmer. On section little or no renal tissue is demonstrable. The tumor has a capsule varying from 2 to 5 mm. in thickness. Passing off from this capsule into the tumor are many connective-tissue partitions which divide the growth into large and small nodules. The greater part of the tumor is soft, and consists of a spongy-looking material resembling carcinoma. The more solid portions of the growth are homogeneous. At numerous points there has been disintegration of tissue. The growth, as a whole, is yellowish-white in appearance, but where degeneration has taken place the tissue is of a bluish tinge or is distinctly yellow.

Histologic Examination.—The capsule is composed of fibrous tissue, sharply outlined from the tumor, which is made up of a glandular growth. The gland type is particularly well marked. There are long stems carrying delicate blood vessels and covered by a single layer of epithelium. Passing off everywhere from these stems are secondary stems, likewise covered by one layer of epithelium. The epithelial cells are low cylindrical or cuboidal in type. Their nuclei are round or oval and stain fairly deeply. In many places a definite glandular arrangement is visible, but the tendency toward the papillary formation is everywhere in evidence. Here and there nuclear figures are visible. The specimen was hardened *en masse*; consequently one would not expect to find many cells undergoing division. In the older portions of the growth the cells have so multiplied that they form a homogeneous mass. At many points coagulation necrosis has taken place and the cells do not stain at all. Such an area, as a rule, is entirely devoid of infiltration, but here and there are aggregations of polymorphonuclear leucocytes. At some points are masses of bright orange pigment, evidently a result of old hemorrhages. Such masses have a fuzzy appearance and closely resemble the outer covering of a chestnut. The specimen is undoubtedly from an adenocarcinoma. The cells of this growth bear a striking resemblance to those of the collecting tubules of the kidney. Sections of the ureter just below the pelvis of the kidney show that it is perfectly normal.

PATHOLOGIC REPORT OF LIVER TUMOR.

The specimen consists of a tumor removed from the lower portion of the liver. Attached to this are the gall bladder and a small amount of normal liver substance. The tumor is 16 cm. long, 14 cm. broad and 13 cm. in its anteroposterior diameters. It is slightly lobulated, has a smooth surface and appears gradually to blend with the healthy liver substance. Passing over the middle of the lower surface of the tumor is a thin strip of liver substance representing all that remains of the lower edge of the liver. The raw surface corresponding to the liver attachment is 16 cm. in length, 6 cm. in breadth. The liver tissue is apparently normal. The growth on palpation seems to be filled with thick fluid contents and strongly suggests an abscess.

The specimen was hardened in alcohol before being opened. It is surrounded by a capsule 1 to 2 mm. thick. The interior is composed of a very soft homogeneous substance. In the outlying portions, however, it is rather spongy. In the hardened specimen the growth presents a mottled appearance; in places it is reddish-blue, at other points yellowish-white in color. Only a small amount of healthy liver substance is present. The gall bladder is firmly attached to the upper and inner side

of the growth and measures 7 cm. in length, 3 cm. in breadth.

To describe the histologic appearance of this tumor would be to repeat what has been said of the histologic appearance of the kidney. It is identical in every particular. In other words, it is an adenocarcinoma secondary to the primary growth in the kidney. It also contains numerous areas of degeneration.

In 1892 T. S. Westcott tabulated 20 cases of partial resection of the liver for Dr. Keen. A second article by Keen, in 1897, contained a tabulation of Cases 21 to 59 by G. W. Spencer. For Keen's article, published in 1899, H. H. Cushing and M. L. Downs have tabulated all additional cases to that date. The accompanying tabulation includes the cases that I have been able to gather from the literature since 1899; for convenience the lines laid down by Keen have been followed.

This case is unusual for the apparently rapid development of the hepatic growth, the symptoms being of less than a week's duration. It is further unusual to find a single metastatic nodule in the liver following a primary growth in the kidney.

Considering the fact that the patient had only one kidney, that the urine contained blood and pus, that the temperature was 102.5 F., and the physical condition poor, I advised letting the growth alone. Dr. Trippe, who was not only the patient's physician, but also a near relative, on the grounds that her condition was rapidly growing worse and that she could not last many days, insisted on its removal. That he was right is clearly shown by the temporary recovery of the patient, and I freely confess that any credit for the operation is due entirely to him. Certainly no case that I could find in the literature offered less prospects of even a temporary success, but Dr. Keen, when reporting his last case, said: "After my experience with these three cases I should hardly hesitate to attack almost any hepatic tumor without regard to its size."

RECLINATION OF THE LENS:

UNDER CERTAIN CONDITIONS A JUSTIFIABLE OPERATION.*

F. T. ROGERS, M.D.

PROVIDENCE, R. I.

A study of the literature of couching reveals the fact that earlier writers made a sharp distinction between depression of the lens and reclination, in the one the lens being pushed bodily downward while in reclination an attempt is made to turn it on its axis so that ultimately its anterior surface is uppermost. In reality, however, an attempt at one procedure is likely to result in the other, so little is the course of the lens under our control, and no distinction will be made in this paper, although as late as 1836 Argyll Robertson¹ said he had never seen the lens rise after reclination, although he had frequently seen it after depression, and that the former, if properly performed, is rarely productive of much inflammatory action.

Depression is the oldest known operative procedure for cataract,² and either by the insertion of the needle through the cornea, keratonyxis, or the sclera, scleronyxis, until the middle of the eighteenth century held complete sway, when it was superseded by extraction, and if since then it has been held in slight respect it is

*Feb. 24, 1905: Dr. Trippe informs me that from time to time a few stitches have come away from the liver, but have occasioned absolutely no inconvenience. The nodules in the cicatrix in the side where the kidney was removed are somewhat larger, but they do not seem to cause her any inconvenience. "Her general condition seems to be very good."

* Read in the Section on Ophthalmology of the American Medical Association, at the Fifty-fifth Annual Session, June, 1904.

1. J. Argyll Robertson: *Edinburg M. & S. J.*, vol. xlvii.

2. Czermak: *Augenärztlichen Operationen*, Wein, 1904, Heft. 13-14, pp. 823-827.