The correction for refraction has been applied in those cases where the difference of N.P.D. made it sensible. — The apparent place of Lac. 1685 has been taken from differential measures with B.A.C. 1511 which was observed at Williamstown. B.A.C. 1564 and 1644 were both observed at Melbourne in 1880. The approximate places of the comparison-stars which are not to be found in any catalogue in my possession are:

Star	Mag.	R. A.	N. P. D.
<i>a</i>	$7\frac{1}{2}$	$5^h 4^m 49^s$	$124^0 41'$
<i>b</i>	$7\frac{1}{2}$	5 8 9	124 37
<i>c</i>	7	5 4 12	123 54
<i>d</i>	$6\frac{1}{2}$	5 5 22	123 40
<i>e</i>	8	5 2 13	116 57

Melbourne 1881, June 6.

Rob. F. Ellery.

Elemente des Cometen *b* 1881. Von Herrn S. C. Chandler jr.

With the permission of the Director, Prof. E. C. Pickering, I communicate the following elements of Comet (*b*) 1881:

$$\begin{aligned}
 T &= 1881 \text{ June } 16.1907 \text{ Washington m. T.} \\
 \pi &= 265^0 7' 7'' \\
 \Omega &= 270 58.8 \\
 i &= 63 30.9 \\
 \log q &= 9.86568 \\
 \text{Motion Direct.} \\
 \text{Middle Place} \\
 \cos \beta \Delta\lambda &= + 0' 1 \\
 \Delta\beta &= - 0.5
 \end{aligned}
 \quad 1881.0$$

These elements were derived by Mr. O. C. Wendell and myself from observations made at this Observatory on June 23^d and June 25th, joined with the approximate position at Rio Janeiro, by Cruls, on May 29th ($\alpha = 5^h 2^m$ $\delta = - 31^0 15'$) assuming the local mean time of observation to be 6 hours. This is the only earlier position accessible to us.

Harvard College Observatory, June 27th, 1881,
Cambridge U. S. A.

S. C. Chandler jr.

Beobachtungen und Elemente des Cometen *b* 1881.

Die Elemente sind hergeleitet aus folgenden, von Herrn Romberg am Meridiankreise der Pulkowaer Sternwarte angestellten Beobachtungen:

mittl. Berl. Zt.	α	δ
1881 Juni 25.42928	$85^0 36' 48''$	+ 53 1 39
» 27.43066	88 4 55	+ 59 57 37
» 29.43376	91 10 29	+ 65 35 15

Dieselben sind auf das mittlere Aequinoctium von 1881.0 reducirt, jedoch noch nicht für Aberration und Parallaxe corrigirt worden. —

Pulkowa 1881 Juli 2.

Die gleichzeitig übersandte Ephemeride stimmt nahe mit der des Herrn Dr. Oppenheim.

$$\begin{aligned}
 T &= 1881 \text{ Juni } 16.5035 \text{ m. Berl. Zt.} \\
 \pi &= 265^0 15' 42'' \\
 \Omega &= 270 58 13 \\
 i &= 63 29 20 \\
 \log q &= 9.86572
 \end{aligned}
 \quad \text{mittl. Aeq. 1881.0.}$$

Darstellung der mittl. Beobachtung: (B.—R.)

$$\begin{aligned}
 \Delta\lambda \cos \beta &= - 2'' \\
 \Delta\beta &= + 0
 \end{aligned}$$

Th. Witttram.

Kr.

Observations méridiennes de la Comète *b* 1881 à Madrid.

1881 Juillet	3	$11^h 51^m 44^s$	t. m. d. Madrid.	α app. = $6^h 40^m 19^s$	00	δ app. = + $73^0 46' 28''$	8	Observateur:	Ventosa.
»	4	11 59 4	»	»	6 51 36.7	»	75 13 48.0	»	La Cruz.
»	5	12 7 33	»	»	7 4 3.3	»	76 30 25.1	»	Ventosa.
»	6	12 17 13	»	»	7 17 41.5	»	77 37 23.7	»	»
»	7	12 28 6	»	»	7 32 32.7	»	78 35 24.7	»	»
»	8	12 40 9	»	»	7 48 34.4	»	79 25 14.1	»	»
»	12	13 37 24	»	»	9 1 45.5	»	81 34 7.0 (?)	»	Jimenez
»	13	13 52 49	»	»	9 21 9.1	»	81 51 6.4	»	»
»	15	14 22 51	»	»	9 59 9.8	»	82 11 47.8	»	»

Les déclinaisons n'ont pas été corrigées de la parallaxe. Celle du 12 est un peu douteuse. Le mauvais temps n'a pas permis de faire d'autres observations.

Madrid, le 16 Juillet 1881.

Pour M. le Directeur:

V. Ventosa.